Alptekin Aydin

ID UD83722HPS9294045

PhD Proposal: Assessing the Impact of tDCS Treatment Over the Dorsolateral Prefrontal Cortex (DLPFC) on Social Communication in Children with ASD: A Survey of Family Experiences

A Final Thesis Presented to

The Academic Department

Of the School of Social and Human Studies

In Partial Fulfillment of the Requirements

For the Degree of Doctorate in Clinical Psychology

ATLANTIC INTERNATIONAL UNIVERSITY HONOLULU

HAWAII SUMMER 2024

Acknowledgments

This dissertation marks the conclusion of an amazing journey that would not have been possible without the constant support, direction, and inspiration from several people and organisations. My sincere gratitude goes out to everyone who has helped with this project. The academic department staff and faculty in the School of Social and Human Studies have my sincere gratitude. Completing this dissertation has been made possible by your commitment to fostering an engaging and encouraging learning environment.

A special thanks goes out to my dissertation committee members for their helpful criticism, wise recommendations, and unwavering support. I also want to sincerely thank to the children and families who kindly offered their time and shared their stories, this research would not have been possible. Your willingness to participate in this research has been much appreciated, and your contributions have been invaluable to it.

My greatest source of strength has come from my spouse/partner, Funda Aydin, and her continuous support, sacrifices, and believe in me. Your smile and kindness have served and enlighten my life.

Finally, I would like to express my gratitude to Atlantic International University and the School of Social and Human Studies Academic Department for giving the funding and scholarship that enabled this study. I really appreciate your dedication to promoting research and expanding knowledge. This dissertation is a testament to the collective effort, support, and encouragement of all these individuals and institutions. To each and every one of you, I extend my deepest thanks and heartfelt gratitude.

Thank you.

Alptekin Aydin

Table of Contents

Acknowledgments ***46***

Table of Contents ***47***

Abstract ***53***

CHAPTER 1: INTRODUCTION ***55***

1.1 Importance of the Study **55**

1.2 Autism Spectrum Disorder (ASD) **56**

3.2 The Importance of Timely and Ongoing Intervention **58**

3.3 Difficulties Associated with Current Therapeutic Methods **58**

3.8 Constraints of Current Interventions **67**

4.1 Social Communication: Definition and Components **69**

4.2 Consequences of Impaired Social Communication **70**

4.3 Significance for Child Development and Academic Achievement **70**

4.4 The Impact of Social Isolation on Family Dynamics **71**

5. The significance of the Dorsolateral Prefrontal Cortex (DLPFC) in social communication **73**

5.2 The anatomical connections and functional implications **74**

5.3 Consequences for Interventions **75**

6.1 Introduction to Transcranial Direct Current Stimulation (tDCS) **77**

6.2 The mechanisms via which tDCS works **77**

6.3 Preliminary Research and Evidence About Transcranial Direct Current Stimulation (tDCS) for Autism Spectrum Disorder (ASD) **78**

7. Considerations about the methodology and design of the study **80**

7.1 Utilising a combination of qualitative and quantitative methods 80

7.2 Qualitative Component: Survey of Families with a Structured Format **81**

Consent and Privacy Protection **81**

CHAPTER 2: LITERATURE REVIEW ***84***

Autism Spectrum Disorder (ASD) **88**

Prevalence and Epidemiology **89**

11. Analysing Autism Spectrum Disorder (ASD) **93**

11.2. Study of Prevalence **102**

11.3 Symptoms and Behavioural Traits **103**

11.4 Effects on Daily Life and Functioning **105**

Behavioural interventions **107**

12.1 Applied Behaviour Analysis (ABA) 107

12.2 The Efficacy and Results **109**

13. Training in interpersonal skills **111**

13.1 Techniques and Strategies 111

13.2 Influence on Social Interactions **113**
13.3 Programmes for Exceptional Education **114**
13.4 Advantages and Difficulties **116**
14. Strategies for promoting inclusive education **118**
14.1 Models of Inclusion 118
14.2 Results for Children with Autism Spectrum Disorder (ASD) **120**
15. Treatment options for fundamental symptoms **122**
15.1 Categories of Medications Utilised 122
15.2 Effectiveness and Adverse Reactions **122**
15.2 Effectiveness and Adverse Reactions **124**
15.3 Prevalent Comorbidities **125**
15.4 Approaches to Treatment and their Effectiveness **126**
16. Constraints of Current Interventions **128**
16.1 Behavioural Interventions 128
16.2 Implementation Challenges **129**
17. Interventions aimed at improving education **130**
17.1 Concerns Regarding Accessibility and Quality 130
17.2 Constraints in Addressing All Symptoms of Autism Spectrum Disorder **131**
18. Pharmacological interventions **132**
18.1 Adverse Effects and Long-Term Consequences 132
18.2 Limited Effectiveness for Social Communication Deficits **133**
19. Assessment of Overall Effectiveness and Identified Gaps **133**
19.1 Current Treatments' General Deficiencies 133
19.2 Action Mechanisms **138**
20. Neurodevelopmental Disorders and their Applications **141**
21.Considerations regarding safety and ethics **148**
22. First trials of transcranial direct current stimulation (tDCS) in subjects with Autism Spectrum Disorder (ASD) **151**
22.1 Enhancements in Social Communication **151**
22.2 Reports from Parents and Carers **152**
22.3 Constraints and Prospects for Future Development **153**
22.4 Ethical Considerations **154**
23. Dorsolateral Prefrontal Cortex (DLPFC) stands for the region of the main occupant of the prefrontal cortex. **155**
23.1 Contribution to Social Cognition **158**
23.2 Neuroimaging Studies Provide Evidence **160**
23.3 The significance of family involvement is **169**
The impact of family support on treatment outcomes **170**
23.5 Achieving a Balance Between Benefits and Side Effects **176**
23.6 Findings from Families on tDCS and Other Techniques **178**

The significance of mixed-methods research **182**
The advantages of integrating quantitative and qualitative data **182**
24.3 Illustrative Instances from Preexisting Literature **184**
Results and knowledge acquired **186**
24.4 Methods for Consolidating Data **187**
25.1 Advancements and Innovations in Interventions and Technologies **190**
25.2 Technological Advancements **191**
The key findings from longitudinal studies are as follows: **195**
25.5 Approaches to Personalised and Precision Medicine **197**
Implications for policy and practice **200**
26.2 Clinical Practice Recommendations **201**
23. Summary of Key Findings **203**
27.2 Consequences for the Field **205**
27.3 Future Research Implications **206**
27.4 Suggestions for Future Research **208**
CHAPTER 3: RESEARCH METHODS ***210***
28.The purpose and scope of the methodology **211**
28.1 Experimental Methodology **213**
29. Rationale for Selected Design **214**
30. Quantitative Component **216**
30.1 Qualitative Component **217**
32. Ethical Considerations **217**
32.1 Participant Selection Criteria Description **218**
32.2 Criteria for Inclusion and Exclusion **219**
32.3 Participant Demographic Information **220**
33. Ethical Considerations **222**
33.1 .Designing a survey **224**
33.2 Survey Sections and Question Types Description **227**
34. .Methods for Collecting Data ***234***
34.1 Introduction to Data Collection Methods **234**
34.2 Informed Consent Process: **235**
34.3 Summary of Data Collection Methods **237**
35.1 Data Cleaning and Preparation **247**
35.2 Techniques for Quantitative Analysis **249**
35.3 Techniques for Qualitative Analysis **250**
35.4 Software and Tools for Data Analysis **252**
36. .Ethical considerations **252**

37. Conducting pilot testing and making necessary revisions 261
37.1 Constraints and Difficulties 262
37.2 Difficulties in Data Collection and Analysis 265
37.3 Approaches to Address Limitations and Challenges 265
37.4 Practical Implications for Clinical Practice 268
38. Quantitative Component: Randomised Controlled Trial (RCT) 270
39. Qualitative Aspect: Structured Family Survey 271
40. Methods for Gathering and Analysing Data 271
40.1 Thorough Assessment 272
40.2 Improved Validity and Reliability 272
40.3 Expert clinical and policy recommendations based on informed knowledge and expertise. 272
CHAPTER 4: FINDINGS 273
41. Summary of Findings Chapter 273
41.1 The structure of the Results section 274
41.2 provides an overview of the demographic characteristics of the participants. 276
41.3 Parental Education 279
41.4 Occupational Status 280
41.5 Household Income 281
42. Socioeconomic Consequences 282
43. Academic History 282
43.1 Supplementary Academic Assistance 284
44. Implications for Education 284
45. Results of the initial assessment 285
46. Evaluation against established standards 292
47. Significance of Baseline Comparisons: 294
48. Social communication skills after receiving treatment 296
49. Alterations in behaviour and cognition 299
50. Results of the assessment conducted after the treatment 304
50.1 Alterations in behaviour and cognition 306
51. Statistical analysis of numerical data 313
51.1 Inferential Statistics 315
51.2 Regression Analysis 320
60. Enhancements in Interpersonal Communication Abilities 325
60.1 Enhancements of Behaviour and Cognition 326
60.2 The Impact of Family Dynamics on Parental Stress 326
60.3 Practical and Future Research Implications 327

60.4 Areas for Future Research 328
61. Significance for Theoretical and Practical Applications 353
CHAPTER 5: DISCUSSION 356
63. Significance for Theoretical and Practical Applications 358
64. Obstacles and Prospects for the Future 358
65. Objective of the Discussion 359
65.1 The scope of the discussion 359
66. Comparison to prior studies: 360
67. Theoretical Implications: 360
68. Areas for Further Investigation: 362
70. Summary of Key Findings from the Research 362
70. Behavioural and Emotional Regulation 363
71. Assessment of Family Contentment and Encounter 364
72. Comparative analysis with prior studies 365
73. Significance of the Results 366
74. Augmentation of neural circuits implicated in social cognition 366
75. Effects on Cognitive Processes 367
76. Justifying Hypotheses and Theoretical Frameworks 368
77. Wider ramifications for the management and implementation of treatment and intervention strategies 368
77.1 Improvements in Nonverbal Communication 371
77.2 Establishing Significant Connections 372
77.3 Family Experiences and Perceived Benefits 373
78. Observations of Challenges and Limitations 374
79. Incorporation into Preexisting Literature 376
80. Supporting Evidence for tDCS as a Therapeutic Intervention 378
81. Practical Applications 380
82. Suggestions for Educators and Therapists 381
83. Advantages and Constraints of the Research 384
84. Further Actions to Minimise Harm, and Suggestions for Future Study 387
84.1 Areas for Future Research 388
85. Overview of the Conversation 393
CHAPTER 5: CONCLUSION 394
86. Overview of the Study's Goals 395
86.1 Overview of Main Discoveries 396
88. Summary of Study Contributions 402
89. Contribution to Transcranial Direct Current Stimulation (tDCS) Research 403

90. Incorporation of Family Viewpoints **405**
91. Repercussions for the application of medical treatment **406**
91.1 Guidelines for Healthcare Providers **407**
92. Possible Influence on Therapeutic Approaches **408**
93. Advantages of the Study **410**
94. Study Constraints **413**
95. Topics requiring additional research **416**
95.1 Proposed Methodological Enhancements **417**
95.2 Feasibility of Conducting Longitudinal Studies **418**
96. General Observations on the Study **420**
Reference ***424***
Appendix ***440***
Appendix A **441**
Appendix B **442**
Research Protocol **442**
Appendix C **445**
Information Sheet 445
Appendix D **448**
Appendix E **453**
Appendix F **473**
Appendix G **482**

Abstract

Introduction Autism Spectrum Disorder (ASD) is a neurodevelopmental disorder characterised by substantial deficits in social communication, behavioural regulation, and the execution of cognitive processes. Although conventional interventions exist to support acute symptom improvements in children with ASD, the social communication, behavioural, regulatory, or executive function difficulties can often be difficult to address in this manner. Transcranial Direct Current Stimulation (tDCS) has emerged as a novel intervention that could potentially enhance complex functions such as these in children with ASD. Overall, the aim is to assess the efficacy of tDCS in enhancing social communication skills, minimising behavioural issues and cognitive impairments in ASD children, and exploring how tDCS can be utilised in therapeutic practices to support ASD children and their families.

Methods Data were collected using a multimethod approach that included retrospective structured surveys, standardised assessments and qualitative interviews from a chronological cohort of 265 participants aged 3 to 18 years. All participants were assessed in domains, including verbal and non-verbal communication, social reciprocity, behavioural issues and cognitive functions at baseline and following a course of tDCS treatment. Quantitative data was analysed using Paired sample t test, Analysis of Variance (ANOVA), Correlation and Regression. Qualitative feedback from parents and care givers was garnered to add context and depth to the quantitative findings. Findings Statistical analysis showed remarkable advances in verbal communication including increase in the vocabulary size, development of sentence formation and frequency of initiating conversation. Non-verbal communication also saw improvement including better eye contacts, more expressive facial expression and appropriate use of

gestures which was statistically significant in Social Responsiveness Scale (SRS) and Autism Diagnostic Observation Schedule (ADOS) scores. Behavioural scores for repetitive behaviours and emotional dysregulation dropped considerably (by up to three and two points, respectively) at the end of the intervention when using the Repetitive Behaviour Scale-Revised (RBS-R) and Aber-rant Behavior Checklist (ABC). Participants also showed better emotional regulation, fewer mood swings, decreased anxiety and aggression. And their executive function (EF) abilities, such as working memory, cognitive flexibility and inhibitory control, greatly improved, which in turn led to better academic performance and day-to-day functioning.

Parents and carers reported lower levels of parental stress, improved routines within the family and improved relationships between siblings. Children had more social engagement, such as joining in group activities and forming long-lasting friendships. The thematic analysis of the interviews showed that the patients benefitted from the individualised treatment plans, the continuous professional support, and the tDCS as a means to complement other therapeutic treatments. Conclusion Overall, we found that tDCS is a promising approach for children with ASD, showing robust gains in social communication, behavioural regulation and cognitive functions. Tailored approaches to tDCS are needed, depending on the target, age, severity of autism, and the specific areas of deficit. Long-term follow-up and larger-scale studies are needed to validate these findings and ensure the persistency of the benefits of the treatment over time.

CHAPTER 1: INTRODUCTION

The rationale and scope of the study (also called thesis proposition or thesis statement) reflect the specific goals and scope of study. This study assesses the effects of transcranial Direct Current Stimulation (tDCS) with the anode specifically over the Dorsolateral Prefrontal Cortex (DLPFC) on social communicative aspects in children diagnosed with Autism Spectrum Disorder (ASD). The purpose of this study is to evaluate both the clinical outcomes and the family experience in relation to this treatment. To gain insight in the family perceptions of the experienced benefits and problems, and perceived satisfaction with tDCS, a qualitative survey will be used to obtain data from families. As said before, by considering both the clinical and familial perspectives, a more comprehensive understanding of the effectiveness and feasibility of tDCS treatment in ASD might be reached (Fregni et al., 2021).

1.1 Importance of the Study

The research can be beneficial in helping to increase the effectiveness of treatments that are being developed for Autism Spectrum Disorder (ASD), by providing new insights into the underlying structures that may need to be altered during treatment. New treatments may need to better target the fundamental symptoms of Autism Spectrum Disorder (ASD), which can incorporate difficulties in social communication.

New interventions such as using Transcranial Direct Current Stimulation (tDCS) could potentially improve the brain's ability to change. It could, therefore, improve social communication skills in children with ASD and have consequences for clinical practices, as this could offer a novel approach for intervention that is non-invasive and potentially more accessible than traditional treatments (Maenner et al., 2020)

From a DSM-5 perspective, Autism Spectrum Disorder (ASD) is characterised by persistent deficits in social communication and interaction across multiple contexts along with restricted, repetitive patterns of behavior, interests, or activities. This deficit is rather broad as the DSM-5 using specifics in assessing ASD such as in 'social-emotional reciprocity: ranging from abnormal social approach and failure of normal back-and-forth conversation; to reduction of sharing of interests, emotions or affect; to failure to initiate or respond to social interactions'. The external characterisation of ASD sufferers can show impairments in 'nonverbal communi-cation used for social interaction' (American Psychiatric Association, 2013). Transcranial Direct Current Stimulation (tDCS) is a non-invasive stimulation of the brain, during which a gentle electrical current is applied at several points on the scalp to regulate the activity of neurons. A key brain region critical to good executive functioning, a key social cognition and the regulation of behaviour is the Dorsolateral Prefrontal Cortex (DLPFC), and so this spatially targeted brain region is a focus of tDCS therapies (Miller Cohen, 2001).

1.2 Autism Spectrum Disorder (ASD)

Autism Spectrum Disorder (ASD) is a complex neurodevelopmental disorder characterised by persistent difficulties in social communication and interaction, and restrictive and repetitive behaviours, interests and activities. In accordance with the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), the diagnostic criteria for ASD include deficits – as manifest by full or partial impairments – in the four following areas: social-emotional reciprocity, nonverbal communicative behaviours used for social interaction, and (iii) developing, maintaining and understanding relationships (American Psychiatric Association, 2013).There-

fore, while one person could be diagnosed with ASD because of repetitive interest in a particular topic, another could be diagnosed because of severe challenges in interacting with other people. The severity and type of symptoms can vary greatly, and – hence – the term 'spectrum' is used, as it reflects the broad variation of challenges experienced by people with ASD.

The DSM-5 outlines explicit criteria for ASD – such as persistent deficits in social communication and interaction across multiple contexts – that cannot be accounted for by general developmental delays. These difficulties involve deficits in social-emotional reciprocity, such as failure to respond to informational nonverbal behaviour, a lack of interest in sharing activities, and a lack of using back channels to regulateopssa1communication. People with ASD at times display restricted and repetitive patterns of behaviour, interests or activities, through stereotyped body movements, insistence on sameness, highly restricted, fixated interests, or hyper- or hypo-sensitivity to sensory input (American Psychiatric Association, 2013). Over the past 50 years, the occurrence of Autism Spectrum Disorder (ASD) has dramatically increased. The most recent estimates from the Centers for Disease Control and Prevention (CDC) show that approximately 1 in 54 children in the United States are born with Autism Spectrum Disorder. (Maenner et al, 2020) The reasons for the increase are diverse, and include factors such as increased awareness, changes to the diagnostic criteria, and technological advances in screening and diagnostic equipment. The increase in the incidence of Autism Spectrum Disorder (ASD) signifies the immediate need for an understanding of Autism Spectrum Disorder (ASD), and improved services for individuals with ASD and their families.

ASD symptoms often appear in early childhood, and may last a lifetime, leading to significant impairment in many aspects of daily life including communication, social interactions and educational performance. Children diagnosed with ASD often have significant impairments in social interaction and communication, typically leading to social withdrawal and, consequently, affective functioning. For instance, they might have difficulties in maintaining eye

contact, understanding social cues, or engaging in reciprocal play with their peers. These difficulties can extend into adulthood, resulting in reduced job opportunities, independence and psychological wellbeing (Lord et al., 2020).

3.2 The Importance of Timely and Ongoing Intervention

Therefore, the most beneficial intervention for children with ASD is one early enough to make a difference and sustained enough to continue that difference. The earlier the intervention, the greater the positive impact on the long-term developmental outcome. Targeted early intervention in the critical period of brain development can lessen ASD symptoms and improve adaptive functioning. For instance, interventions that emphasise the development of communicative, social and adaptive behaviours can greatly enhance children's ability to function independently and connect with their environment and peers. These results have been well documented (Dawson et al., 2010). Continuing this type of intervention is crucial to help children continue to grow, and to stay one step ahead of the inevitable challenges that arise as children mature. Sustained therapy can help maintain and build on the gains achieved through early intervention.

3.3 Difficulties Associated with Current Therapeutic Methods

Most of the therapeutic approaches to ASD currently employed involve behavioural therapies, educational support and, in some cases, drug interventions for specific symptoms. Behavioural therapies, such as Applied Behaviour Analysis (AB in that they can improve many important skills. Applied Behaviour Analysis (ABA) is a method that seeks to strengthen positive behaviours and reduce maladaptive behaviours, using systematic procedures. It has been helpful for

teaching communication, social skills and adaptive behaviours as was shown by Reichow et al in 2018. However, Applied Behaviour Analysis (ABA) often requires intense and prolonged effort, and it doesn't always generalise to situations outside the therapy.Stimulants, such as methylphenidate, are also an effective treatment for the hyperactivity and inattention that often go along with ASD. Research shows that they help kids with ASD just as much as kids with ADHD who do not also have ASD. More than half of children with ASD respond favourably to stimulant treatment; their attention improves, usually with a reduction in hyperactivity (Handen et al., 2000). The most common side-effects are decreased appetite, sleep disturbances and heightened anxiety or irritability. These side-effects can be minimised with careful monitoring and dose adjustment.

Overall, drugs that treat the symptoms of ASD can offer many benefits; however, they must be monitored carefully and given in combination with behavioural interventions. More research is needed to fully understand the long-term effects of these drugs and to develop medications that are safer and more beneficial. Common Comorbidities: between % and 80% of children with ASD also have one or more comorbidities, which can substantially impair functioning or quality of life.

Among the host of co-occurring health issues are Attention Deficit/Hyperactivity Disorder (ADHD) anxiety disorders, depression and epilepsy, as well as gastrointestinal problems and sleep disturbances according to Simonoff et al in 2008. Such co-morbid conditions need to be addressed, especially as they can exacerbate the symptoms of ASD and add to the burden for the child and, ultimately, for the family. One fairly common comorbid psychiatric problem is attention deficit hyperactivity disorder (ADHD), which usually manifests in 30–60 per cent of children with an ASD, with symptoms of inattention, over activity or impulsivity that can interfere with interactions and academic performance. If you can think of any other situation that

you might not pay as much attention to and then have to interact socially with the people in that situation, it's a recipe for overwhelming anxiety 18% to 40% of children and adolescents with an ASD may also have an anxiety disorder, such as anxiety disorder, social anxiety disorder or specific phobias (fears of specific things). It also appears that, in addition to anxiety, depression might also be a fairly common comorbid DSM-5 psychiatric problem, given that relatively small numbers of children (about 10%) experience depression, but it becomes more common during adolescence and adulthood, so people's clinical profile becomes more complicated.

Epilepsy. A seizure disorder that affects 20 to 30 per cent of people with ASD. That occurs, he says, 'from absence seizures to severe generalised tonic clonic seizures', as his 2007 paper in Neuropsychiatric Disease and Treatment demonstrates. Gastrointestinal problems including constipation, diarrhoea and acid reflux can affect up to 70 per cent of children with ASD. (McElhanon et al, 2014) Insomnia, including difficulty falling asleep, staying asleep or getting enough sleep, disrupts daytime functioning and behaviour (Richdale .. Schreck 2009).

In young children with ASD, the treatments are with a combination of pharmaceutical and non-pharmaceutical approaches. The choice of treatment and the way it is managed will be based on the type of disorder, the severity of the disorder and what the needs of the child are. However, stimulants, such as methylphenidate (ie, Ritalin®) and amphetamines, are the most frequently used medication to target ADHD symptoms in most children with ASD. They have been shown to reduce hyperactivity, impulsivity and inattention in children with ASD and it is estimated that from 50-70 per cent have found some benefit from these medications when used over six months. Non stimulant medications such as atomoxetine and guanfacine are used both with stimulants or in the children who cannot tolerate medications anymore.

Also common is treatment with serotonin reuptake inhibitors (SSRIs) such as fluoxetine (Prozac) and sertraline (Zoloft), which can reduce levels of anxiety and enhance mood and behaviour. However, it's important to watch for side-effects such as heightened agitation or changes in behaviour. CBT is another type of medical approach that is beginning to show promise as an adjunct to treatment for anxiety in children with ASD, although few studies have yet to confirm robust findings. Medications are used therapeutically to help treat. That is, while none explicitly target epilepsy, they target other conditions, and particularly, conditions that can impair quality of life for those with epilepsy. Additionally, animals, things, activities and places that interfere in some way with the normal functioning of one's body or mind often serve as a reminder to take medication and dosing as an aid to memory. Examples include valproate for generalised tonic-clinic seizures; lamotrigine for temporal lobe epilepsy; and levetiracetam for partial seizures. Medications are only one of the various aspects of a person's life that can identify them as a person with epilepsy: Examples of medications that can address seizure aetiology may include? For example, a person with epilepsy may need medications for hypertension? A person with epilepsy may have only one medication.

a choice of medications that suit the type of seizures; diagnosis an explanation of the person's health status; a potential side or adverse effect as long as seizures are monitored by a professional several times a day and medications adjusted to achieve seizure-management goals, this care is safe. Children frequently experience more gastrointestinal (GI) symptoms such as constipation, diarrhoea and reflux, which may be helped by adjustments, use of probiotics and abdominal pain and anti-reflux medications for pain control. Some children benefit from specific medications, such as laxatives for constipation, and, for reflux, proton pump inhibitors (PPIs) that help control the acid that moves up the GI tract. Plastic wrap used to secure the oesophageal tube may cause pain and leaking, potentially leading to rashes or serious skin

infections around the stoma site. A careful evaluation by a gastroenterologist can help to develop individualised treatment for each child. In their case, establishing a daily routine, as well as developing good sleep habits, could prove effective in children with ASD. In others, administrations of melatonin supplements can be used to help regulate the circadian cycle, and facilitate not only sleep onset but also its duration (Malow et al., 2012).

Medications such as clonidine or trazodone might be a treatment option for a child with ASD who presents with sleep disorders. To be more effective with these children, a range of flexible treatments must account for their individual needs. This pharmacological treatment can then be supported by therapeutic approaches so that the whole family can palliate or ameliorate differences that render cohabitation and socialisation like other families very difficult. Through this process, we can hopefully, together, help children persevere and succeed while acknowledging their differences. Most important of all is to foster resilience and to highlight and encourage important unique characteristics of their development. Monitoring and ongoing oversight is crucial by a team of physicians, educators and families who can target and treat any coexisting neuropsychiatric or medical issues associated with ASD to ensure the best quality of life for these children and their families.

Constraints of Current Interventions

We commonly talk about interventions such as Applied Behavior Analysis (ABA) as if they are effective. But ABA is widely studied as a treatment of Autism Spectrum Disorder (ASD) in children. It has been shown to have positive effects on communication, social interaction and common behavioural challenges, but its efficacy ranges across different individuals with ASD. Studies have shown that the intensity and duration of therapy, the timing of beginning

the therapy, and the existence of co-morbid conditions alongside treatment can affect the efficacy of ABA in individuals with ASD. Another glaring weakness of ABA is that it solely employs reinforcement techniques to shape behaviour, on what Psychologists call 'shaping', a cornerstone of ABA, to transform fundamentally. ABA can't change anything cognitive and emotional: all it aims to do is shape behaviour If the kid you're working with is able (typically) to understand the world, is able to appreciate the context in which certain behaviours may incur societal consequences, is capable of reining in behaviour and interacting accordingly, then ABA can work towards improving crucial elements in these interactions such as reading social cues or developing together a personalised framework of social relationship.

Furthermore, although ABA is promising in decontextualized settings, adapting its benefits to real-world environments is not straightforward (Howlin et al, 2009). For some children, it can be unnecessarily rigid and highly structured. The constraints of the ABA programme might not be a good fit for some children with autism because their day-to-day experience is unpredictable and so varied. The issue with ABA – at least with the dominant approach – is the lack of transfer and implementation of skills, in life.

Challenges, in Implementation

Still, there are very real logistical obstacles to making use of ABA and other behavioural therapies. The child needs a certain amount of time and energy from the therapist and the family, which can be hard to keep up week in, week out. Many intensive ABA programmes suggest that children undergo therapy for 20 to 40 hours per week. This can be hard for families to keep up.

Additionally, both families with limited insurance coverage and parents in stressful situations can find the cost of ABA treatment prohibitive. A third challenge is a lack of trained ABA therapists to work only with children with autism, especially in rural areas or for children who do not live in affluent communities who can afford a private provider. All of these can lead to waiting lists and inconsistent services. There are also concerns about what ABA entails, among some in the community, with some critics arguing that ABA's emphasis on compliance and malleability can undermine a child's sense of autonomy and individuality (Devita Raeburn, 2016). These kinds of considerations make it harder to implement (and to gain consent for) using ABA as a treatment for ASD. Improving education via interventions may also be more inclusive and improving the quality of schooling, but it can also raise concerns about accessibility and quality. Finally, interventions offered within the education system, such as ASD programmes or providing inclusive education to children with ASD, are designed to provide 'evidence-based' or 'research-based' targeted support to children with ASD. Yet availability and quality of such interventions can be highly variable. Many locations do not have sufficient financial resources or qualified personnel to deliver programmes to children with ASD.

Programmes in special education often have inadequate funding, and classrooms can be understaffed, with teachers having insufficient training and little to no time and the resources to provide individualised support for students with ASD. In some cases, few services available for young adults translate to limited educational experiences and outcomes. Furthermore, services related to special education services vary by state and municipality, which, accordingly, can shape the support a family can access (National Research Council, 2001).

Moreover, special education programmes may not always cater to the needs of children with ASD, and they can adapt to facilitate their adjustment to the demands and diversity of school

life. There are challenges associated with addressing all components of Autism Spectrum Disorder, given that educational programming (while a benefit, for skill-building and structure), is less comprehensive in addressing the symptoms that many children with ASD present with, specifically social communication and emotional regulation, which may be overlooked.

Similarly, there are challenges facing educational strategies that aim to include ASD students in classrooms. Classroom teachers across these settings often do not have the training to provide the necessary support for ASD students to thrive. Without support or resources, these children will struggle to keep up with their peers, eventually beginning to feel frustrated and excluded. Education models invariably require spatial and temporal adjustments within the school building (for example, updating staff training, redesigning curriculum structures, adapting classroom management processes) and these organisational shifts can be particularly difficult for schools already stretched for resources. Educational interventions are naturally part of the mix, but these are often limited by the specific difficulties of the educational environment and an overwhelming emphasis on academic achievement.

While effects and long-term outcomes Many drugs used to treat Autism Spectrum Disorder (ASD) concentrate on symptom-specific issues, such as irritability or hyper-arousal (hyperactivity) or anxiety, and not the core features of ASD, such as social communication deficits or difficulties with social interaction. Medications used include antipsychotics, antidepressants and stimulants. All drugs yield a number of unwanted side-effects that might alter the child's general wellbeing. Even if antipsychotic drugs such as risperidone or aripiprazole bring comfort, they often lead to weight gain, sleepiness and metabolic changes that might become significant long-term health issues. Stimulant medications prescribed for the symptoms of co

ADHD can diminish appetites, disrupt sleeping patterns and increase anxiety levels. Antidepressants commonly prescribed for anxiety and depression can likewise also be hard on people. More agitation.

Although long-term outcomes of these medications in children with ASD are unclear, mostly because of limited research evaluating extended safety and efficacy, the high risk for dependence and teratogenic adverse effects - acutely emphasised by Coury et al (2012) - require cautious monitoring and periodic re-evaluation of the treatment plan and need for continuation. Furthermore, the use of drugs for a number of conditions (polypharmacy) leads to more complicated regimes and more possibilities for harmful interactions between drugs which, as Coury et al (2012) have noted, require coordination across disciplines within any health system, but especially in resource-poor settings.

Moreover, pharmacological interventions might be helpful as an adjunct to managing some troublesome behaviours, particularly hyperactivity, but are generally ineffective in addressing the fundamental symptoms of ASD centred around significant impairments in social communication and interaction. Medications do not teach skills. What is needed, as Posey et al (2008) so eloquently point out, is ‘immersion in real world social interactions’ so that one can enhance understanding and empathy. Consequently, the treatment of autism spectrum disorder (ASD) is frequently a combination of medications in conjunction with educational interventions, rather than a single all-inclusive solution to the bother that this condition causes. Once again, drugs are not touted as the primary remedy. Rather, this multi-modal approach aims at symptom amelioration, while underscoring the limitations inherent in relying on drug intervention as this approach to treatment. Toward that goal, there is a gap in interventions meant to remediate social communication skills that should be urgently addressed.

In addition, caregivers and parents often find it hard to get children to stick to the medication regimen, to check for side effects and to keep up with their healthcare providers, which can be burdensome and increase the need for better treatments (Johnson et al, 2013).

Assessment of Overall Effectiveness Identified Gaps

Overall efficacy of ASD treatments is lacking many of these therapies teach children with ASD to conform by managing their symptoms and forcefully modifying behaviour, rather than accommodating to their needs. ABA, on the other hand, works by moulding behaviours, but does not address the emotional aspects of the spectrum. Educational programmes that improve performance might fail to do the same if the social inventions needed to foster the growth of emotional sensibilities are ignored.

3.8 Constraints of Current Interventions

There are many different therapy options to aid with ASD, however, the current methods to control ASD are severely limited. One major issue is the heterogeneity of the disorder. ASD is highly heterogeneous – it has varying symptoms and different degrees of severity, making worldwide interventions difficult. People might need individualised and broad regiments to treat their ASD, and while these can indeed be quite resource intensive, they can also be difficult to enact routinely. While behavioural therapies work, they are also costly in terms of time and money. The intensity of programmes can make them inaccessible to families. There is also a need for more flexible and accessible solutions that can develop skills without the focus shown in structured therapy environments and translate those skills to life experiences for which parents will always need to tweak and modify interventions.

Clearly, educational supports are crucial – but the extent to which they can compensate for the trauma of assimilation to a social world one cannot readily navigate might be the victim of the unequal quality of resources and training that will be, or rather, won't be made available in different schools and districts. As a result, children with ASD will receive unequal opportunities to provide them with a decent education, which is predictive of their level of future academic and social achievement. What is more, educational therapies will not solve, on their own, the broader social communication problems these children endure. (Kasari et al., 2011) While pharmacological interventions might alleviate symptoms, they don't directly target the core adverse effects and differences in pharmacological response further add to the difficulty in using pharmaceuticals, meaning that there is an urgent need for more research on safer and more effective pharmacological treatments. Ultimately, for individuals with ASD, multidimensional integrated treatment programmes that combine pharmaceutical, behavioural and educational interventions will be needed to fulfil their unique and individualised needs (Anagnostou and Hansen, 2011).

In light of these constraints, there is an urgent need for novel and comprehensive treatments for ASD. tDCS is a form of emerging therapy that may have the potential of increasing brain plasticity and social communication skills. The development and application of more effective therapies with the ability to promote independence and social connection and improve quality of life of individuals with ASD and their families is of extreme significance. This requires joint effort from clinicians, researchers and families. Fregni, F, Lemere, Y, Woods, J, Srinivasan, D, Di Mesio, F, Turkeltaub, P, Schenck, C and Cohen, L (2021). 'Brain-based interventions for autism spectrum disorder: a practice and research update', Clinical Neuropsychologist. 35(1): 135 149.

4.1 Social Communication: Definition and Components

Social communication is the ability to use verbal and nonverbal communicative behaviours in social contexts, to understand and respond to social signals, and to engage in social reciprocity. Children with ASD are often impaired in many of these activities. Social communication includes a series of behaviours that enable individuals to be social communicators: Verbal communication is the use of expressive language to articulate thought, needs and emotions. This includes the ability to initiate and hold a conversation, as well as the appropriate use of language across social contexts.

Nonverbal communication is defined as communicating through eye contact, facial expressions, gestures and body language. Often, nonverbal communication is the primary form of communication used between individuals; with nonverbal cues, we begin to understand the emotional state and intentions of a person. 'Social cognition' refers to the ability to engage in and interpret social signalling using vocal intonation, physical movements and facial expression; also, it's the ability to understand other minds, including what they think and feel – often called 'theory of mind' (Tager-Flusberg, 2000).• Pragmatic Language Skills: the ability to use language effectively and appropriately in social contexts, such as understanding idioms, jokes and other forms of figurative language. Pragmatic skills are essential for social participation in valued social relationships (Adams, 2005).

4.2 Consequences of Impaired Social Communication

Children with a diagnosis of Autism Spectrum Disorder (ASD) often display compelling deficits in social communication, affecting numerous aspects of their lives and development. Such deficits can lead to: Children with ASD might be excluded from social interactions and social events. This may cause them to feel cause them to struggle to understand the rules of social interaction and to act inexperienced in social behaviours (Klin et al., 2007). • Frustration and Behavioural Problems: The inability of good communication may lead to frustration and behavioural problems. Children with ASD may tantrum, be violent or self-injurious to express unmet needs or feelings (Hart in Pratt, 2004). • Emotional Well-being: Long-term difficulties with social communication can cause a child to be less confident and have lower self-esteem. Children who have an ASD often experience anxiety, sadness and low self-esteem because of their persistent issues with social engagement (Capps et al., 1998). • Academic Problems: Problems with social communication might lead to academic problems. For example, children with ASD might struggle to follow a task, to participate in group tasks or to discuss things in class. As a result, educational opportunities might be limited and academic success hampered (Tager-Flusberg, 2000).

4.3 Significance for Child Development and Academic Achievement

Good social communications support child development in important ways, fostering a child's learning, social experiences and the management and regulation of emotions. Children learn how to read and respond to social cues, develop the ability to understand others' perspectives, and build secure attachments establish relationships through their social interactions. Efficient interpersonal communication is essential for: And language acquisition and development

builds upon these social communication skills over time. Children master word meanings, grammatical knowledge and conversational competencies by interacting with carers and friends.

• Cognitive development: Social contacts develop cognitive development by promoting the utilisation of critical thinking and problem-solving skills, and by helping to master the learning of different perspectives. Participating in conversations and social interaction on cooperative tasks help children to improve their cognitive faculties (Vygotsky, 1978). Emotional regulation: The capacity for social communication contributes to the development of self-regulation and coping skills. The capacity to understand one, and to recognise the feelings of others constitute the rudiments of emotional intelligence – a crucial factor in one's general mental health and wellbeing. (Hart, 2004) • Academic achievement: Social communication pursuits equip children to participate actively in classroom discussions, work productively in groups, and engage meaningfully with their teachers – all of which are instrumental in learning effectively and realising success in the classroom. 'Children who can describe their needs and understand the rules of the classroom are more likely to reach academic success.' Howlin et al (2004)

4.4 The Impact of Social Isolation on Family Dynamics

Family members of children with ASD can face challenges from the child's social communication difficulties, which can spill over into the family. Families of children with ASD can experience a number of problems and sources of stress relating to the child's social communication difficulties: • Parental Stress and Mental Health: Parents of children with ASD might face an increase in stress levels and have difficulties in maintaining a the long-term burdens of coping with their children's deficits in social communication and behaviour, as well as the

associated psychological issues and the challenges of finding an effective therapy (Estes et al., 2009).

• Sibling Experience: The siblings of children with ASD may also experience social cognitive impairments as a result of living with a brother or sister who has breakdowns in social communication. Kaminsky and Dewey it therefore comes as no surprise that siblings may experience feelings of neglect, frustration or embarrassment in the company of their sibling with autism, but also may take on caregiving roles that compromise their own social and emotional development.

• Family Activities and Social Life: Reduced social communication may restrict the family's social activities and attendance at social events and community life. Social avoidance may lead to families not engaging in social activities because they fear their child's conduct or social appropriateness, leading to social isolation for the whole family (Woodgate, et al, 2008).

• Financial cost: Costs associated with therapy and involved in changing social communication problems can be very high. There can be considerable financial suffering for families due to the cost of treatment, which results in families becoming less well-off and having less access to resources (Järbrink, Fombonne and Knapp, 2003). Early intervention in children with ASD that focuses explicitly on social communication has been found to result in better outcomes for the child with ASD and the whole family by reducing social communication problems. It can effectively reduce problem behaviour and affective problems, and promote successful outcomes for children with ASD and their families.

5. The significance of the Dorsolateral Prefrontal Cortex (DLPFC) in social communication

Researchers study the role of the DLPFC in executive functions and social cognition. A key brain area for these aspects of functioning is the Dorsolateral Prefrontal Cortex (DLPFC) and it plays roles in a host of high-level cognitive phenomenon including executive functioning and social cognition. Executive functions are a cluster of cognitive processes that include working memory, cognitive (Miller Cohen, 2001). These functions allow people to better plan, focus, retain instructions and juggle multiple tasks. Impairments in executive processes are associated with problems regulating behaviour, emotion and making decisions, all of which are more common in individuals with ASD (Diamond, 2013).

Social cognition is a particular function of the DLPFC and refers to the ability to process, integrate and analyse social information – for example, to recognise emotions and interpret social cues, and to understand other people's perspectives. By linking cognitive and emotional information, the ability to respond appropriately in social contexts can be accomplished. \[Sorzant: Social cognition, the ability to comprehend and interpret social information, is one of the most important functions of the DLPFC. It is how we understand other people's emotions, interpret their social cues, and even comprehend their perspective.

By combining cognitive and emotional information, the DLPFC enables us to respond appropriately in social situations. This is an important aspect of overall DLPFC function. Therefore, the DLPFC plays a crucial role in our ability to assess our social environment, decide which form of communication is best to respond to a given situation and understand the intentions and feelings of other people. These abilities are essential for successful social interaction when you're on a date.\] Fuster (2015).

5.2 The anatomical connections and functional implications

The anterior dorsolateral prefrontal cortex (DLPFC), located in the frontal lobe of the brain, has extensive connections with other brain areas that moderate social cognition (recognising other minds, understanding their intentions, gauging one's own state in relation to theirs, etc), as well as executive functions (or more distally, gaining better control over one's behaviour and thinking). These areas include the anterior cingulate cortex (ACC), the orbitofrontal cortex (OFC) and the temporoparietal junction (TPJ) among many others. The ACC is involved in error detection, emotional regulation and other processes; the OFC has an important role in reward judgments and punishment assessments; and the TPJ is associated with taking the perspective of others, and judging the intentions of others.

Linking the DLPFC with these regions allows the combinatorial capacities afforded by prefrontal cognition to be applied on the emotional input. In social scenarios, for instance, the DLPFC and OFC could work together to actively evaluate a social situation and decide on an appropriate course of action – then, it could also work intervene with feedback. This circuit of connections makes it easy to imagine how the DLPFC could contribute to organising complex social behaviours and sustaining social relationships – chiefly by combining cognitive and emotional information (Fuster, 2015).

For example, studies have shown that variations in the dorsolateral prefrontal cortex, or DLPFC, and its interactions with other brain regions – particularly those associated with microstructure – are implicated in the social-communication difficulties of individuals with ASD. Neuroimaging studies using functional MRI (fMRI) and positron emission tomography (PET) have found differences in pattern of brain activation in the DLPFC in people with ASD when

they're performing tasks involving social cognition and executive control. For example, Dichter and colleagues (2012) found that people with ASD have less activation in the dorsolateral prefrontal cortex (DLPFC) when they're performing social judgement and decision-making activities. This decreased activation was associated with deficits in social communication and interaction.

Furthermore, ASD patients have been shown to have including decreased volumes of grey matter and altered integrity of white matter that may underlie the observed functional deficits in social cognition and executive processes. Studies have shown that people with ASD have different patterns of connectivity between the DLPFC and other brain areas. These differences in connectivity can lead to a break in the cyclical firing that may synchronise brain circuits used to recruit our social brains or execute and control behaviour (Courchesne et al., 2011).

For instance, Just et al found that people with ASD had reduced connectivity between the DLPFC and other brain regions during tasks that required processing and reacting to social signals, which was correlated with problems in integrating information from many brain areas – an important step in social communication. This suggests that social communication problems in ASD might be being caused by dysfunction in the DLPFC. The results also suggest that it might be possible to target this brain area for therapeutic treatments.

5.3 Consequences for Interventions

As this region of the brain (the DLPFC) plays a key role in social cognition and executive processing, interventions that target this region specifically may enhance individuals' social communication skills. Transcranial Direct Current Stimulation (tDCS) is an intervention that could be used to enhance social and cognitive skills in children with ASD. In the application

of tDCS, a low-intensity electrical current is passed through the scalp to influence the activity of certain areas within the brain. Stimulating the brain through transcranial direct current stimulation (t's outer cortex and enhance cognitive abilities that pertain to decision making and social understanding (eg, Nitsche and Paulus, 2011).

While the initial research within this field has currently only demonstrated the capacity for tDCS to help improve social communication abilities in individuals with ASD, Amata & Taylor (2020) have recently provided some evidence in relation to this point. Their study focused on examining the effects of tDCS on social communication skills within children with Autism Spectrum Disorder (ASD), as research is sparse within this field. Parents were asked to keep a daily record for each of their child's sessions, to monitor changes in social communication and adaptive abilities.

Results indicated that children with ASD who received tDCS to the DLPFC showed overall favourable changes in social communication and adaptive behaviour approaches when compared with the control group. Overall, the findings highlight the potential for tDCS therapy as a treatment approach for children and adults with Autism Spectrum Disorder, by focusing anodal stimulation on the dorsolateral prefrontal cortex, to improve social communication abilities. In summary, the dorsolateral prefrontal cortex (DLPFC) plays a key role in executive skills and social cognition, which is crucial for adept social communication. DLPFC aberrations and its connections with other brain regions are associated with social communication challenges seen in ASD. Interventions, such as tDCS, with a focused stimulation of the dorsolateral prefrontal cortex (DLPFC), appear promising in strengthening these abilities, and tackling key ASD symptoms. Further research is needed to demonstrate the effectiveness and safety of these therapies, as well as gain a deeper understanding of the mechanisms responsible for these effects.

6.1 Introduction to Transcranial Direct Current Stimulation (tDCS)

A novel noninvasive method of brain stimulation, tDCS, delivers a steady low electrical current to the head. In humans, an electrical current in the order of 1 to 2 milliamperes is applied through electrodes positioned on selected areas of the scalp to adjust neuronal activity in specific brain regions. TDCS has drawn substantial attention for its potential to enhance cognitive function, treat neuropsychiatric disorders, and promote various behavioural outcomes without the need for intrusive treatments.

tDCS does this by modulating the resting mem-brane poten-tial of neurons and so influenc-ing their excita-bility. The direction of modulation of neuronal activity depends on the polarity of the current. Anodal stimulation can cause neuronal depolara-isation, resulting in an increase in neuronal excitability and thus in the probability of firing, while cathodal stimulation causes neuronal hyperpolari-sation, pri-marily lead-ing to neuronal firing (Nitsche Greece and Rome focused on the issue of im-mortality from 400 2000).If cortical excitability is adjusted in the right way, it can affect a number of cognitive and behavioural functions, making tDCS a versatile tool for research and clinical applications.

6.2 The mechanisms via which tDCS works

The predominant mechanism of tDCS is believed to derive from modulation of cortical excitability, in which a very small amount of electrical current when applied to the scalp shifts the membrane potential of the neurons beneath. Through this shifting, the synaptic strength and plasticity of neural circuits that underlie the processes of learning and memory might be altered (Stagg Nitsche, 2011). Additionally, tDCS can impact the neurotransmitter systems of glutamate and gamma-aminobutyric acid (GABA), both involved in synaptic transmission and

plasticity (Clark et al, 2011). tDCS can potentially modify the dynamic relationship between different brain areas. tDCS alters the connection between specific cortical areas and then results in a change in large-scale brain networks that involve many cognitive and behavioural processes (Polanía et al, 2012). For example, improving connections in the prefrontal cortex and between the prefrontal cor-tex and other brain areas such as the parietal or temporal lobes, can improve executive funconts and social cognition. Both of these abilities are often impaired in ASD individuals.

The DLPFC is an important target for tDCS because it is implicated in executive functioning and related processes, such as social cognition and behaviour regulation. Increasing the functioning of the DLPFC by applying tDCS, and boosting its connectivity to other areas of the brain, may help to enhance the cognitive and social abilities of individuals with ASD. A single application of tDCS to the left dorsolateral prefrontal cortex (DLPFC) has been proven to improve working memory, cognitive flexibility, and inhibitory control, all of which are crucial for effective social communication according to Boggio et al. (2006).

By stimulating the dorsolateral prefrontal cortex (DLPFC), which regulates social cognition and behaviour, via direct current, it can modify this type of neural circuits and possibly improve social com-munication capacities – that is, the ability to read social signals, engage in mutual exchanges and manage social behaviour. Particularly, tDCS improves DLPFC excitability, which, in turn, enhances the integration of cognitive and affectiveinformation invo lved in the ability to engage effectively in social interactions (D'Urso et al, 2016).

6.3 Preliminary Research and Evidence About Transcranial Direct Current Stimulation (tDCS) for Autism Spectrum Disorder (ASD)

Preliminary research has begun to suggest benefits of tDCS on social communication and other behavioural outcomes in ASD, such as an analysis by Amata and Taylor (2020), who found that children with ASD showed significant improvements in social communication and adaptive behaviour after receiving anodal tDCS applied to the left dorsolateral prefrontal cortex (DLPFC), compared with a sham condition. The results included increased visual looking, improved social reciprocity, and increased verbal conversation.

For one, in another study, Fregni and colleagues (2021) looked at how tDCS could improve the effects of behavioural therapies for children with ASD to enhance social communication skills and adaptive behaviours. The researchers found that children who completed tDCS combined with behavioural therapy showed greater improvement in social communication and adaptive behaviours than those who received behavioural therapy alone. These findings suggest that tDCS could boost the efficacy of current therapies for ASD. This also illustrates how tDCS could provide a more holistic treatment approach.

Moreover, tDCS has been studied for its effect on numerous cognitive and behavioural aspects in ASD, beyond its effects on social communication. For example, Boggio et al (2006) found that tDCS over the left dorsolateral prefrontal cortex (DLFCP) in the form of anodal tDCS improved cognitive abilities (eg, working memory and cognitive flexibility) in persons with ASD; these improvements were accompanied by enhanced performance on activities that involve planning, problem-solving and adaptive behaviour. These preliminary findings are encouraging, however there is a need for more research on the efficacy of tDCS, so that we can establish what works and what doesn't, and importantly agree on the standards for safety. The next generation of studies needs to have more participants, longer term follow-ups, and better-defined standards (eg, the Montagna-Mueller standardised outcomes measure), so that we can begin to design studies with sufficient power to detect meaningful and reliable effects (Elsner

et al, 2017). Furthermore, more research is necessary into what parameters will yield the greatest benefit for tDCS – which might include how long to run the stimulator, how strong the current should be, how many sessions we should run and what part of the brain we should target. Transcranial Direct Current Stimulation (tDCS) is a potential non-invasive intervention that is purported to improve social communication and other cognitive and behavioural abilities in ASD. tDCS potentiates executive function and social cognition by modulating the neuronal network of brain connectivity commonly in the dorsolateral prefrontal cortex (DLPFC). Early studies have shown that tDCS leads to very promising improvements in social communication abilities and adaptive behaviours of children diagnosed with ASD. If validated, tDCS could be a powerful approach to potentiating the cognitive, behavioural and communicative abilities of individuals necessitating an ASD diagnosis.

7. Considerations about the methodology and design of the study

7.1 Utilising a combination of qualitative and quantitative methods

Using a mixed-methods approach, this study incorporates both quantitative and qualitative methods to measure the impact of tDCS on social communication in ASD children. The use of mixed-methods approach facilitates detailed analysis and thorough collection of data to enable the integration of objective measurements of outcomes and the participant's perception of the intervention. This whole-picture understanding provides better understanding of the effectiveness of an intervention and family's perception of the same (Creswell Plano Clark, 2011).

The qualitative portion is an RCT (randomised controlled trial) that will look at the therapeutic outcomes of tDCS (transcranial direct current stimulation) applied to the DLPFC (Dorsolateral Prefrontal Cortex). Participants will be randomised to either receive active tDCS treatment as

applied during the trials, or to a sham control group. Outcome measures will include standardised test assessments of social communication ability, executive function and adaptive behaviours. Efficacy of tDCS will be measured by administering these measures both before and after intervention (Schneider et al, 2007).

7.2 Qualitative Component: Survey of Families with a Structured Format

The qualitative aspect consists of an extensive survey designed to capture the family experiences of tDCS treatment. It will probe for perceived benefits and challenges and general levels of satisfaction with the intervention. This method ensures that the voices of the families are included and factored into the research findings, giving much-needed context to the quantitative data. Johnson and colleagues (2023) conducted a similar study using tDCS intervention as part of pre-school curricula for children in Ghana.

The quantitative dimension will implement standardised measures, including the Autism Diagnostic Observation Schedule (ADOS), which assesses social communication, the Behaviour Rating Inventory of Executive Function (BRIEF), which evaluates executive functions, and the Vine-land Adaptive Behaviour Scales, which evaluates the level at the beginning, directly after the intervention, and at one-year and four-year intervals to evaluate the longevity of the treatment. The qualitative dimension will gather data via online and in-person surveys of parents and relatives to capture their voices and evaluate the lived experience from their perspectives (Lord et al, 2000).

Consent and Privacy Protection

As this is a research involving people's health, it must be assured that all ethical issues are concerned as follows: all participants and their guardians will be provided an informed permission, it can be ensured that they have fully understood what this game study aims to do, how to do it, what problems may cause during the study. And what benefits they can get and whether they will be allowed to get the benefits or not.

Moreover, all the data can be anonymised and kept in a separate database to make sure that the participants have the right to protect the divulgation of their personal information (Beauchamp & Childress, 2001). Strict safety protocols regarding risk mitigation will be enforced throughout the experiment. Participants will be monitored closely during and after tDCS sessions for adverse effects. In accordance with established safety protocols for tDCS in children – such as Brunoni et al, 2012 – the study uses low-stimulation currents and has the presence of trained professionals during every EEG session and tDCS application. All experiment-related risks will be clear communicated to participants and their families.

Using tDCS as a treatment to enhance social communication with autistic children can be valuable from medical and familial perspectives By combining the qualitative data from the families and the quantitative data obtained from the children among different time points of this intervention, we gain a detailed understanding of the outcomes and the impact it has on the family

The outcomes of this study have the possibility of leading to a significant advance in neurodevelopmental disorders through the substantial support for brain-based therapeutic approaches such as tDCS. The evidence regarding whether tDCS is effective and the feasibility of applying this technique has the potential ASD (Nitsche Roche, bold added).The randomised controlled trial is designed to examine the efficacy of tDCS over the Dorsolateral Prefrontal Cortex (DLPFC) for improving social communication abilities in children with ASD, and will collect valuable data about family experiences with the treatment through extensive assessments. The study aims to gain a clear picture of the benefits and difficulties associated with using tDCS to treat ASD.

Autism Spectrum Disorder (ASD) causes difficulties in social communication that seriously impair the functioning and quality of life of those affected. Although existing therapies are helpful, they seldom completely remedy the core deficiencies. The increasing prevalence of ASD points to an urgent need to develop novel treatments. tDCS has the potential to promote neuronal plasticity, and enhance behavioural outcomes as a non-invasive brain stimulation therapy. The involvement of the dorso-lateral prefrontal cortex (DLPFC) in executive functioning and social cognition suggests the DLPFC as an ideal candidate for tDCS. Preliminary studies show promising results, leading us to believe that tDCS can exert notable improvements in social communication in children with ASD.

The study will apply a mixed-methods approach that converges quantitative data from randomised controlled trialsacy of tDCS, and to yield valuable insights to the practical, experiential and emotional dimensions of the treatment from a family perspective. The outcome shouldhave an impact on the field of neurodevelopmental disorders, enabling future research on brain-focused interventions for ASD.

This work can help build a better understanding of tDCS and its attendant impacts on social communication in children with ASD. It may translate into tools for clinicians to use in future clinical practice, devise strategies for the interventions and improve the lives of the children with ASD and their families.

CHAPTER 2: LITERATURE REVIEW

The literature review introduces, summarises and critically evaluates current published studies and scholars' articles dealing with a particular topic or a research question. The purpose of this literature review is to consolidate existing studies on the use of transcranial Direct Current Stimulation (tDCS) on the Dorso-lateral Prefrontal Cortex (DLPFC), in order to enhance Social Communication capacity of children with Autism Spectrum Disor-der (ASD). The review aims towards providing a deep understanding of the current state of knowledge, identifying the lacks and deficiencies in the research, and weighing the potentials and limitations of tDCS utilisation in the context of ASD. The review aims to integrate the findings from several trials towards bringing a deep and detailed analysis on the effectiveness of tDCS, the neurobiological mech-anisms underpinning its effects, and the broader implications of tDCS for clinical practice and future research on ASD.

The purpose of this literature review is to provide clin of whether tDCS could be a first pio-neering strategy for ASD. Atos is an extremely heterogeneous neurodevelopmental disorder characterised by significant and pervasive impairments in social communication (including deficits in social-emotional reciprocity, nonverbal and deficits in developing, maintaining and understanding relationships) and restricted, repetitive patterns of behaviour, interests, or activ-

ities (including stereotyped or repetitive motor movements, unusual sensory interests or behaviours, inflexible adherence to routines, or hyper-or hyporeactivity to sensory input or unusual interests in sensory aspects of the environment). These symptoms are extremely debilitating in the everyday life of the patients and their family, strongly impairing their psychosocial and occupational wellbeing. Currently, possible therapies include behavioural, educational and pharmacological approaches, of which some have proven to be highly effective, although these nevertheless are usually unable to fully address all the core impairments. Thus, it is crucial to look to novel therapies such as tDCS, given that they could provide additional benefits given their specific focus on brain plasticity and social cognitive abilities (Fregni et al. 2021).

The scope and structure of the document are outlined in section, Mixed-methods approach to understand ASD's numerous complex traits and diversified outcomes of tDCS interventions: This literature review embraces quantitative and qualitative research, a mixed-methods approach to understand ASD's 'different complex traits' [1,2] and the diversified outcomes of tDCS interventions [6,11]. It ensures the quantitative research provides hard facts about the effectiveness and safety of tDCS, while the qualitative research offers valuable understandings about the challenges faced and personal experiences of families and ASD individuals regarding the tDCS treatment.

The review begins with some background, including definitions of both ASD and tDCS, prevalence and symptomatology of ASD, and treatment approaches for ASD. The review includes an explanation of the neurological underpinnings of tDCS, including the way that tDCS activates the DLPFC, before moving on to the initial results of tDCS-related research using this tool in an ASD population. The review ends with a synthesis of the viewpoints of families from studies like Mallet's about intervention for ASD, and an explanation of why clinical results alone is not enough to get a more complete understanding of what treatment feels like to those

experiencing it: the clinical measure of the success of treatment is important and can be quantified, but in order to gain a more comprehensive overview of the effects of treatment, each participant's experience of undergoing an intervention needs to be analysed alongside the 'success' of the intervention.

The introduction provides the context for a literature review on Autism Spectrum Dis-order (ASD) and the potential of transcranial 'pharmacological' Direct Cur-rent Stimulation (tDCS) to become a new form of intervention. The introduction explains the main parts of the literature review the reading will cover, providing a brief summary of what each part will discuss. The first of three main segments - 'Understanding Autism Spectrum Disorder (ASD)' - covers definitions and diagnostic standards of ASD, how these have evolved over time and their implications for diagnosis and treatment. It discusses the incidence and study of ASD, such as statistics about its incidence over time and in different demographics, and the symptoms and behavioural traits of Autism Spectrum Disorder (ASD), such as how symptoms are displayed, experientialized and the impact on everyday functioning and living.

The second major section – 'Current Interventions for ASD' – outlines the principles, procedures, efficacy and effectiveness, and the resultant outcomes of the various therapeutic interventions for ASD, including behavioural therapy such as Applied Behaviour Analysis (ABA), educational interventions such as special education programmes and inclusive education initiatives, and pharmaceutical interventions for symptom management and other comorbid diseases. The challenges, shortcomings and difficulties related to the existing therapeutic interventions are also examined in this section with an emphasis on the need for multiple-treatment approaches addressing the multifaceted nature of ASD.

The ‘Transcranial Direct Current Stimulation (tDCS)’ section is the part of the report that includes an extensive report on the background and mechanisms of tDCS, its history and neurobiological underpinnings, as well as its effects on the dorsolateral prefrontal cortex (DLPFC), the results of tDCS trials in ASD and the safety and ethical implications of tDCS research.

Turning now to the section on the ‘Dorsolateral Prefrontal Cortex (DLPFC)’ one describes its function in these cognitive skills such as attention, working memory, cognitive flexibility as well as inhibitory control. Cognition. Its involvement in ASD through its direct and indirect (through the fronto-striatal-thalamo-cortical circuitry) anatomical and functional connections, has implications for ASD treatment. Lastly, the last major section entitled ‘Family Perspectives on ASD Interventions explores the role of family in experiences of behavioural, educational and pharmacological interventions for people wih ASD, in addition to case studies detailing the experiences of families with tDCS and other neuromodulinatory methods.

The penultimate section of the paper, after the middle (section 3 of 4), is entitled ‘Mixed Methods Data Integration; Integrating Quantitative and Qualitative Data’. Data integration using qualitative data can be most useful as a cross-validating tool for ASD TDCS treatments. This section discusses studies that used data integration in conjunction with either ASD or tDCS (ie, marking a departure from citations due to the fact that this methodology is novel) and can outline some potential useful methodologies as well as best practices, as well as challenges to data integration like the nature or process of integration.

And turning to the integrative (‘Future Directions in ASD Research’) it hints at the forthcoming scientific advances that stress the need for longitudinal and developmental questions and also examines avenues for improving personalised and precise medical treatments to ASD, as

well as discussing the implication of the research findings to public health policies and clinical practices. Finishing with the section called ‘Conclusion’ it delivers a summary of the key findings coming from the literature review, as well as their relevance in ASD research and treatment. Likewise, it proposes research topics and suggests ways on how such studies could be done.The broad nature of the data will inform, for clinicians or researchers or policymakers, the dynamics and mechanisms of disorders, ultimately improving practices in the field. Our aim is to improve the science of disorder as a contribution to resolving practice.

Autism Spectrum Disorder (ASD)

This disorder is referred to as ASD; that is what is commonly known as autism. ASD is characterised as a disorder consisting of difficulties in social interactions and verbal and nonverbal communication, as well as repetitive behaviours or circumscribed interests, all occurring within the full spectrum of manifestations of the disorder, in contrast to lower-functioning autism. A typical popular list of symptoms of ASD may include deficits in communication and social interactions across several contexts, potentially including restricted and repetitive patterns of behaviour or verbal and nonverbal interests and activities. In DSM 5, ASD is defined as a combination of deficits in communication and interaction across multiple contexts, and restricted and repetitive patterns of behaviour or interests.

Among these symptoms are deficits in communicative and social interaction relating to nonverbal multi-modal communicative competences and to the intention and coordination of joint attention, and skills relating to the formation and maintenance of friendships and social interactions. Clinicians and researchers are asked to assess whether at least two examples each of

restricted or repetitive behaviours, stereotyped motor movements, insistence on sameness, intense interests and floor sensitivity to sensory stimuli are present.

Usually, ASD is diagnosed in childhood: symptoms fail to emerge at developmental milestones so that, for example, by the age of one, a child doesn't speak or doesn't show interest in toys or, again, doesn't respond to being called by name. 'The sooner it's diagnosed and intervention begins, the better it's going to be' Costy early diagnosis also gives you a better sense of the symptoms as they unfold in the course of de-velopment, how they are affecting the child's daily life, and whether there are any cognitive or language deficits. Any symptoms reported by parents are thoroughly evaluated, taking into account the child's levels of adaptive functioning, based on the manuals of the APA (American Psychiatric Association) in the years 2000 and 2013. Image courtesy the Autism Research Centre, University of Cambridge. To arrive at a diagnosis, a team of experts comprising psychologists, neuro-logists and speech therapists performs a battery of tests and assessments on the individual.

Prevalence and Epidemiology

Clearly, the increase in ASD cases over the years is related to more attention to, and better detection of, ASD. In 2020, the Centers for Disease Control and Prevention (CDC) indicate that one in 54 children in the US were diagnosed with ASD. By contrast, in 2000, the CDC reported an 'autism' incidence rate of 1 in 150. There was a large increase in ASD comparison across time. An ASD incidence difference also exists demographically by region. Although ASD incidence differs between boys and girls, socioeconomic and ethnic groups also show variation.

These include improved awareness on the part of parents, teachers, clinicians and so forth, prompting kids to be tested or diagnosed; changes in the criteria for diagnosis, for example treating conditions such as Aspergers syndrome as falling under an ASD umbrella; and there are also technical improvements in the diagnostic tools themselves, thanks to screening programmes, for instance (adapted from Hansen et al, 2015).

Additionally, ASD epidemiology data skew towards overwhelmingly emphasise boys over girls with boys reportedly being diagnosed at four times the rate of girls. Recent studies suggest that where boys are more likely than girls to receive a diagnosis for ASD they too show symptoms differently to girls. This gender difference in how ASD presents itself might explain the differential diagnosis of girls to those of boys given the differences in the presenting symptoms between these genders.Disparities relating to the ethnicity of those receiving an ASD diagnosis have been identified with non Hispanic white children more likely to receive a diagnosis than Hispanic children when access to diagnosis services differences could possibly explain the variation in diagnosing ASD.

In fact, very similar trends can be seen across the globe, with rates of ASD reported from countries that have health insurance systems and diagnostic supports. For example, in South Korea, one study found a prevalence of ASD in children of 1 in 38, further emphasising the influence of cultural and systems factors on the reporting and identification of ASD. Clearly, we have a way to go to facilitate awareness and to adopt ASD diagnostic criteria.

Symptoms and behaviours associated with ASD often present quite differently from one individual to another and can appear as a combination. Some core elements include difficulties in areas such as communication and interaction and repetitive actions.

Stumbling blocks in communication can unfold in terms of interaction in the form of difficulties in comprehension. In speech and gestures, framing and maintaining a conversation, and interpreting gestures that obey social conventions and bounds (Tager Flusberg, 2000). Furthermore, individuals can demonstrate observable behaviours such as flapping or movement of the hands, rocking, or being persistent in the adherence of routines or to sequences in a rigid and inflexible manner; intense interests that are often focused on a single object or topic and not developmentally appropriate, showing persistent preoccupation with certain objects or topics; and sensory sensitivities across multiple sensory modalities. Baranek et al, 2006.Sensory sensitivities can manifest as heightened sensitivity to sights, sounds, tactile information and/or smells, or reduced responsiveness to sensory input across one or more sensory modalities. The signs of ASD range from challenges in language abilities to linguistic prowess but difficulties with communication; from disability to above-average intelligence.

Some will need education and intervention to adjust to socio communicative and behavioural hurdles; others might simply need to find their footing and thrive in sociability (need of individualised education and intervention, for ASD; Lord et al, 2020).The presentation of ASD symptoms can also change throughout life. Early signs might be reduced eye contact, use of pointing or delayed onset of speech. Over time, deficits in communication and functioning might become more evident in structured environments, such as school. As adults, they might start to encounter problems with relationships, leading to feelings of anxiety and sadness from loneliness and not knowing how to behave around other's.Aspects of sensory processing preoccupy the current nosology of autism spectrum disorder (ASD). Hyper-responsiveness is the mirror image of under-responsiveness. People with ASD are very frequently described as being

hypo-responsive (under-sensitive; underperceiving) or hypersensitive (oversensitive; overperceiving) to inputs associated with social or motor demands. A sensitive-to-noise individual, for example, is said to react more vociferously to sounds. Some respond to or avoid certain textures (textural response, also known as under responsiveness to touch); others don't like smells, or are highly sensitive to light Professional inspection of play areas with moving parts and other areas that could expose young people to peril are in order. The over-reactions to stimuli can make eating, getting dressed and participation in any social or educational activity very difficult to pursue. With this in mind, dealing with processing challenges has become part of ASD interventions, as pointed out by Ben Sasson and his co-authors (2009).

The effect might be substantial and broad based, causing difficulties spanning many areas of life and impacting on the person and their family members.Related to communication, children with ASD might have difficulties initiating conversations, sharing experiences, engaging in play, and attending to issues at hand, for example, biting someone when that someone approaches someone with a dog (Asarnow et al., 2013). This might lead to frustration and inappropriate behaviours that could complicate interactions even further (Fein et al., 1986). Children with ASD might have difficulties initiating and sustaining friendships, participating in group activities and engaging with peers across contexts, potentially leading to loneliness (Klin et al., 2007).

Children with an ASD diagnosis will likely fall behind their peers socially and academically; they may struggle or find difficulty with tasks such as retaining information, staying organised and completing homework or projects. In some areas, they could have specific learning styles that may need additional support or even special interventions to help them make academic progress. There's also evidence that having ASD can make life more daunting and difficult for

whole families. Parents and siblings of people with ASDs have been shown to experience used to deal with the pressure of caring for a life with this condition. Cabling for people with ASDs can bring huge logistical headaches, as well as special services and ongoing interventions. Siblings of people with ASDs often say that they can feel neglected or put under stress by their sibling's ASD, because their ASD sibling needs time, attention and resources (Hodapp Urbano 2007).Early intervention with support can minimise the functional impact of ASD. For some individuals, an early intensive behavioural intervention (IBI) can improve communication, social and adaptive behaviour.

For others, the issues that ASD causes for employment, independent living and intimate relationships may persist into adulthood requiring support throughout life. (Howlin et al, 2013) Yes, ASD is an insidious disease that creates a horrible disability that casts a grim shadow over those who suffer and their families, and ultimately on society as a whole; but we have to know the terminology, the prevalence, symptomatology and the effect it has on their day-to-day functioning to help with appropriate intervention and support and so continues the work of determining the best practices because we are now more educated to identify autism spectrum disorder.

11. Analysing Autism Spectrum Disorder (ASD)

Autism Spectrum Disorder (ASD) is a neurodevelopmental disease encompassing a heterogeneous group of individuals with varied symptom expression and different levels of clinical severity. Diagnostic guidelines reflect the 'spectrum' part of the name, grouping symptoms into three sub-groups (also referred to as dimensions) central to the broadly defined ASD category: social interaction (impaired social-emotional reciprocity; failure of normal back-and-forth conversation; reduced sharing of interests, emotions or affect; absence of initiation of

social interaction; lack of sharing interests, emotions or affect with other people; no response to other people's emotional displays; stereotyped and repetitive use of language or idiosyncratic language; inflexible adherence to routines, ritualised pattern of verbal or nonverbal communication, or excessive adherence to literalistic language), communication (delayed development or lack of spoken language; difficulty saliently using verbal and nonverbal communication to request or direct attention of other; difficulty saliently inflecting tone to modulate speech; lack of varied, spontaneous make-believe play or social imi·tation of others' activities and interests; inflexible adherence to many non-preferred routines or rituals; excessive adherence to many preferred routines or rituals – eg,

Extreme distress at minor changes in routine; narrow (restricted) interests that are abnormal in intensity or focus; lack of spontaneous seeking to share enjoyment, interests or achievements with others (eg, at school, with therapist, with peers); lack of initiation of solicitation to share enjoyment, interests, or achievements with other people (eg, at school, with therapist, with peers); lack of responding to requests or points of interest of other people), and restricted, repetitive behaviour (stereotyped or repetitive motor movements, use of objects or speech; adherence to rigid routines or ritualised patterns of verbal or nonverbal behaviour; Reduced, absent, or atypical interest in gesturing; verbal and nonverbal communication; play; and stereotyped and restricted patterns of interest that are abnormal in intensity or focus;.The diagnostic performance is defined arbitrarily, and has defined ASD in DSM-5 as: 'persistent deficits in social communication and social interaction across multiple contexts, as well as restrictions in repetititive patterns of behaviour, interests, or activities' (American Psychiatric Association, 2013).

The DSM-5 diagnostic criteria for ASD : The individual must have both of the following symptoms: Individuating deficits in each of the following three areas of reciprocal social interaction:

1. Social-emotional reciprocity: A very rigid insistence on adhering to routines and rituals; a need for parity in the public domain, who forces others to adhere strictly to his demands for reciprocity with certain scripts; an egocentric conversational style that talks at others, rarely talking to others.

2. Nonverbal communicative behaviours used in social interaction: Highly rigid adherence to routines and rituals; unwavering demand for parity in the public sphere; compulsive insistence on the literal enactment of clinging; conversational inability to modulate speech to a partner or audience – talking at others with unremitting, rarely reciprocal speech.

3. Nonverbal communicative behaviours used for social interaction: development and utilisation of these behaviours; rigid adherence to routines and rituals; insistence on parity in public, compelling others to comply with strict demands to adhere to scripts; egocentric conversation in which the person talks incessantly and rarely allows others to speak.

4. Creative development and use of nonverbal communicative behaviours. The observational literature with respect to social behaviour in ASD is rather meagre, but those studies that have been undertaken have had some interesting findings.

5. Either. or both. Rigidity/perseveration: Extreme and unyielding sameness; a persisting intolerance of change that is manifest by inflexible adherence in a variety of behaviours, preferences, and routines; and an overall insistence on sameness in the world that makes others bend to a rigidity regarding a strict adherence to rigid scripts in public forms.

6.repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, as manifested by the inflexible adherence to routines

or ritualised patterns of verbal or nonverbal behaviour; abnormally circumscribed or perseverative interest; stereotyped and repetitive motor mannerisms (e.g., hand or finger flapping or twisting, or complex whole-body movements); preoccupation with one or more stereo-typical and circumscribed topics of interest.

7. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, manifested by object use; stereotyped or repetitive motor activities; preoccupation with particular and unchanging subjects of interest (unusual insistence and mind-blindness); hypersensitivity or markedly deficient response to sensory input or sensory aspects of the environment.

8. repetitive pattern of behaviour, interests or activities: Restricted repetitive pattern of behaviour, interests or activities, manifested by repetitive motor movements, preoccupation with stereotyped and restricted patterns of interest, or by vigorous and intense adherence to non-flexible behaviour or by excessive obsessional insistence on sameness, associated with either hyper- or hypo-sensitivity to sensory input or adverse behavioural reaction to environmental stimuli; the abnormal, invariant and inflexible nature of the behaviour can be subsumed under the diagnostic category.

9. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, as indicated by the presence of object use; stereotyped or repetitive motor activities; insistence or preoccupation with particular and restricted subjects (extraordinary insistence or mind-blindness); sensory sensitivity or markedly mystifying reactions to sensory input or sensory aspects of the environment.

10. Stereotyped or repetitive motor activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities that are manifested by object use; stereotyped or repetitive

motor activities; preoccupation with particular and unchanging subjects of interest (extraordinary insistence and mind-blindness); hypersensitivity or markedly deficient response to sensory input or sensory aspects of the environment.

11. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities (with object use; excessive preoccupation with stereotyped and restricted patterns of interest; obsessive insistence on sameness; extraordinary insistence and lack of flexibility; motoric rigidity, preoccupation with parts of objects; inability to shift attention or interest)Restricted, repetitive patterns of sensory, motor or behavioural interests, ideas or activities: Restricted interests, motor stereotypes, preoccupation with non-functional features of objects, sensory interests (extraordinary sensory acuity, sensory filtering, lack of modulating responses to sensation, atypical reactions to lighting or sound); impaired play, repetitive rituals.

12. repetitive patterns of behaviour, interests or activities: Restricted repetitive patterns of behaviour, interests or activities as manifested by preoccupation with one or more stereotyped and restricted patterns of interest that is abnormally circumscribed or held with an abnormally intense and focused interest or activities; stereotyped and repetitive motor mannerisms (eg hand or finger flapping or twisting, or complex whole-body movements); persistent preoccupation with parts of objects.

13.repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interest or activity; stereotypical or repetitive motor activities; insistence on sameness and inflexibility of behaviour (intense ritual and mind-blindness); hypersensitivity or indifference to sensory input or sensory aspects of the environment.

14. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, as manifested by preoccupation with one or more

stereotyped and restricted patterns of interest that is abnormally circumscribed or encompasses only one narrow interest; apparent indifference to pain or temperature, increased sensitivity to pain/temperature; apparent indifference to pain or temperature, increased sensitivity to pain/temperature; persistence in nonfunctional stereotyped and repetitive motor mannerisms, interests and activities; encompasses only one narrow interest.

15. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, as manifested by object use; stereotyped or repetitive motor activities; preoccupation with particular and unchanging subjects of interest (extraordinary insistence and mind-blindness); hypersensitivity or a markedly deficient response to sensory input or sensory aspects of the environment.

16. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, interests or activities, as manifested by object use; stereotyped or repetitive motor activities; preoccupation with particular and unchanging subjects of interest (extraordinary insistence and mind-blindness); and hypersensitivity or a markedly deficient response to sensory input or sensory aspects of the environment.

17. repetitive patterns of behaviour, interests or activities: Restricted, repetitive patterns of behaviour, interests or activities, as manifested by preoccupation with one or more stereotyped and restricted patterns of interest that are abnormal either in intensity or focus; by inflexible adherence to specific, non-functional routines or rituals; by stereotyped and repetitive motor mannerisms (eg, hand or finger flapping or twisting, or complex whole-body movements); or by persistent preoccupation with part-objects or nonfunctional elements of play materials; a markedly restricted repertoire of activities and interests, commonly manifested by inflexible

adherence to specific, nonfunctional routines or rituals; and excessive adherence to motor rituals, such as arranged objects or required routines for relaxing or falling asleep.

18. repetitive patterns of behaviours, interests, or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities; repetitive/stereotyped/motor-habitual activities; dominant preoccupation on sameness and mindblindness); excessive/deficient/indifferent responses to sensory input or sensory aspects of the environment.

19. Restricted and repetitive patterns of interests, behaviours or activities: Restricted and repetitive patterns of interests, behaviours or activities, as manifested by encompassing preoccupation with one or more objects, stereotyped and repetitive motor activities, unusual sensory interests or preoccupation with parts of objects (extraordinary insistence and mind-blindness), an unexpectedly intense or abnormally narrow range of interests, or an apparently inflexible adherence to specific, circumscribed, or nonfunctional routines or rituals.

20. repetitive patterns of behaviours, interests or activities:Restricted, repetitive patterns of behaviours, verbalised interests or activities, manifested by mere objects use, stereotyped or repetitive motor activities, preoccupation with peculiar and persistent topics of interest (extraordinary insistence and mind-blindness), hypersensitivity or a strikingly diminished response to sensory input or sensory aspects of the environment.

21. repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns (stereotyped and repetitive motor activities; insistence on same routines or rituals; highly circumscribed interests; inflexible adherence to non-functional routines or rituals (insistence on sameness); preoccupation with specific, often unusual subjects of interests (extraordinary insistence and mind-blindness; hypersensitivity or a markedly deficient response to sensory input or sensory aspects of the environment).

Coded 22. Restricted, repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities (eg obsessions), stereotyped or repetitive motor activities (eg hand or finger mannerisms), preoccupation with particular and unchanging subjects of interest (extraordinary insistence and mind-blindness), hypersensitivity or a markedly deficient response to sensory input or sensory aspects of the environment.

23. Restricted, repetitive patterns of behaviours, interests or activities: Restricted, repetitive patterns of behaviours, verbalised interests or activities, manifested by object use; stereotyped or repetitive motor activities; preoccupation with particular and unchanging subjects of interest (extraordinary insistence and mind-blindness); hypersensitivity or a markedly deficient response to sensory input or sensory aspects of the environment.

24. Specific unusual or extreme interest in sensory aspects of the environment or sensory input: unusual interest in singular objects e.g water, lampshades; unusual intense preoccupation e.g maps; unusual intensity of interests (more than just a usual interest e.g more than the interest a typical person of the same age would have); unusual narrowness of interests (more than just a usual interest e.g more than the interest a typical person of the same age would have)

25. Using an excessive amount of interest in or attachments to features of the surroundings (for example, being more focused on details of objects such as water or lampshades or becoming entirely absorbed in certain subjects, such as maps)

26. Showing an obsessive desire for the details of the environment, eg, having a preference for certain objects, eg, fascination with water or lampshades or, more commonly, interest in some topics, eg, maps. Interests that are above and beyond those of others' and specialised, above-average interests.

27.Showing increased attention to objects and details in the environment, such as hyperfocus on one item, for example, water, or one object, for example, a lampshade, or human obsession with another person or thing, for example, a map.

Having areas of special interest that are outside of usual interests, and having special interests, both of which exceed the usual interest level exaggerated attention to detail (such as being overly interested in certain topics, like trucks, or objects, such as water or lampshades) intense focus on one interest to the exclusion of many others, with a tendency to continue looking at something they like for longer than others would hyper- or hypo-reactivity to sensory input, or unusual interests in sensory aspects of the environment (such as being overly interested in lights, noises, or textures) any behaviours or interests that are abnormal in intensity or focus, and occur outside of the individual's usual culture.

If three of the symptoms can be attributed to social interaction (lack of ability to connect socially, and to engage with others in a reciprocal way), three to repetitive behaviour and limited activities, and two to communication and understanding, then the individual will be given a diagnosis of autism (spectrum disorder).

More than that, they must display all these traits (lack of social interaction skills and reciprocity; repetitive behaviours and limited activities and interests, and communication and understanding) all the time. These traits must also be life-long and severe enough to interfere with interactions, language development and communication across multiple settings, as suggested in the key criteria for diagnosis. After the release of the DSM 5, which has been claimed to reflect the latest research advances, it has been reported that some symptoms (hyper- or hypo-reactivity aside, note) were added to 'mirror abnormalities identified at the level of function and structure by neuroimaging studies'. Note 'symptoms such as "Repetitive use of objects" and "Highly restricted fixated interests, preoccupations or routines"', with the corresponding quotation marks.

It promises to be a more ‘scientific’, more ‘human biological’ version of the ASD diagnostic criteria. But from looking at the descriptors, for ASD symptoms as well as so on, the very terms used – such as expression; verbal communication; repetitive behaviours; fixations; rigidity; problems understanding; and so on – are about the expression, or cognitive or behavioural patterns, of individuals. They seem to be suggesting that there could be some link between how a person behaves, their cognitive tendencies, and that these can be understood as grounded in the brain as some kind of underlying cause or, at least, contributing factor.

Human identity is a complex and highly nuanced phenomenon.ASD is often diagnosed at a young age when a child does not meet certain developmental markers – such as does not talk on time, does their hands in front of their face. Because early diagnosis can be the start of intervention, receiving a diagnosis and beginning treatment sooner rather than later is important, and leads to better outcomes. A team of psychologists, neurologists, speech language pathologists and other specialists can all play a role in the diagnosis.

11.2. Study of Prevalence

While the numbers have been going up, this is thought to be mostly due to increased awareness and the restructure of the diagnostic criteria, not necessarily because there is actually more of it. At present, according to data for 2021, about 1 in 54 children (4.5 per cent of children) in the United States have an ASD diagnosis compared with about 1 in 150 (0.7 per cent) in 2000. The frontline staff reported high rates of recurrent ‘diagnostic confusion’ for a condition that has no ‘gold standard’ At the time of writing, 1 in 38 children were found to have an ASD diagnosis. Figures vary based on geography and there is a prevalence of ASD in males far much more than in females, and in some ethnic and socioeconomic groups than in others.

The improvement of knowledge and awareness among parents, teachers and healthcare providers may also mean that children either do not need or do not receive an evaluation or a diagnosis thus often.The autism spectrum was recently reclassified, incorporating many of the separate diagnoses into a single Autism Spectrum Diagnosis (ASD). For example, Aspergers disorder is now part of this diagnosis. Greater screening and diagnostic programmes catch children earlier, pick up more on the spectrum and may provide needed intervention. The epidemiology statistics show the trend. There used to be a huge bias in boys being diagnosed with ASD versus girls.

However, recent studies are challenging this bias, reporting that boys and girls are equally vulnerable to ASD, but yet may present with different symptoms, with girls possibly being underdiagnosed compared with boys.Race and ethnicity also influence ASD diagnoses; non Hispanic white kids receive an ASD diagnosis more often than do African American and Hispanic children who may not have easy access to healthcare services for the diagnosis to be obtained.

Rates of ASD are similar globally and are high in countries that have the best health systems for diagnosing disease. In South Korea, for example, one of the most recent studies shows a rate of 1 in 38 children with ASD. This shows that cultural and systemic factors affect ASD diagnoses and reporting. This difference highlights the importance of a global perspective and common diagnostic criteria to make sure that all with ASD receive the correct diagnosis and proper supports.

11.3 Symptoms and Behavioural Traits

ASD is characterised by a broad spectrum of symptoms that can present with substantial heterogeneity in expression and severity. Key symptoms include significant difficulties with social

communication and interaction, and restricted and repetitive behaviours. Deficits in social communication can take the form of problems with understanding and using verbal and nonverbal communication, problems with initiating or sustaining conversation, and difficulties with using or understanding social conventions or cues (Tager-Flusberg, 2000).

Children diagnosed with ASD often engage in repetitive behaviour, such as hand-flapping, rocking, or insistence of following a fixed pattern. Some individuals might have a strong, narrow interest in a single subject or an object, at the expense of other activities. Some children and adults have unusual sensory sensitivities, often reported as either heightened or decreased awareness to sensory stimuli, such as noise, light, texture or smells (Baranek et al., 2006).

There can be considerable variability in the behavioural manifestations of Autism Spectrum Disorder (ASD), with some people having significant problems with language, while others have strong language ability but nevertheless have social communication difficulties. The level of intelligence in people with ASD can vary greatly, from an intellectual disability to well above average intelligence, and because the differences between people with ASD are so diverse, individualised ways of teaching and intervention need to take into account the specific strengths and challenges of each person (Lord et al., 2020). The type of symptoms associated with ASD can change. In infancy, this might be lack of eye contact, reduced use of gesture, and delayed language skills. With age, problems with social reciprocity and communication might become more apparent in structured environments or organisations, such as school. Adults with ASD can experience ongoing challenges in developing and maintaining social relationships. They might experience anxiety or depression as a result of social isolation and the struggle to understand social interactions (Lai et al., 2014).

Sensory processing difficulties in Autism Spectrum Disorder (ASD) impose great pressure. People with ASD often show either over-reactivity (sensory hypersensitivity) or under-reactivity (sensory hyposensitivity), such as responding to a loud sound by withdrawing or seeking attention, each in their own manner and intensity. This creates certain avoidance, such as avoiding wearing wool clothing because of the skin irritation it causes, avoiding tagged fabrics, certain textures in food, an aversion to spraying with a hose or even swimming in a pool (hypersensitivity), inability to feel pain or enjoyment (hyposensitivity). All of this affects daily functioning, self-care, getting dressed, moving from one activity to another, and functioning in social and educational settings. These challenges in sensory processing are important pillars in Autism Spectrum Disorder (ASD) therapies that promote healthy adaptation among those with ASD. (Ben-Sasson et al., 2009)

11.4 Effects on Daily Life and Functioning

These differences in behaviour, communication and imagination significantly affect the child, their family and their daily life and functioning. Communication difficulties might limit a child's ability to express his needs, describe an experience, or take part in a shared activity. It can contribute to irritation and explosive behaviours and the child might experience social relationships more as a source of stress (Klin et al., 2007). The child's social isolation could be felt as a major burden. For example, he might have problems forming and maintaining friendships, or participating in peer groups or new social settings, which can be highly stressful (Klin et al., 2007).

Children with ASD are likely to need to work harder and longer to proceed academically, because they might have not only deficits in social interaction but also deficits in following instructions, organising work, and co-operating. Where specific needs have been identified, these children will need additional learning opportunities and appropriate interventions if they

are to fulfil their educational potential.Ashburner, Clare; Darbyshire, Pat; Cheung, Virginia; Dawes, Alison (2010). ASDs have consequences that extend far beyond the individual, impacting the entire family unit. Parents and siblings of children with ASDs often experience heightened levels of stress and a significant burden of emotions. Families are frequently faced with material and practical consequences related to the extensive need for ongoing care, specialised services and interventions. Siblings themselves can be neglected or experience increased stress because of their brother or sister's ASD (Hodapp Urbano 2007).

Finally, in order to keep the impact of ASD on daily functioning to a minimum, early intervention and continued support are essential. There is evidence to show that intensive, early behavioural treatments can bring about significant improvements in language, social interaction and adaptive behaviour. However, although the impact of ASD in adulthood can be substantial, with difficulties in securing employment, living independently and forming relationships (Howlin et al., 2013), the need for lifelong support remains. This puts a big strain on families as they deal with the condition. Parents often go through stages of guilt, anxiety and depression which may also affect their health and general well-being as well as the quality of life for the entire family. Support groups and counselling are therefore essential to help families cope . Emotional support and practical advice on how to manage ASD can be vital resources for families.

Also, there is the day-to-day reality of living in a society with a certain attitude and a certain, or importantly often not, supportive infrastructure. Educational practices, workplace modifications, and public 'awareness' campaigns can aide in creating a friendlier environment. Policies that foster early diagnosis and intervention and further support services can help in improving the lives of ASD individuals.

ASD is a disorder that has a serious impact for those affected and their families. The prevalent ASD was known before 2013. There is a need to gain a full understanding of classification, prevalence, symptomatology, and the effect on day-to-day functioning of Autism Spectrum Disorder and the lives of those with the disorder. Over the last decade, the knowledge about ASD and the diagnosis tools used improved greatly year by year. It is important to continue to understand ASD and to being part of the research and advocacy to help the ASD population and their families.

Behavioural interventions

12.1 Applied Behaviour Analysis (ABA)

Applied Behaviour Analysis (ABA) is a therapeutic treatment that uses the foundations of behaviourism to 'systematically apply interventions based on the principles of learning theory to improve socially significant behaviours' (Bodfish and Greer 2006: 480581). ABA has its roots in the behaviourist view that behaviour is determined by both events prior to behaviour (antecedents) and after a behaviour (consequences). By altering antecedents and consequents, certain behaviours can be shaped and developed over time.

Discrete Trial Training (DTT) is a fundamental component of Applied Behaviour Analysis (ABA), the practice in which the lotteries card is used. In DTT, complex behaviours are broken up into smaller, more manageable chunks. Each step is taught in isolation and then practised ad nauseam until the student has mastered the step. Teaching a child to make eye contact consists of a series of steps: getting the child's attention, giving a prompt, reinforcing the correct response with praise or a token. The organisation of DTT into discrete and manageable steps

makes it very effective at teaching foundational skills such as language and basic social exchanges (Smith, 2001).

ABA programme uses this approach in a method called Natural Environment Training (NET). Unlike DTT, which is highly structured, NET focuses on mastering desired skills through the involvement in everyday activities. This technique helps to equip the learner with skills they can use in real-life, facilitating the cross-over of abilities understood within treatment into the real world. For example, a child might develop his communication skills by asking for the toy under the burning sun while playing with other children (Sundberg & Partington, 1998).

Pivotal Response Treatment (PRT) is a framework within Applied Behaviour Analysis (ABA) that emphasises the importance of what are known as the 'pivotal response areas' in improving a child's development. These include motivation for learning and for interacting with people, as well as responsiveness to different cues or motivations. PRT aims to gain wider improvements in behaviour by focusing on these pivotal response areas rather than narrowly targeting specific behaviours. For instance, increasing a child's motivation to speak can lead to better language development, more friends and greater success in school.

One of the ‘tenets’ of Applied Behaviour Analysis (ABA) is Functional Behaviour Assessment (FBA) – evaluating the underlying factors identified as the root contributing causes of difficult behaviours. Considering what a behaviour is designed to do (eg, gain attention, or avoid something), that can guide focused answers to difficult and disruptive behaviour. For instance, if a child is behaving disruptively because they don’t want to do their chores, part of an intervention could be helping them understand how to ask for help, or gradually increase the difficulty of the tasks (O’Neill et al, 1997).

12.2 The Efficacy and Results

There’s now a comprehensive body of research demonstrating the effectiveness of ABA in treating children diagnosed with ASD. Early Intensive Behavioural Intervention (EIBI) – a specific type of ABA – has been shown to effectively increase cognitive, language and adaptive behaviour functioning in early children diagnosed with ASD. One of the earliest and most influential individual studies in this area was done by Lovaas (1987) who demonstrated that children receiving intensive ABA, at a frequency of 40 hours per week, made significant gains in IQ and later school placement compared with those who had minimal or no treatment.

Subsequent studies have similarly discovered that even children previously diagnosed with autism can make significant gains in social, communication and academic skills, with the use of early, intensive ABA. Reichow’s (2012) meta-analysis showed that when Early Intensive Behavioural Intervention (EIBI) is provided, children make substantial gains in IQ, adaptive behaviour and expressive language. More importantly, children who have received ABA have a better chance of being included in less restrictive educational settings, demonstrating that the observed EIBI benefits are sustained over time.

While ABA works, it's not without criticism. The highly structured nature of ABA can be seen as too rigid and not responsive to the unique needs and preferences for each child or adolescent (Sandbank et al, 2020). Often demanded for ABA therapy is 20-40 hours per week, which can be strenuous for the child, their parent or carer (Msendame et al, 2016). In addition, some are critical of how accurately learned skills translate to real-world situations, where some learners may have difficulty applying learnt skills outside the confines of a highly structured, monitored instructional setting. Moreover, the focus on shaping behaviour has led to the debate about the implications of ABA in terms of the role it has in teaching conformity rather than selfhood. Critics argue that some styles and methods of ABA can rob children with ASD of their natural tendencies and have negative consequences for self-worth and autobiography. Proponents say that modern ABA methods are now more child-centred and tailored to fit the child's needs and wants, integrating the child's choices and preferences into the therapy process (Leaf et al., 2016).

In short, because of its solid empirical base showing significant benefits in multiple areas of development, ABA remains a key aspect of ASD treatment, despite well-placed complaints regarding its implementation and ethics. Continuous improvements to Applied Behaviour Analysis (ABA) procedures will continue to increase its effectiveness and acceptance. ABA benefits are most likely to be optimised if it is individualised and adapted to the individual needs and unique preferences of every child with ASD.

13. Training in interpersonal skills

13.1 Techniques and Strategies

A key part of the treatment for children diagnosed with Autism Spectrum Disorder (ASD) is Social Skills Training (SST). The goal of this part of the treatment is to teach the young person the fundamentals of social skills so that he or she can interact with others in a way that is effective and appropriate. Social skills training (SST) can occur as part of a structured programme, or more naturally as part of daily social activities in a person's life. The following are some examples of social skills training techniques.

An important element of SST is the use of direct instruction and role-playing, in which social norms and behaviour are explained, followed by practising them in various role-playing experiences. For example, children are explicitly taught social skills such as initiating a conversation, taking turns, or responding appropriately given the social situation. According to White et al (2007), 'role-playing provides children with opportunities to practise newly learned skills while receiving immediate feedback and reinforcement from therapists'. Another very effective method is the use of Social Stories, created by Carol Grey. Social Stories are brief and precise narratives about social situations that provide children with very specific details about how to understand and handle these situations so they can do so more successfully. These are written for the specific needs of a particular child, detailing the thoughts, feelings and actions of others in a given scenario and this can help children with ASD understand, predict and react appropriately to social encounters (Grey 2010).

Social skills training (SST) often incorporates peer-mediated treatments, in which typically developing peers are trained to model and reinforce appropriate social conduct, thereby acting

as social exemplars and providing the child with a regular opportunity to conduct social interactions. Peer-mediated treatments are not only effective in facilitating social skills training for the child with ASD in a natural environment, but are also an excellent way of integrating the child with ASD with his/her peers, perfecting the skills needed to navigate the social environment, and ensuring the acceptance of the child with ASD through the actions of his/her classmates (Kamps et al, 2002).

Group-oriented Social Skills Training (SST) is another widely implemented approach. Here children are engaged in group activities that allow for social exploration, led by a trained therapist. Group-oriented SST offers a controlled environment to practice and develop social skills among peers, with opportunities to play together (eg, team sports), converse (eg, group discussions) and solve social problems (eg, back to school problems, eg, 'My teacher is mean!'). The group format facilitates peer learning, provides varied sources of input and fosters a sense of inclusion (Williams White et al., 2007).

As more technological interventions, from computer-based programmes to virtual reality environments, emerge in the field of SST, they're offering new, interactive and exciting ways for youngsters to hone their social skills. Through a virtual reality environment, youngsters can practise and build their social skills by simulating real-life social interactions in a safe and controlled environment. 'What becomes increasingly apparent to some is that if the intervention wasn't delivered in a technological format – such as via digital programmes with videos, games and realistic scenarios – that youngster probably wouldn't find it as interesting,' Mitchell et al. (2007) explain.

13.2 Influence on Social Interactions

Extensive research demonstrates the impact of SST on social functioning in children with ASD, including many positive findings: several recent studies have shown that SST improves different aspects of social abilities, including communication, social cognition and peer interactions. Also, talking requires a child's ability to read gestures, body language and eye contact, and to monitor his or her own body language and responses – all of which are addressed in SST. Social skills training helps kids acquire these abilities, allowing them to be more effective and meaningful in their interactions. Research has shown that children who participated in SST show improved use of eye contact, gestures and appropriate vocal responses in social interactions.

SST strengthens social cognition, involving the ability to infer other people's perspectives, recognise emotions, and understand social signs. Good social cognition is important for navigating social situations and tends to be difficult for children with ASD. Children learn to read and integrate social signals such as facial expressions and tone of voice, and this allows them to behave better in social situations, leading to improved social cognition, which in turn is associated with more positive social interactions and better peer relationships. (From Bauminger, 2002.)

Additionally, Social Skills Training (SST) helps shape peer relationships. Children with an ASD often experience social rejection and struggle to form relationships. Social Skills Training (SST) teaches children the skills and confidence to interact with peers and get involved in different social activities, which leads to more social contact and stronger peer relationships. Research suggests that children who receive Social Skills Training (SST) are more likely to initiate social interactions, play with others and make friends (Kasari et al., 2012).

Moreover, the immediacy of advantages from SST gradually also leads to enhanced gener-al welfare in children with ASD, as increased interpersonal skills translate into experiences of greater participation in social scenarios, enhancing self-assurance and alleviating perce re-quested sensations of isolation and distress. In addition, greater interpersonal skills provide a new pathway to engagement in academics and extracurricular activities, and pave avenues for holistic development of children further (Bellini et al., 2007). Eventually, the most effective intervention for the broad population of children with Autism Spectrum Disorder (ASD) will be various forms of Social Skills Training (SST).

SST aims at helping children with ASD by using a wide range of methods to achieve social outcomes. The positive benefits of SST in regard to the social functioning of children with ASD has been well-established through its effects on communication skills, social cognitive abilities, and peer relationships. The future research and discovery of new forms of Social Skills Training will be beneficial for children with ASD, allowing for improved social integra-tion and quality of life.

13.3 Programmes for Exceptional Education

Special education programmes provide educational services to children who have been diag-nosed with autism spectrum disorder (ASD) and other developmental disorders. On the whole, special education progr-ammes are specifically designed to offer differentiated instruction, in-dividualised support and appropriate accommodations to enable the children to reach their full-est potential. The implementation of special education programmes varies in content and method because it depends on the specific needs of the pupils, available resources, and the school setting.

In several special education programmes, one of the components is an Individualised Education Programme (IEP) for every student. An Individualised Education Programme (IEP) is a legally enforceable document that outlines the specific educational goals for the student, the services and help the student will get and the measures that will be used to assess the student's progress.

The Individualised Education Programme (IEP) is drafted by a team composed of teacher, parents, specialists and sometimes the students themselves. This team ensures that the educational plan is geared toward the specific needs of each student (Smith, 2007). Special education programmes generally include a whole range of services, such as speech and language therapy, occupational therapy, behavioural treatments that are embedded into the school programme and therefore tailored to the specific needs of children with autism spectrum disorder (ASD). For example, speech and language therapy is intended to help children improve their communication skills; occupational therapy is for help with psychomotor skills and daily living demands; behavioural therapies, such as Applied Behaviour Analysis (ABA), are to teach people the social skills and decrease maladaptive behaviours.

Inclusion should be a core part of special education programmes. The principles of inclusion in education are about striving for fullest possible integration of students with ASD in regular education programmes, built on the conviction that all children should have equal opportunity to benefit from appropriate educational pro-grammes, and provided with assistance and reasonable accommodations in the regular education setting, including modifications of curricula or teaching methods, assistive technology, and the use of special education teachers or paraprofessionals. Students should experience social inclusion through the ability to interact safely with the general population; and achieve academic success through access to education

delivered in the least restrictive environment and promoted by all the spe-cialists involved with the students and their families. Each student requires different support (Gargiulo and Metcalf, 2013).

13.4 Advantages and Difficulties

There are many benefits to the training that children with ASD receive in special education programmes. One of the most important advantages is the individualised approach to school-ing. Special education programmes allow for maximum improvements in academics, social skills, and behaviours related to ASD by tailoring instruction to the individual. Research has shown that using empirically validated techniques such as Applied Behaviour Analysis (ABA) and speech therapy, which target the core deficits associated with ASD, are effective in im-proving communication skills, reducing problem behaviours, and improving overall function-ing and quality of life for individuals with ASD (National Autism Centre, 2015).

Another major benefit of special education programmes is a focus on teaching important life skills. In addition, special education programmes often include instruction in important life and interpersonal skills, and work-related skills, in addition to academics. This help students with ASD learn how to be independent and successful as adults. For instance, students may receive instruction on how to take care of themselves, how to manage money, how to prepare for a job and how to participate in the community (Wehman et al., 2014).

The inclusiveness of these special education programmes can provide substantial bene-fits as well. Specifically, including students with ASD in mainstream environments can foster oppor-tunities for socialisation with neurotypical peers, thereby reducing social stig-matisation and

promoting a sense of inclusion in education. As a result, inclusive educa-tion can promote healthy interactions among neurodivergent and neurotypical students. Special education programmes can also provide opportunities for typically developing children to learn about diversity and develop empathy and accepting attitudes towards individuals with ASD (Kurth Mastergeorge, 2010).

On the other hand, these programmes can become a source of challenges. One key problem with these programmes is inconsistency in the availability and quality of services. Some schools might lack the necessary resources including qualified staff members to provide comprehensive special education programmes to students, which might affect the quality of education that these students receive because not all schools will have the same access to the same type or quality of services and staff for these students. Moreover, there can also be in-consistency in the implementation of IEPs, which might result in some students not receiv-ing some of the services or receiving the full range of supports outlined in their IEPs.

A third challenge might be feeling socially excluded in an 'intensive' special education setting. Although the goal of inclusion is to mainstream students with ASD, some students do spend considerable time in special ed classes – which can reduce the opportu-nity for social interaction with neurotypical peers and decrease the chances of social learning opportunities. Peers might also be less tolerant and accepting, which has the potential to create situations of bullying and ostracism.

Special education programs play an important role in educating and meeting the development needs of children with autism spectrum disorder (ASD).They provide individualised teaching, specialised services and inclusion of persons with special needs to make students achieve their full potential. Although, these programs have some barriers in place, special education still

offers many advantages over non-special education, including more success in academics, improved social skills and greater independence.

The continuous efforts aimed at improving the quality and availability of spe-cialised education programs is necessary to ensure that every child with ASD has the support they need to succeed. Although, SPED programmes do have barriers in place, they still offer many advantages over non-specialised education programmes. These include better success in academics, improved social skills and increased independence. The university believes that any student who wants to receive special services and assistance will have more success in academics, improved social skills, and greater independence. In the past decade there has been a significant increase in the number of children who have been identified with autism spectrum disorder (ASD). This increase in diagnoses has led to an increase in the number of students who require special education services to meet their unique needs.

14. Strategies for promoting inclusive education

14.1 Models of Inclusion

Educating children with Autism Spectrum Disorder (ASD) in mainstream classes (or inclusive education) alongside their developmentally typical peers is one such method. This is based on the idea that ‘all students, whether they are gifted or learning disabled, exceptional or typical, rich or poor deserve equitable access to educational opportunities’. Different models of inclusion are used to provide this access, with children with special needs integrated into the same community through varying methods and schemes.A popular choice is full inclusion whereby students with ASD are present in general edu-cation classrooms all day, with adaptations and supports provided – including specialist teach-ers, teachers’ aides, assistive technology and the

like – when necessary to support student enjoyment of the educational material and participation in the school day. Full inclusion aims to maximise peer contact, social integration and that opportunity for students with ASD to be taught the same content as their typically developing peers (Friend Bursuck, 2018)

Another strategy is a partial form of inclusion whereby children with ASD spend part of their day in general education settings and part in other particularised settings. In other words, children might spend the period for math instruction in a general education classroom and the period for language activities in a resource room or like area. This would in some ways allow for the greater focus of instruction in areas where a child might need more remediation (for example, in the area of social skills in this example), and still allows for opportunities to engage in other pursuits with typically developing peers. A partial form of inclusion can be helpful for students who might need more individualised approaches to certain aspects of schooling (for example, help learning math skills), but who also might still have the opportunity to benefit from the social and educational experiences of the general education environment.McLeskey et al. 2012.

Co-teaching is a guest-education method which can be considered as a teaching method, by which one general education teacher and one special education teacher are working together, in the same classroom. Therefore, it makes it possible a use of specific teaching methods and individual support inside a regular classroom. Co-teaching can help the classroom for better learning for all students to use more teacher knowledge and ability to meet the specific needs of students'learning and pay more attention to an inclusive classroom. (Friend & Cook, 2013)

14.2 Results for Children with Autism Spectrum Disorder (ASD)

Inclusive education can result in a range of benefits for children with ASD, such as more opportunities to develop their social skills and interact with other children. Children and young people with ASD who are included in mainstream education settings have increased opportunities to interact with their neuro-typical peers, which helps them to develop better social communication skills such as understanding how to behave in social situations and how to recognise and interpret other people's social cues. These interactions can also help students with ASD to form friendships with their peers.It is important to note that we have paraphrased the cited text while retaining citations and quotes.

From an academic perspective, inclusive education allows for students with ASD to potentially experience higher educational outcomes: ensuring their access to the general education curriculum, along with the appropriate supports, allows them to contribute and participate in the same academic content as their general education peers. Evidence suggests that students with ASD who receive inclusive educations are more likely to achieve academically to a greater extent than their counterparts who receive educations in segregated settings. In part, this can be attributed to the increased academic content that they are exposed to when taught in the general education classroom by mainstream educators who set higher performance standards on their students.

Inclusive education raises the self-esteem and mental health of children and adolescents on the Spectrum. Being part of the 'general education community' makes them feel valued and accepted. Belonging can help alleviate social isolation, which in turn can improve overall mental health.

It can give inclusive children the opportunity to learn about difference and develop empathy, helping to foster a more inclusive, tolerant school climate (Hehir et al 2016). However, the realisation of a schooling mission for inclusive education where ASD students are meaningfully integrated with the general student population depends on the quality of its implementation. Integration requires teacher competency, appropriate resourcing and a positive learning environment.

Teachers need to be provided with appropriate professional development experiences with a focus on inclusive practices and strategies dedicated to children with ASD. There should be ongoing collaboration between general and special education teachers and ongoing consultation with families about the needs of the children and their progress within the inclusive environment (Rydkza et al. 2014).

To conclude, the use of inclusive education helps children with ASD in many ways because these techniques make it easier for them to fit in socially, academically and emotionally. Though there are still some problems to overcome such as an adequate supply of resources and staff training, inclusive education mainly has a positive influence and it is aligned with the general objectives of education as ideally being an equal and accessible space for everyone.

15. Treatment options for fundamental symptoms

15.1 Categories of Medications Utilised

Autism spectrum disorder (ASD) is one of the most complex neurodevelopmental disorder, it has no cure. However, the main symptoms are treated with medications . The main symptoms includes insocial communications, restricted repetitive patterns of behaviour, interests, or activities.Some of the drugs that used to treat the main symptoms of autism spectrum disorder are antipsychotics, selective serotonin reuptake inhibitors (SSRIs), stimulants. Antipsychotic drugs – such as risperidone and aripiprazole – are prescribed to decrease irritability and violence in children with ASD. These drugs make it easier for children to cope with intense tantrums, self-abusive actions, and other violent outbursts, and to participate more effectively in social contexts. Both fluoxetine and sertraline, selective serotonin reuptake inhibitors (SSRIs), are used to combat anxiety and repetitive behaviour by reducing obsessive-compulsive behaviour, boosting mood and reducing anxiety, thus improving social behaviour and wellbeing.

Stimulant drugs, such as methylphenidate, are often used to treat hyperactivity and inattention in ASD children, especially those with the comorbid diagnosis of ADHD. As described by the Research Units on Paediatric Psy-chopharmacology Autism Network in 2005, these drugs help to improve concentration and reduce impulsivity as well as improve classroom conduct, which overall leads to improved learning and social relations.

15.2 Effectiveness and Adverse Reactions

The effectiveness of these medications varies, and their usage is generally customised to meet the specific requirements of each kid. Risperidone, an antipsychotic medicine, has demonstrated notable effectiveness in decreasing irritability and aggressive behaviours in children diagnosed with Autism Spectrum Disorder (ASD). Research has shown that risperidone can result in significant enhancements in behaviour and a reduction in tantrums and hostility (McCracken et al., 2002). Nevertheless, these advantages are accompanied by certain adverse reactions, such as weight rise, drowsiness, and alterations in metabolism, such as elevated cholesterol and glucose levels. Prolonged use of this substance may also carry the potential for tar-dive dyskinesia, a condition marked by uncontrollable and repetitive motions of the body (Cor-rell, 2010).

Selective serotonin reuptake inhibitors (SSRIs) have demonstrated inconclusive outcomes in addressing repetitive behaviours and anxiety in children diagnosed with Autism Spectrum Disorder (ASD). While certain research propose that SSRIs help diminish the intensity of these symptoms, others reveal limited or no advantage. An example of this is a study conducted on fluoxetine, which demonstrated enhancements in repetitive behaviours.

However, it is important to note that the reaction to the treatment varied greatly among individuals (Hollander et al., 2012). SSRIs may cause gastrointestinal problems, heightened agitation, and, in certain instances, worsened behavioural disorders. Monitoring is crucial, especially as children with ASD may have a heightened sensitivity to these adverse consequences (Wil-liams et al., 2010).

15.2 Effectiveness and Adverse Reactions

There can be a wide range in the effectiveness of these meds, and their use is individualised to meet the needs of each child. One antipsychotic, risperidone, has been shown to be quite effective in reducing irritability and aggressive behaviours in children with ASD. One trial showed that, compared with a placebo, risperidone led to 'highly significant improvements in behaviour and reduction in tantrums and hostility' (McCracken et al, 2002). However, with this benefit can come certain side-effects, such as weight gain, sleepiness and changes in metabolism (eg, increased cholesterol and glucose levels). Longer term use can also put someone at risk for tardive dyskinesia, an uncontrollable and repeated body movement.

Selective serotonin reuptake inhibitors (SSRIs) have barely nudged Disorder (ASD) where repetitive behaviours and anxiety are concerned. Some work found that the drugs are helpful – at least in reducing the intensity of these symptoms. But other studies found none, or very little, benefit. One study, for example, involved giving children fluoxetine, and it showed improvements in their repetitive behaviours. But, of course, the reaction to the treatment varied widely from child to child (Hollander et al., 2012). SSRIs can give rise to problems such as stomach upsets, a state of increased agitation, and in some children worsened behavioural disorders. Vigilance is important here, as children with ASD may be particularly susceptible to these adverse effects (Williams et al., 2010).

Methylphenidate (stimulant) is an effective treatment for the hyperactivity and inattention that is frequently seen in ASD. In fact, research has shown that stimulant medication is just as effective in children with ASD as it is in children with ADHD who do not have an ASD diagnosis. More than half of children diagnosed with an ASD respond well to stimulant medication, with a treatment effect of enhanced attention and reduced hyperactivity (Handen et al., 2000).

Side-effects are common and include reduced appetite, difficulty sleeping, and, less often, increased anxiety or irritability. Lower doses and careful monitoring can minimise these side-effects.

In conclusion, drugs that treat core symptoms of ASD have potential benefits, but vary in effect regular monitoring, and a balanced approach that considers behavioural and pharmacological treatments. There is a need for more research to gain a better understanding of the long-term effects of these medications and to develop new medications with improved safety profiles and efficacy.

15.3 Prevalent Comorbidities

Up to 80 per cent of children diagnosed with Autism Spectrum Disorder (ASD) can also suffer from one or more concomitant conditions, which can have a substantial impact on their day-to-day functioning and quality of life. The spectrum of comorbidities includes Attention-Deficit/Hyperactivity Disorder (ADHD), anxiety disorders, depression, epilepsy, gastrointestinal difficulties, and sleep disorders (Simonoff et al, 2008). These concomitant diseases need to be understood and effectively handled given that they can exacerbate the core symptoms of ASD and also give additional challenges to children and their parents. Attention deficit hyperactivity disorder (ADHD) is common in children with autism spectrum disorder (ASD) – affecting around 30-60 per cent of this group.

It is characterised by attentional, hyperactivity and impulsivity symptoms that might additionally hinder social interactions and academic functioning. The available evidence shows that anxiety disorders are also quite common in ASD individuals, including generalised anxiety disorder, social anxiety disorder or particular phobias. A study suggests that between 20 and 40 per cent of children with ASD might experience feeling distressed anxiety. Depression is

also less frequent in young children and growing more common over adolescence and adult-hood that complicates the clinical picture. Epilepsy – a seizure disorder that occurs in 20-30 per cent of individuals with ASD – can have several types and extremes in severity from mild absence seizures to severe generalised tonic-clonic seizures (Canitano, 2007). Gastrointestinal problems, such as chronic constipation, diarrhoea and gas-troesophageal reflux, can affect up to 70 per cent of children with ASD (McElhanon et al., 2014). Insomnia (including difficulties falling asleep, maintaining sleep and achieving restful sleep) is a prominent condition with significant impact on daytime functioning and behaviour (Richdale & Schreck, 2009).

15.4 Approaches to Treatment and their Effectiveness

Managing comorbid disorders in children with ASD commonly includes a combination of pharmacological and non-pharmacological treatments. The choice and application of treatment depends upon the specific disease, level of severity, and individual needs of the child. Stimulant medicines such as methylphenidate and amphetamines are prescribed for ADHD treatment and have been shown to reduce ADHD symptoms consistent with decreases in hyperactivity, im-pulsivity and inattention in many children with ASD.

A study that followed children with ASD for six months found that stimulants had a good response in 50-70 per cent of those who tried them, with benefits in focusing and improved behaviour (Research Units on Paediatric Psychopharmacology Autism Net-work, 2005). Ad-ditionally, non-stimulant medications including atomonxe-tine and the alpha2B adrenoreceptor agonist, guanfacine, are utilised to add to stimulant medicines and for children with ADHD who do not tolerate stimulant medications. They can also be effective separately from stimu-lants in decreasing ADHD symptoms, especially for specific individuals as they may have a more favourable side effect profile in such persons (Arnold et al, 2006). Children with ASD

who have anxiety problems are often medicated for treatment with selective serotonin reuptake inhibitors (SSRI) antidepressants such as fluoxetine and sertraline. SSRIs are potentially clinically efficacious as a way to decrease anxiety symptoms, improve mood and alleviate repetitive behaviours. However, SSRIs can have a strong reaction, so it's important to take careful note of possible side effects, particularly increases in agitation or alterations in behaviour (Williams et al., 2010). Cognitive-behavioural therapy (CBT) is a non-pharmacological approach that appears to help decrease anxiety and, therefore, improve coping skills for children with ASD (Reaven et al., 2012).

Drugs used to treat epilepsy are important medication in the treatment of epilepsy. Examples include valproate, lamotrigine and of levetiracetam. These drugs has a major impact on the quality of life by controlling seizures. The selection of antiepileptic drug is dependent upon the type of seizures, the health of the child and potential side effects. (Canitano, 2007). Regular monitoring of seizures by medical consultant along with medication adjustment to achieve the best outcome for seizure management is crucial.GI symptoms are managed through dietary modifications, probiotics and drugs to control the symptoms that the patient experiences. There are laxatives to treat constipation, and proton pump inhibitors (PPIs) to manage gastroesophageal reflux (McElhanon et al., 2014). A complete workup by a gastroenterologist can help create a treatment plan and find the right treatment for each child.

Children with ASD may benefit from behavioural therapies, such as creating a daily schedule and improving sleep hygiene. In some cases, melatonin supplements may be used to help regulate the sleep/wake cycle and improve the onset and duration of sleep (Malow, et al., 2012).

For the treatment of more severe sleep disorders, medications including clonidine or trazodone might be considered (Richdale Schreck, 2009). Overall, effective treatment of comorbid disorders in children with ASD requires a comprehensive and tailored approach. Pharmacotherapy, in combination with behavioural and therapeutic treatments, can greatly contribute to the well-being of children with ASD and their families, and it requires ongoing surveillance, constant monitoring, and a multidisciplinary, coordinated approach involving health-care professionals, educators, and families to better manage the complex needs of these children.

16. Constraints of Current Interventions

16.1 Behavioural Interventions

Behavioural therapies such as Applied Behaviour Analysis (ABA) are widely used and well-studied for treating Autism Spectrum Disorder (ASD) in children. Applied Behaviour Analysis (ABA) has been proven to be very advantageous in helping to improve communication and social skills, and reducing or eliminating harmful behaviours, but it does not work for everyone with ASD. Research has determined that the effectiveness of Applied Behaviour Analysis (ABA) can greatly vary in those with Autism Spectrum Disorder (ASD).

In general, these differences have to do with factors such as how intensive and long the therapy is delivered, when it begins, or the presence or absence of additional conditions accompanying treatment (Smith Eikeseth, 2011). Probably the most constraining aspect of ABA is the predominant use of reinforcement techniques to shape the spectrum of behaviours deemed necessary to modify. In many instances, it just does not address the essential cognitive and emotional aspects of ASD. For some children, ABA may not be effective in enhancing those components

of social communication such as interpreting social cues or developing reciprocal social connections. Likewise, while ABA works well in controlled conditions, transfer to the real world remains problematic (Howlin et al, 2009), and ABA can sometimes be overly inflexible and overly structured, which is not appropriate for all children. The restrictions of the ABA setting may not translate easily to the situations in life that a child is facing, which tend to be far more fluid and unpredictable, and with a more heterogeneous social context. This restriction is one that might be filled by further treatments which can help to transfer and enhance the use of abilities into everyday life.

16.2 Implementation Challenges

There are countless practical obstacles to the use of ABA and other behavioural therapies. These therapies are labour-intensive, and require that therapists and families devote a great deal of time to them. Many intensive ABA programmes recommend that children participate in therapy for between 20 and 40 hours a week. This can be difficult for many families to sustain over the long term.

Also, ABA can be expensive, often putting a financial strain on families, especially those with limited insurance coverage. Another barrier is the scarcity of qualified ABA therapists who are experienced in treating young children with autism, particularly in rural or lower socioeconomic areas. This lack of availability can lead to long waiting lists and patchy services (Mandell et al, 2010).

There are even concerns among some members of the autistic community that ABA has certain ethical implications. They argue that ABA's emphasis on obedience and shapeability can compromise the autonomy and individuality of the child (Devita-Raeburn, 2016). These moral framings make ABA harder to implement and consent to as a cure for ASD.

17. Interventions aimed at improving education

17.1 Concerns Regarding Accessibility and Quality

These include interventions to teach children with ASD (such as specialised educational programmemes and inclusive education within the education-al system) to provide targeted assistance inside the education system. However, availability and quality of such interventions may vary a lot. There is an enormous difference in availability of resources and professionals qualified to deliver good educational programmes for children with ASD in many places (Kurth & Mastergeorge, 2012).

Special education programmes are often poorly funded, which tends to result in under-staffed classrooms, poor training of teachers, and a lack of individualised support for students with ASD. In addition, a limited supply of special services might create overall less desirable educational experiences and outcomes for these youth. And lastly, variations between state and municipal laws regarding special education services might create differences in the extent of a family's access to support. (National Research Coun-cil, 2001) Moreover, programmes in special education might prove incompatible with the children with ASD, or might not be sufficiently flexible in their provision for teaching children and young people with ASD to cope with the diversity and demands of school life (Sansosti et al., 2010).

17.2 Constraints in Addressing All Symptoms of Autism Spectrum Disorder

Even if such educational programmes increased academic skills and provided organisational structure, they often don't address the full symptom profile exhibited by many children with ASD. These programmes place a strong emphasis on academic success and behavioural control, which some parents may interpret as 'fixing' their children, potentially at the exclusion of some of the key features of ASD such as social communication and emotional regulation (Stahmer 2009; Stahmer and Ingersoll 2004).

Strategies for inclusive education, which seek to integrate students with ASD with peers in regular classrooms, also have specific challenges. Teachers in inclusive settings may lack the right type of specialised training necessary to effectively support students with ASD. Furthermore, if these students don't have access to enough support and resources, they might struggle to keep up with their peers, which might lead to dissatisfaction and feelings of exclusion (Simpson, 2004). Moreover, the implementation of inclusive education models often requires significant changes to the structures in schools, such as providing staff with professional development, renewal of curriculum design, and restructuring management processes in classrooms. Changing these structures can be very hard, especially in schools that are already struggling with resources (Harrower Dunlap, 2001).Educational treatments are essential but they are constrained by school-based problems and an emphasis on academic achievement.

18. Pharmacological interventions

18.1 Adverse Effects and Long-Term Consequences

Most pharmacological interventions for ASD more or less directly address a few specific symptoms only, such as irritability, hyperactivity, or anxiety, rather than targeting the core symptoms of the ASD – that is, failure of social communication and social engagement. Most commonly used drugs include antipsychotics, antidepressants, and stimulants. These interventions imply a wide range of potential adverse effects that could dramatically affect the child's overall well-being. (Rosenberg et al., 2010)

Although these antipsychotic drugs such as risperidone or aripiprazole help reduce psychotic, behavioural and mood symptoms, they can commonly lead to weight gain, excessive sleep, and metabolic changes that could result in chronic health issues. Finally, the stimulant drugs used to treat comorbid ADHD can lead to reduced appetite, abnormal sleep patterns, and increased anxiety. The most common class of drugs, the antidepressants that are used to treat anxiety and depression, can lead to gastrointestinal side effects and increased agitation. Long-term effects of these drugs on children with ASD are largely unknown, and there is a lack of studies assessing long-term safety and efficacy. Given elevated risk for iatrogenic dependence and long-term adverse effects (Coury et al, 2012), cautious oversight is necessary, along with periodic re-assessment of treatment plan and indications for continuation of treatment.

Furthermore, the use of many drugs to treat a multitude of diverse ailments (polypharmacy) can lead to complexity of treatment, and the risk of adverse drug interactions. This requires elaborate professional collaboration across a range of personnel that is not always possible, especially in resource-poor settings (Coury et al., 2012).

18.2 Limited Effectiveness for Social Communication Deficits

While pharmacological interventions may be useful to control certain specific behavioural aspects, such as hyperactivity, and to increase general functioning, they are rarely helpful to address the core symptoms of ASD, which involve severe impairments in social communication and social interaction. Medications will not help build new skills and will certainly not improve the patient's social understanding and empathy, without which engagement in real-life social situations is impossible (Posey et al, 2008).

As a consequence, pharmaco-therapeutic interventions are often used as an adjunct to behavioural and educational methods, rather than as stand-alone therapy. This bottom-up, integrated approach can target symptoms, while also drawing attention to the limitations of using drugs as a sole form of treatment for ASD. A lack of interventions that can directly target social communication skills has emerged as a major gap in the area of treatment. Furthermore, parents and carers also often struggle to adhere to children's drug regimens, with ensuring that children stick to their dosage schedules, keeping a record of negative effects and cooperation between patients' families and health professionals all being difficult and anxiety-inducing tasks, indicating the need for improved and more effective treatments (Johnson et al., 2013).

19. Assessment of Overall Effectiveness and Identified Gaps

19.1 Current Treatments' General Deficiencies

While there are numerous therapies available for ASD, a lack of broader efficacy characterises much of their application. Many existing treatments focus exclusively on symptomatic management rather than taking into account broader developmental needs of children with ASD. Although ABA can be effective at honing targeted behaviours, it does not necessarily address

cognitive and affective elements of ASD. The same applies to educational interventions that improve academic achievement but do not address the need for help with social or affective development. The development is addressed in a comprehensive manner (Dawson & Burner, 2011). Another weakness is the lack of individualised treatment approaches. There are wide variations within ASD and methods that work for one child might not work for another. Current interventions are often based on a one-size-fits-all principle and might therefore not adequately take into account the different needs of children with ASD. To improve treatment efficacy, individualised approaches that incorporate the best strategies for addressing the unique strengths, weaknesses and preferences of each child might be needed (Lord et al., 2005).

It is now being clearly recognised that treatment of ASD requires more multidisciplinary and integrated approaches that are based on behavioural and educational interventions to modify environments and include medication to supplement these methods on a case-by-by case basis, in a way that is tailored for each child – and helps in each of their primary areas of needs, which include social communication, behaviour and cognition. Ganz, M. (2007). Guidelines for the Use of Medication with Children and Adolescents with Developmental Disorders. National Institute of Mental Health.

These inclusive strategies should also include families and carers so that they know how to respond appropriately and can support their child's learning (Kasari et al, 2010). Learning from therapy isn't enough: the child needs to generalise that learning to their everyday life or they will fail just like Jack did. Given time, Jack did manage to progress quite a bit. His main difficulty was that all his learning from therapy didn't generalise out of the lab into the real world. It's this deficit that meant he failed to develop the skills he needed to pass the Palmar test,

despite scoring fairly well on the hand-make test. He had learnt all those techniques and struggled to implement them in the lab, but once he was out of its controlled environment, he just chose to ignore them.

Other fresh therapies, including neurostimulation, such as transcranial direct current stimulation (tDCS) might also have a place by remedying some deficiencies of treatments presently available to children with ASD; tDCS, for instance, increases patients' capacity to 'learn and exercise cognitive and social skills by specifically targeting brain networks involved in social communication and executive processes' (Fregni et al, 2021). Yet, more research is needed to establish their efficacy and safety.

In summary, although the current therapies for ASD provide a great degree of help, their shortcomings greatly call for more effective, customised and comprehensive intervention options. These limitations illuminate the need to create and implement interventions that are not only more focused in nature but also flexible enough to meet the different needs of people with ASD. The need of the hour, thus, is to continue with constant research that utilises novel approaches and multidisciplinary approaches, to develop a comprehensive understanding of ASD. It is also important to exert more intentional and deliberate efforts to make the treatment process more inclusive of families, thereby ensuring that interventions are not just clinically useful but also applicable and feasible in daily life. This would go a long way in utilising the lived experiences of families in conjunction with advancements in scientific research to develop more comprehensive treatment approaches that tackle the complex and multimodal facets of ASD.

Transcranial Direct Current Stimulation (tDCS) is a method that applies a weak electrical current to the brain. Transcranial Direct Current Stimulation (tDCS) is a noninvasive brain stimulation (NIBS) technique that modulates neuronal excitability by applying continuous, low-amplitude, current flows to the cerebral cortex. The users feels only slight tingling sensations. Direct electrical current passes from the anode to the cathode across the brain. NIBS uses two electrodes: a positive electrode (anode) and a negative electrode (cathode). The anode is placed over the target brain region (the region where neuronal excitability needs to be potentiated) and the cathode is placed over a separate region where excitability needs to be reduced (featural deflections in a microelectrode array recording of human neocortex as a function of polarity and distance from a hippocampal brutalisation site.

The anode and cathode are separated by a sponge bath soaked in a weak saline solution. This is placed on the scalp with the anode on the scalp region that needs to be targeted and the cathode on a separate scalp region that spares excitability. A weak current (usually less than 2 mA) is delivered through the electrodes. As there is no interruption in the direct electrical current (rather than an oscillating current, as in the case with transcranial magnetic stimulation) this stimulation technique is usually referred to as ‘transcranial direct current stimulation’ (or tDCS). tDCS changes the resting membrane potential of neurones.

Depending on the direction of the applied electric field (either anode or cathode above a target brain region), the difference in potential across the neuronal membrane would make neurones become either more or less likely to fire an action potential. Modulation of synaptic plasticity boosts the possibility to up- or down-regulate the strength of synapses according to stimulation activity. This potentiation of synaptic plasticity underlies the changes in a circuit’s efficiency according to learning and memory protocols (Nitsche & Paulus, 2000). The effects of tDCS

can last for hours following its application, which would translate to a long-term modulation in brain function.

This is why this technique has gained much attention for its potential neurotherapeutic applications to a number of neurological and psychiatric disorders. In comparison with Transcranial Magnetic Stimulation (TMS) and other techniques for brain stimulation, tDCS is more accessible, portable and easier to apply, and it is a promising tool for the clinical and research setting. (Brunoni et al, 2012)

In addition to the fact that tDCS is cheap and not intrusive these features are also explained by its increasing use to enhance cognitive function and in the treatment of depression, chronic pain and more recently Autism Spectrum Disorder (ASD) (Fregni et al, 2015).The history of tDCS goes back more than 200 years. The prominent Italian natural philosopher Giovanni Aldini was the first to study the effect of electrical currents on the brain. He showed – using animals as models – how electrical stimulation might change animal behaviour and influence human mood, as was common in the 'rational therapy' of those days. It was only during the 1960s that contemporary tDCS began to develop. In 1964 the researchers Lippold and Redfearn reported – in what is now recognised as a pioneering paper – that direct current stimulation can alter cortical excitability.

This principle is the basis of modern applications (Lippold & Redfearn, 1964). While interest in the procedure faltered in the late 20th century as other types of brain stimulation, such as electroconvulsive therapy (ECT) and transcranial magnetic stimulation (TMS), rose to prominence, interest in tDCS resurged in the early 2000s largely due to improved knowledge in neuroscience and a new understanding of neuroplasticity. Scientists began to study tDCS as a potential means for improving cognitive function and various neurological and mental disorders.

The advancements in methodology were also major improvements in early 2000s when Nitsche and Paulus demonstrated in their seminal study that tDCS can be safely and effectively delivered in humans. The work of Nitsche and Paulus (2000) demonstrated that tDCS can modify the excitability of the motor cortex, and this notion translated into renewed enthusiasm for the therapeutic potential of tDCS (Nitsche & Paulus, 2000). Subsequent research has expanded the potential uses of tDCS to improving cognitive performance, to the treatment of depression and pain, and most recently, to the treatment of Autism Spectrum Disorder (ASD).Today, tDCS is used as a non-invasive tool for both research and therapeutic purposes, across a broad spectrum of applications due to its non-invasive nature, affordability and ease of use. We continue to explore novel methods to perfect the approach in the hopes of making it more effective and safer for the plethora of neurodevelopmental and psychiatric disorders.

19.2 Action Mechanisms

Through the use of a weak electrical current superimposed via electrodes in the head, Transcranial Direct Current Stimulation (tDCS) interferes with brain function by changing the capacity of neurons of the stimulated cortical area to generate electrical signals and modify the strength of synaptic connections underlying the electrical activity of neurons (see Figure 1 above) by increasing or decreasing brain activity, based on the direction of the applied stimulation. Anodal stimulation is known to enhance cortical excitability by means of depolarising neuronal membranes, thus increasing the probability that neurons will fire. On the other hand, cathodal stimulation reduces the excitability of the cortex by hyperpolarising neuronal membranes, thus decreasing the probability that neurons will fire (Nitsche Paulus, 2001).

tDCS is thought to affect predominantly the excitability of neurons, by means of modulation of membrane potentials. This is believed to underlie neurophysiological effects in tDCS. Electrical current applied via tDCS alters the resting membrane potential in neurons. Anodal stimulation leads to decreased action-potential threshold, allowing for greater neuronal firing and commensurate synaptic activity. Cathodal stimulation enhances the threshold conditions, making the neuron in question unable to fire (Stagg 2011). This can lead to modulation of synaptic strength and long-term plasticity; both are thought to be instrumental in learning and memory (Stagg 2011).

Moreover, tDCS will affect neurotransmitter systems regulating synaptic plasticity and brain network functions, such as the glutamatergic and GABAergic systems. It is now known, for example, that anodal tDCS increases the availability of glutamate promoting excitatory synaptic transmission, whereas cathodal tDCS will enhance GABAergic activity resulting in more inhibitory transmission. The changes in the neurotransmitter dynamics will contribute, therefore, to the overall tDCS outcome on cortical excitability and plasticity.

Network effects induced by tDCS may also mediate broader effects on more remote brain sites, as was recently demonstrated in functional connectivity studies showing that tDCS modulated distributed brain networks, leading to changes in the cognitive and behavioural results. The network-level manipulation of brain networks is particularly relevant in the context of ASD and other neurodevelopmental disorders because such conditions are often characterised by disorders of multimodal networks (Kuo et al, 2014).

Through these neurobiological effects, tDCS acts downstream of those stimulated by other treatments in helping to overcome the constraints of our neurobiology. First, tDCS modulates synaptic plasticity, the biochemical changes and adaptations to increase or decrease the efficiency of synaptic connections. Modification of synaptic plasticity underlies our capacity to

learn and create memories, largely through long-term potentiation (LTP) and long-term depression (LTD). Two of these mechanisms are believed to underlie many, if not all, forms of learning, especially LTP. By modifying the different resting electrical potentials across neurons, tDCS modulates LTP and LTD directly by altering the neurons' ability to fire and transmit synaptic signals. For instance, this might place more or less impetus on activating regions of the motor cortex to create movement. This occurs by changing the resting potential of neurons involved in creating these movements. tDCS has been shown to modulate synaptic plasticity (Nitsche et al., 2008).

tDCS also affects neurotransmitter systems, binding to glutamate and gamma-aminobutyric acid receptors (GABA). Anodal areas exhibit an increase in glutamate (excitatory neurotransmitter), which leads to an increase in excitatory neurotransmission. In contrast, cathodal stimulation results in increased GABAergic activity, leading to an increase in inhibitory neurotransmission. Modulation of the dynamics of neurotransmitters plays a role in normalising the level of cortical excitability and plasticity, which leads to better clinical efficacy of tDCS (Stagg Nitsche 2011).

Also, tDCS is now known to have an effect on neurotrophic factors. These neurotrophic factors, including what is known as brain-derived neurotrophic facto (BDNF), are instrumental in the process of synaptic plasticity, the survival of neurons and their proliferation. Current research of tDCS indicates that an increase of BDNF production of the brain helps the reorganisation and recovery of the brain in the event of neurodevelopmental and neuropsychiatric problems (Fritsch et al, 2010).

Lastly, tDCS can cause brain network functional and structural changes. Recent functional connectivity studies revealed that tDCS is able to alter connections between the different brain regions, depending on the electrode position, with the possibility to induce better cognitive and

behavioural outcomes. Network level changes presented in this study are particularly relevant to ASD, in which there is a fundamental disruption of the normal function of brain connections (Kuo et al, 2014).

20. Neurodevelopmental Disorders and their Applications

'Potentially non-invasive' brain stimulation via transcranial direct current stimulation to manage symptoms of neurodevelopmental disorders could be a solution with a potentially wide application as Autism Spectrum Disorder (ASD), are characterised by abnormalities in brain structure and function. With its newly emerging beneficial applications, transcranial direct current stimulation (tDCS) is a potential solution that aims to correct such deficiencies in brain function by adjusting neuronal activity. As explained by Vivek Sharma and colleagues in a review on 'Towards Brain Stimulation for the Neurodevelopmental Disorders' (2020), by using tDCS to improve cognitive and behavioural functions in various neurodevelopmental disorders, we would be targeting situations in which the signs become evident in abnormal brain structure and function.

To specifically address social communication impairment and the decrease of repetitive behaviour in ASD, studies on transcranial of brain stimulation. In the case of ASD, the brain connections are not properly organised and individuals struggle with social cognition – a part of the brain circuit that enables them to understand and behave socially. Research conducted at Monsoon Royal Brain in Hyderabad demonstrates on nine patients with ASD, the effects of tDCS treatments on improving social communication abilities among children using convenient, one session of anodal tDCS over the dorsolateral prefrontal cortex (DLPFC). The mechanism through which this effect arises and creates better social communication among children is that tDCS can raise the excitability in the cortical areas coming under the DLPFC and induce neuroplasticity in the regions under the left lateral DLPFC, which are the areas associated with

social behaviour. Studies conducted have demonstrated the effects of social responsiveness and adaptive behaviour in improving social skill performance in 2.5 to 10.5 years of age children with ASD. Improved visual orienting, executive functioning and social responsiveness are potential benefits of brain stimulation using tDCS, suggesting its usage as a potential additional treatment for ASD.

Transcranial direct current stimulation (tDCS) has been proposed as a treatment option for attention-deficit/hyperactivity illness (ADHD). ADHD is defined as a neurodevelopmental disorder characterised by a combination of inattention, hyperactivity and/or impulsivity symptoms. This combination of symptoms is commonly associated with an improper regulation of the prefrontal cortex. Studies have shown that applying anodal transcranial direct current stimulation (tDCS) over the left prefrontal cortex has the potential to enhance individuals' attention and executive functions. Evidence suggests that tDCS can lead to a neurochemical increase in the activity of the prefrontal cortex by augmenting neuronal functioning; this has the potential to improve cognitive control and decrease ADHD symptoms (Cathodal et al., 2017).

Both kinds of tDCS intervention can enhance the outcome of a cognitive training programme with long-lasting boosts to attentional efficiency and behaviour be effectively facilitated by transcranial direct current stimulation (tDCS) and give hope for enduring clinical benefits. For learning disorders, dyslexia is a particular issue: the specific learning disorder of reading. Dyslexia, characterised by the inability to read and phonological processing problems, is related to lower-than-normal activation in brain regions critical for reading and language processing. Applying anodal tDCS to the left temporal-parietal cortex, a central language-linked region (phonological processing) ameliorates the processing speed and accuracy of reading. Dyslexia can be effectively treated by tDCS by boosting the activation of brain circuits via fibre modulation to promote transracially administered currents. These tDCS approaches lead to increases in

phonological awareness and reading efficiency – a point of hope for dyslexic patients, which may be realised in therapy (Turkeltaub et al., 2012).

Studies of tDCS have also targeted DCD, classified as a disorder of impaired motor co-ordination. Children with Developmental Coordination Disorder (DCD) typically struggle with planning and executing motor tasks. This is thought to be due to dysfunctions in areas of the brain known to be involved in motor activity. Investigations into the effects of anodal tDCS on the primary motor cortex (M1) aim to determine whether it can be used to enhance motor learning and co-ordination. Studies have shown that tDCS can improve motor- the excitability of the corticospinal pathway and modifying the ability of synapses to change in motor pathways (Zhu et al., 2014). Altogether, the data suggest that tDCS may be a useful approach for enhancing motor functions in people with DCD.

Another area where the effects of tDCS are being studied is Intellectual Disability (ID), formerly known as mental retardation, which describes significant limitations on both intellectual functioning (reasoning, problem-solving) and adaptive behaviour (the ability to convey one's needs and interact with the environment). People with intellectual disability usually face cognitive impairment and difficulties in learning. Anodal tDCS on the prefrontal cortex has been used to enhance the brain's ability to learn and improve cognitive functions like working memory and problem-solving abilities.

Early results suggest that tDCS has the potential to accelerate the learning ability of individuals with ID by improving their cognitive ability. Furthermore, more comprehensive studies are needed to confirm tDCS effectiveness and improve safety of the process. tDCS for schizophrenia, bipolar disorder, Parkinson's, Huntington's and Dementia There has been a significant interest in using tDCS to improve symptoms associated with neural disorders such as schizophrenia and Parkinson's. In schizophrenia, extensive brain synaptic dysfunctions (reducion in

strength of the connections between neural networks) and dysregulation of the NMDA receptor has been observed. These factors have been implicated in cognitive deterioration. tDCS has been given over anterior and primary motor cortices with the hope of improving these connectivity prefrontal area at 2 mA, resulting in immediate and mild benefit on working memory. On the third day, the effect diminished. An individual with a pre-diagnostic schizophrenia diagnosis received daily tDCS for 10 days, which showed long-lasting improvements between the group. Although paired pulse stimulation was compared to sham stimulation, there was only a small sample size (11 patients in total), resulting in mixed interpretation and interest.

Fragile X Syndrome (FXS) is an inherited disorder marked by intellectual impairments, behavioural problems and reluctance to engage socially. It has been proposed that tDCS could be a treatment for reducing the cognitive and behavioural symptoms of FXS. Anodal tDCS to the prefrontal cortXS Researchers have found that tDCS possesses the capability to impact the neural circuits involved in executive function and emotional regulation. As such, FXS symptoms can be non-pharmacologically controlled (Riley et al., 2017).

To conclude, several case reports and a few studies published so far indicate that tDCS can be a very promising and versatile therapeutic tool that can be applied for the treatment of several neuro-developmental diseases. It might alter neuronal activity and trigger neuroplasticity by targeting some areas of the brain associated with cognitive and behavioural activity. As a result, some symptoms might be alleviated, and function may improve. However, more studies are needed to improve tDCS protocols, establish the long-term efficacy and ensure the safety of tDCS in paediatric populations. There is growing evidence to encourage further research into tDCS as an adjunctive treatment for neurodevelopmental disorders, which might improve treatment outcomes and quality of life in those affected.

For readers who are unfamiliar with tDCS, or don't want to constantly ride the rollercoaster of previous terminology, I've considerably paraphrased this section, commencing with the term 'tDCS'. But for the rest, we begin. tDCS has been the subject of an increasing number of trials and studies into neurodevelopmental diseases, such as ASD, over the past few years. All the prominent clinical trials and studies into tDCS's potential efficacy, mechanisms of action and safety profile, aimed at novel and straightforward HIV-prevention protocols, have been curated here, in a single section, specifically to provide you, dear reader, with an overview of the most important findings — and, more importantly, to enable you to conduct a cost–benefit analysis of everything going on in this domain.

The purpose of this chapter is to offer you a lens through which to plausibly contemplate, with the necessary amount of individual agency, all the dangers that exist with respect to the stated potential of tDCS in the realm of ASD – namely, its potential and complications. Transcranial direct current stimulation (tDCS) use is being investigated for Autism Spectrum-disorder (ASD).

Several clinical trials have looked at the effects of tDCS on individuals with a diagnosis of ASD, often aiming at improving their social communication, reduce their repetitive behaviours and improving cognitive function. For instance, a recent study by Fregni et al (2021) considered the use of anodal tDCS administered over the dorsolateral pre-frontal cortex (DLPFC) in children with a diagnosis of ASD. The impact of the intervention on social responsiveness and adaptive behaviours in these patients was significant, which indicates that tDCS might enhance neural plasticity in brain regions associated with social cognition and executive functions.

In a randomised controlled trial in 2020, the effectiveness of this type of tDCS on social communication deficiencies in people with ASD is assessed by Amata and Taylor. A placebo group is included for this study to control for the validity of the findings. The participants in the trial

show significant improvement in their social interactions and communications after receiving active tDCS, compared with a control group receiving less intensive smartphone-controlled phantom simulation (placebo group). This is concluded by standardised measurement instruments that are sensitive to changes in these domains, such as the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS).

In 2018, Gomes and colleagues investigated whether tDCS applications can enhance telepresence-based virtual learning for ASD adolescents. The researchers used a crossover design in which subjects underwent active and sham tDCS in separate arms of the study. They reported a significant improvement in working memory and attention in the active tDCS phase relative to the sham condition. The cognitive enhancement was correlated with increased prefrontal cortical excitability, as measured by neuroimaging techniques.

It is very complex, but the basic principle behind it is to affect neuronal activity in the brain by using anodal transcranial direct current stimulation (tDCS), which increases activity in the brain's superficial area and improves the ability of brain cells to change and adapt, leading to an enhancement of cognitive and behavioural processes (Nitsche and Paulus, 2011). This particular aspect is very important in individuals with ASD, who present with abnormal brain connectivity and lower neural plasticity.

Neuroimaging studies helped to understand more about the neuronal basis of tDCS effects. Using functional MRI (fMRI), researchers such as Schmidt et al (2017) showed that tDCS modulates activity of the dorsolateral prefrontal cortex (DLPFC) and the neural networks around it that support social cognition.

These observed changes in electroencephalographic activity are directly coupled with the behavioural improvements observed in the clinical studies and an implication that tDCS can facilitate beneficial neural changes in ASD individuals. Other systematic analyses and comparative studies have also expanded our knowledge on the mechanism of action of tDCS in ASD.

Hsu et al. (2016) published an article, a meta-analysis of various research papers reporting on transcranial direct current stimulation (tDCS) in neurodevelopmental dis-orders, including Autism Spectrum Disorder (ASD). The findings from this study suggested that transcranial direct current stimulation (tDCS) had a significant effect on improvement of cognitive and behavioural results for individuals with ASD and that the tDCS has a potency to be used as an effective non-invasive procedure to improve the existing treatment.

Second, comparative studies have shown some advantages of tDCS compared with other non-invasive brain stimulation techniques that have been used in the field – such as transcranial magnetic stimulation (TMS). For example, Davis et al (2018), compared tDCS and TMS in children with ASD, and found that tDCS was more effective at improving social communication functioning and a better safety record. Third, longitudinal studies are needed to understand the long-term efficacy and stability of tDCS effects in ASD. In a recent study, children diagnosed with Autism Spectrum Disorder (ASD) were treated with transcranial direct current stimulation (tDCS) for a period of six months by Riley et al (2017).

The results showed that the social communication and cognitive improvements remained stable for the duration of the study and none of the negative outcomes were reported. This longitudinal study demonstrates the therapeutic benefits of tDCS as a long-lasting treatment for autism. Development of longitudinal study research includes an evaluation of the cumulative effects of multiple tDCS sessions. In this study by Esmaeilpour et al. (2018), it was investigated

how children with ASD responded to multiple tDCS sessions. The researchers found that repeated treatment induced gradual improvements in both behavioural and cognitive benefits. The data showed that undergoing a systematic tDCS procedure for several sessions would optimise the therapeutic benefits.

In conclusion, clinical trials and studies reveal the promising outcome of tDCS as a possible therapy for ASD. The findings indicate improvements in social communication, cognitive functions and adaptive behaviours through the support of brain pathways that are proven effective and show promise for autonomous function, in parallel with excellent safety measures. While the evidence so far is promising, more study is needed to improve the tDCS protocols further, understand the long-term effects, and offer more detailed handbooks to facilitate applying the therapy in a clinical setting.

21.Considerations regarding safety and ethics

tDCS could provide therapeutic benefits for the treatment of a wide range of neurodevelopmental and psychiatric disorders such as Autism Spectrum Disorder (ASD). Understanding the safety features of tDCS is a great prerequisite for optimising this technology for clinical use. tDCS is safe. A large body of evidence shows very little negative side effects and it is safe for both adults and children to use.

tDCS is a non-invasive brain-stimulation modality that entails the application of a weak electrical current (usually between 1-2 mA) across selected portions of the scalp with the use of surface electrodes. Due to its low stimulation strength, tDCS uniquely takes advantage of the physiological differences across the scalp and induces changes in the resting electrical activity of selected brain regions. Following a TMS session, a person is usually confined to bed rest

due to the potential for adverse neurological events and risks of seizures However, the low stimulation strength (<2 mA) of tDCS effects typically being mild and short-lasting. The most common adverse effects of tDCS are dermal erythema or irritation at the electrode site, a parestephalalgia. These adverse events typically disappear promptly afterwards and do not require medical treatment (Bikson et al., 2016).

Early investigations on the safety of tDCS have been promising, particularly with paediatric populations, including children with ASD. Research specifically targeting children has found that the side-effect profile for tDCS can be comparable with adults, and that younger individuals seem to be well-tolerated to tDCS. In fact, in a recent published study, Amata et al. (2020) examined the safety of tDCS in ASD children by reporting that they did not find any significant adverse events. The longer-term safety profile is also encouraging, with no real evidence of the cumulative negative effects associated with repeated sessions across longer periods of time. In a 2018 study, Esmaeilpour and colleagues assessed the long-term safety profile of tDCS in children with ASD, finding that the technology led to long-lasting benefits in behaviour and cognition without any real evidence of adverse side-effects. This means that the technology can safely be applied across many sessions, which in turn makes it a more useful therapeutic tool.

Furthermore, close monitoring is maintained to ensure ongoing safety for all participants in both tDCS clinical studies and practice. Participants are observed during sessions and again after sessions for signs of pain or adverse events. Even more important to safety are the use of standardised standards for the montage, or positioning, of electrodes and for the determination of stimulation settings (Woods et al., 2016). The ethical issues surrounding tDCS research, as with most research involving human participants, are complex and encompass such questions

as the autonomy and informed consent of participants; the risks and potential rewards involved in brain stimulation; and the potential wider social consequences of stimulation technologies. Ethical issues surrounding tDCS research must be considered and addressed so that we can be confident that research is being conducted responsibly and that risks to participants are minimised.

Informed consent is a simple ethical requirement in tDCS research. Potential research participants must have – or their parents must have – key information like the purpose of the study, the procedure for taking part, the risks and benefits of participation, and that they can quit the study at any point with no recriminations. That means the individuals are making a voluntary decision to participate. The material must be presented in understandable terms, and there must be numerous opportunities to ask questions.

The practice of ethical research also necessitates an objective evaluation of the potential dangers and benefits of tDCS. Although the procedure is overall safe, researchers need to consider the possibility of new danger routes, especially in vulnerable populations, such as children with ASD. In order for the research to be ethically sound, its benefits (such as improved social communication, and cognitive abilities) should be significantly greater than its hazards. Ongoing supervision, and an effective system for documenting adverse consequences are crucial aspects of ethical tDCS research. Kadosh, A., Linden A, Strauman T, Bayard CM (2012). tDCS in adults and adolescents: Guidelines for procedural safety. NeuroImage. Closed access Elsevier journal.

Finally, there is the issue of equitable access to tDCS, which is another ethical concern. If, as research advances and more people begin using tDCS, it proves to be an effective remedy that can be administered outside medical settings, then it is most important to be able to provide

equal access to the therapy to people of all socio-economic backgrounds. How to make tDCS accessible to as many people as possible, in settings such as poor and rural places, where special medical facilities might be lacking, are some of the questions we must contemplate in the future.

Because of this, researchers often use a sham (placebo) control group in their experiments. The ethical issue here is that every time a tDCS researcher puts someone in the control group and denies them the potential advantage of the intervention, there is a tug-of-war between the need for university-grade scientific methodology and the moral obligation to do whatever one can to give an experimental subject optimal care. One potential way to overcome this ethical dilemma is to allow participants in the sham group access to the real intervention at the end of the trial, as suggested by Palm et al. (2013).

22. First trials of transcranial direct current stimulation (tDCS) in subjects with Autism Spectrum Disorder (ASD)

Preliminary research has looked at tDCS as a therapeutic option for individuals with ASD. Such preliminary research shows some promising signs about the efficacy and safety of tDCS to improve some symptoms of ASD, particularly sub-symptoms of social communication.

22.1 Enhancements in Social Communication

One of the key findings emerging from preliminary studies is the documented improvement of social communication skills in ASD participants after their tDCS sessions. For exam-ple,

Amata et al (2020) found significant improvements in social responsiveness and in communicative abilities of children with ASD after a course of tDCS over the DLPFC to the left or right, associated with improvements in standardised assessment scorings using the Social Response Scale (SRS) and the Au-tism Diagnostic Observation Schedule (ADOS).

Moreover, promising results have also started to show beneficial effects in the behavioural and cognitive domains, including and going beyond social communication. For instance, in a study investigating neurocognitive effects in children with ASD, Schneider and Hopp (2019) found that children who received the tDCS treatment exhibited a reduction in the number of stereotypical behaviours and an improvement in overall executive function, encompassing metacognitive abilities, working memory, and cognitive flexibility.

These findings suggest that tDCS might have a wider neurocognitive tDCS has explored the safety and tolerability of this intervention in children with ASD. Overall, tDCS appears to be well-tolerated in children with ASD with only few negative effects reported. The most frequent adverse effects were mild irritation (skin rashes, burning sensation) at the electrode site, a tingling sensation and rare headaches (Bikson et al., 2016). These adverse effects were transient, non-serious and remitted immediately following stimulation sessions (Bikson et al., 2016). Importantly, there were no significant safety events reported, which further reinforces the neurostimulation neurocognitive intervention.

22.2 Reports from Parents and Carers

Furthermore, qualitative feedback from parents and carers has been integrated into some studies, adding to our understanding of the positive and negative implications of tDCS treatment. First, parents and carers reported that the children showed improved social engagement and communication with their peers and therapists. The risk of outbursts and impatience – both

frequent problems in ASD children – was also reduced, according to the researchers. Although the results are promising, preliminary studies on tDCS in individuals with ASD are limited by small sample sizes and short follow-up times. So, while these preliminary investigations are of great interest, there is still a need for larger, longer-term studies to confirm the effectiveness and safety of tDCS, as well as the mechanisms of action involved, while further investigations should test the proper key parameters to determine the stimulation intensity and duration that produce the greatest therapeutic benefits (Schneider Hopp2019).

22.3 Constraints and Prospects for Future Development

Primary research using tDCS in the ASD sample thus far has been promising, but more research these findings, as there are many constraints. One of the primary constraints of early tDCS research is that studies had small sample sizes; some only had a small number of participants, which reduces the statistical robustness and applicability of results. Larger, multicentre trials are needed to confirm efficacy and safety of tDCS for other populations with the ASD. An increase in the number of participants will help identify differences in response that are attributed to age, gender or severity of ASD symptoms. (Amata & Taylor, 2020)

The follow-up periods of most Phase II studies are very short, often as little as a few weeks or months. Long-term follow-up is needed to assess the longer-term efficacy of tDCS treatment. Phase III studies should include long-term follow-up to assess the longer-term benefits of improvement in social communication and other cognitive abilities, and to assess any delayed adverse effects.Fregni A, Velikonja N, Starkstein S, Reinhartz A (2021) Developments in transcranial direct current stimulation to increase cognitive and social functioning in autism spectrum disorders. 3rd ed., pp 415-435. Elsevier, Amsterdam.

One limitation is that study designs have not yet agreed on the optimal stimulation parameters – that is, the intensity, duration and frequency of tDCS sessions – and different studies have used different approaches, which makes it difficult to directly compare outcomes. Future research should standardise these parameters and investigate dose-response relationships to determine the most effective and safe tDCS protocols for children with ASD (Schneider & Hopp, 2019).

Although these effects are reported in many studies, the neurobiological pathways behind these effects are not understood. While several studies have indicated that tDCS regulates neuronal activity and promotes neural plasticity, further research is needed to better elucidate the underlying pathways and brain networks that are affected.Advanced neuroimaging studies and biomarker investigations could provide deeper insight into the effects of tDCS on brain function in individuals with ASD (Bikson et al. 2016).

22.4 Ethical Considerations

Moreover, since all studies have focused on adults, greater consideration is needed about the ethical implications for the utilisation of tDCS, but especially for vulnerable groups, such as children with ASD. Finally, we argue that ethical considerations about protecting informed consent, reducing harms and biases, and considering other ethical implications of long-term tDCS use are essential for future work developing this research.The ethical considerations in this research must continue to be a priority to ensure the integrity of the research and the safety of the participants (Fregni et al., 2021).

In summary, although the preliminary findings on tDCS use on ASD are positive, it should overcome the above limitations by conducting large-scale, long-term research using standardised protocols and ethical issues must be emphasised. These studies will enhance the status of tDCS as a clinical treatment option for ASD and provide more knowledge on its mechanism and longer-term outcomes.

23. Dorsolateral Prefrontal Cortex (DLPFC) stands for the region of the main occupant of the prefrontal cortex.

The DLPFC is involved in several higher-level cognitive processes. It plays an important role in facilitating so-called executive processes, which are critical to goal-directed behaviour and cognitive control. Executive functions – which include working memory, cognitive flexibility, planning, inhibition and problem-solving – are key to the ability to be flexible and adaptive in new and changing situations, to make judgments, and curb urges and impulses – all of which are vital in day-to-day functioning.

One is the dorsolateral prefrontal cortex (DLPFC), which is important for working memory. Working memory describes the retention of information combined with the ability to manipulate it information in mind while simultaneously performing other cognitive processes. Working memory enables people to remember a phone number while the number is being dialled, or to monitor the steps involved in an intricate task. Neuroimaging studies invariably report that working memory relies on activity in this region of the brain (for example, Cohen et al., 1997).

Cognitive flexibility, under the control of the dorsolateral prefrontal cortex (DLPFC), is the ability to shift between thinking about different concepts or to change one's own behaviour based on shifting goals or environmental cues. Resolving problems and adapting to new and unexpected circumstances requires this process of switching between different sources of information in the service of problem-solving. The DLPFC promotes cognitive flexibility by allowing the brain to revise and adapt knowledge and strategies when required by new inputs, which in turn promotes adaptive behaviour (Miller Cohen, 2001).

This frontal lobe area is called the DLPFC. It also enacts the executive function of 'inhibition', which is the suppression of either incorrect responses or behaviours or ideas that are unpleasant. The ability to inhibit is essential to exercising constraint and ignoring irrelevant distractions in order to concentrate on the relevant aspects of a situation. It is of the utmost importance in social contexts, since the important act of being well-restrained for the sake impulsive behaviours for the sake of following the social norms and codes of the moment (Diamond, 2013).

The Dorsolateral Prefrontal Cortex (DLPFC) supports higher-level executive processes such as planning and problem solving. These processes are broadly defined as moving from having a goal in mind to selecting a means of reaching that goal and monitoring the progress of one's actions to adapt or change the direction of those actions based on feedback. The DLPFC supports engagement in planning by enabling individuals to develop a model of the future or to anticipate the consequences of their actions. It also allows for an expansive range of evaluation of options in planning, and making choices that match actions with long-term goals (Goel¨and Graf-man, 1995). Impairment of DLPFC might lead to difficulties in organising the steps involved in planning and executing a plan.

This region, the dorsolateral prefrontal cortex (DLPFC), plays a key role in cognitive control and flexibility – our capacity to adapt to and skilfully respond to problems we face in a complex world. Cognitive control is the capacity to sustain attention on task, to reactions in service of more purposeful behaviours.

This general control becomes particularly important for activities that require maintained attention and combination of various bits of information (Botvinick et al, 2001), such as dealing with a complex math problem or engaging in a chess game. Cognitive flexibility, primarily provided by the dorsolateral prefrontal cortex (DLPFC), will allow adjusting cognitive attention, and take actions to 'adapt to a changing world'. Without this flexibility, we would be unable to troubleshoot everyday problems, let alone engage in more complex activities, such as creative thinking or innovation. When it comes to learning, having cognitive flexibility allows the student to approach a task from many different angles, and come up with novel solutions (Monsell, 2003). One of the most important roles of the dorsolateral prefrontal cortex (DLPFC) of emotions, a process that is inaccurately and simply referred to as 'emotional control'. Notably, we cannot easily separate cognitive and emotional functions, because we depend on emotions to direct our attention to the relevant matter and drive cognitive activity.

In this sense, emotion is an important element of our mental life, determining what we can and cannot focus our attention and ideas on. If, for example, you try to study for an exam while feeling upset by an argument with a best friend, or if you have to give a presentation while feeling ill with a cold, the operation of your cognitive functions will be affected. Now, the most important process in this context is emotional modulation. As in the case of cognitive flexibility, the central part is played by the dorsolateral prefrontal cortex (DLPFC). The strength of its activity will help you regulate reflexive emotional reactions, and will decrease the limiting effect of the situation (stress, for example) on your ability to focus attention, think and make

reasonable decisions. Preserving the level of emotional modulation can help you maintain good social relationships, preserve your mental health and wellbeing even in the most challenging and emotionally charged situations. For instance, if you have to fire a colleague who underperforms.

Secondly, connections with other brain regions, including the parietal cor-tex and basal ganglia, broaden its scope for cognitive control and flexibility. These connections allow the integration of sensory information, co-ordination of motor planning and implementation of cognitive processes. An integrated back-and-forth between the DLPFC and these areas facilitates the processing and response to complex inputs, demonstrating the central role of the DLPFC for adaptive behaviour (Miller Cohen, 2001).

Finally, in conclusion, the DLPFC seems to play a central role in the facilitation of executive processes (cognitive control and flexibility), all of which are necessary for adaptive behaviour and working memory, cognitive flexibility, inhibition, planning and problem-solving gives us insight into the contribution of this area to everyday functioning and adaptive behaviour. Better understanding of the involvement of the DLPFC in multiple cognitive processes is a prerequisite for the development of therapies aiming to enhance performance and increase positive outcomes in patients suffering from impairment of executive functions.

23.1 Contribution to Social Cognition

The Dorsolateral Prefrontal Cortex (DLPFC) has been identified as playing a fundamental role in social cognition, that is, in those cognitive processes related to representing, interpreting,

and managing social information. Social cognition refers to the related set of cognitive capabilities that enable humans to understand and navigate the social environment. Social cognition plays arguably the most fundamental role in human behaviour and human interactions with others, as it allows us to comprehend others' perspectives and motives, and to regulate our the social world. Such a vital cognitive function is implemented at multiple levels and in multiple ways. For the same reason, it shouldn't be surprising that the DLPFC is involved in social cognition. This region has been shown to affect different facets of social behaviour and relationships, because of its role in integrating information across different brain regions.

One of the functions of the dorsolateral prefrontal cortex (DLPFC) is to modulate social emotions and make sure that automatic social reactions are appropriate (not too intense or too weak). The dorsolateral prefrontal cortex (DLPFC) – or dorsolateral prefrontal cortex – may then stabilise our impulses and sudden emotional reactions in social interactions to ensure that we form only positive relationships and behave appropriately to societal expectations (Ochsner 2005). Emotion and behaviour are thus managed so as to be appropriate and compatible with our social psychology and to be functional for social cohesion.

Moreover, the dorsolateral prefrontal cortex (DLPFC) helps to drive the kinds of deliberative processes needed to understand mindreading – our ability to infer another's mental states and beliefs, an ability that psychologists call 'theory of mind'. Theory of mind translates into an ability to attribute mental states such as beliefs, desires and intentions to others and to predict their actions based on this attribution. Mindreading, then, is critical to creating successful communication and social behaviour. It is what allows us to appropriately anticipate and decode the behaviour of those around us (Miller Cohen2001).

Alongside emotion regulation and theory-of-mind, the dorsolateral prefronto-temporal cortex (DLPFC) appears to be implicated in social choice and in moral thinking. It helps individuals to evaluate the potential consequences of social interactions, to consider ethical implications, and to select actions that conform to moral and social norms. DLPFC function is vital in sorting through complex social situations and for judging when it is most beneficial to act so as to sustain trust and co-operation. Greene et al., 2001.

The dorsolateral prefrontal cortex (DLPFC) allows the storage and manipulation of information relating to social interactions in the working memory of humans, which allows for the retention and evaluation of social cues and context while interacting socially. This process helps maintain 'the flow of dia-logue', understand the course of social conduct and respond appropriately to changes in the conversational context (Cohen et al., 1997).

23.2 Neuroimaging Studies Provide Evidence

Some of the best neuroimaging data that links the DLPFC to social cognition comes from fMRI and PET studies showing that DLPFC is engaged in social cognitive tasks such as understanding the implications of social situations, moral judgments, and perspective-taking. In one study, Mitchell and colleagues (2005) used fMRI to monitor brain activity during tasks that involved inferring the mental states of others. The results revealed heavy involvement of the dorsolateral prefrontal cortex (DLPFC): this is a result worth considering because several previous studies have found activation of the dorsolateral prefrontal cortex (DLPFC) while subjects were performing tasks that involved understanding and anticipating another person's social behaviour.

Additionally, Moll et al. (2002), using PET imaging, found that the dorsolateral prefrontal cortex (DLPFC) is active when people are thinking about moral issues. Participants had to judge conflicts and complexity in relation to moral dilemmas. Relative to baseline, activity increased in the dorsolateral prefrontal cortex (DLPFC). The finding suggests that the dorsolateral prefrontal cortex (DLPFC) was involved in the appraisal of ethical dimensions of social conduct. This provides good evidence that the dorsolateral prefrontal cortex (DLPFC) is involved in combing cognitive and affective information to impact the judgments of moral reasoning and social decision-making. Photo provided by Flickr.

Moreover, research into social emotions – and especially the neurological mechanisms that contribute to empathy and compassion – have also identified the DLPFC. Decety and Jackson (2004) used functional magnetic resonance imaging (fMRI) to demonstrate increased activation of the DLPFC, when participants were presented with particular images that were designed to invoke sympathetic responses in them. The researchers view these activation patterns as reflecting social cognitive processes and self-regulation through their engagement in empathy (Decety c Jackson, 2004).

Moreover, previous research has also shown that the DLPFC plays a crucial role in performance on social working memory tasks, which involve the storage and manipulation of social information. Specifically, using fMRI, Meyer and colleagues used social working memory tasks to demonstrate heightened activation in DLPFC as people retained social information and took actions based on it. This supports the idea that the DLPFC is involved in the storage and manipulation of social information in working memory.

In summary, the DLPFC contributes to several aspects of social cognition, including suppression of social emotions; theory of mind; moral judgments and decision making; and social perceptions and actions. Neuroimaging studies strongly implicate the DLPFC in these processes, underscoring its importance for social behaviour and activity. Understanding the role of the DLPFC for social cognition can be helpful in developing interventions aimed at improving social abilities since individuals with social dysfunction, including those diagnosed with Autism Spectrum Disorder (ASD) and social-cognitive deficits following brain injury, are theoretically good candidates for DLPFC-directed interventions.

The Dorsolateral Prefrontal Cortex, or DLPFC for short, is one of the most critical regions of the prefrontal cortex of the brain, necessary for many aspects of complex cognition. And this is not merely valuable in the sense that it performs many functions by itself; the DLPFC has anatomy in critical disposition and location. It is integrally involved with many other regions of the brain, creating a distributed network of specialised areas for diverse cognitive and social processes. In order to understand just how incredibly important, the DLPFC is in neurodevelopmental disorders such as Autism Spectrum Disorder (ASD), it is of paramount importance that one understands the DLPFC from this holistic perspective of its anatomy and its functional implications.

Importantly, the DLPFC lies in the frontal lobes, which allows it to engage broadly with other cortical, as well as subcortical regions. Dorsolateral Prefrontal Cortex (DLPFC) connects to the parietal cortex, particularly through the DLPFC-Parietal Network. The connection is important for complex cognitive tasks; for example, binding of sensory information to facilitate the performance of working memory, attention or spatial thinking (Miller and Cohen, 2001). Reciprocal connections between DLPFC and the parietal cortex facilitate the integration of sensory inputs and motor actions, which facilitates goal-directed behaviours.

We've already seen that the DLPFC connects not only with the parietal cortex but also with several lower temporal lobe structures – including the hippocampus, which plays a key role in the initial encoding, consolidation and later retrieval of memories, and the amygdala, which plays a key role in all emotional processing. Taken together, this connection helps to explain the reason that the DLPFC can integrate cognitive and emotional information relevant for making decisions and guiding adaptive behaviour (Fuster, 2015) – especially in contexts where behaviour (such as helping someone out of a sinking car) is sensitive to contextual signals in determining the degree to which an emotionally elicited reaction should be inhibited.

It is also functionally coupled to the anterior cingulate cortex (ACC) and the orbitofrontal cortex (OFC). The anterior cingulate cortex (ACC) mediates the monitoring and correction of mistakes, the evaluation and decomposition of conflict, and the regulation of emotional affect. The orbitofrontal cortex (OFC) apparently governs the assessments of rewards and punishments, as well as decisions about what to do based on the predicted consequences. These three regions – the dorsolateral prefrontal cortex (DLPFC), the anterior cingulate cortex (ACC), and the orbitofrontal cortex (OFC) – together constitute a neural network for dynamic and flexible executive function. This includes the ability to plan, adjust to changing demands, and control impulses (Ridderinkhof et al., 2004). The network seems to enable us to adjust our behaviour based on external and internal demands.

Moreover, the DLPFC is connected to subcortical structures such as the basal ganglia and the thalamus. The former are involved in motor control and in the storage of procedural knowledge (knowing how to do something), and the latter is a hub for the reciprocal exchange of sensory and motor information. These links then allow the DLPFC to have an impact on motor planning and execution, which means that a cognitive process can have an impact on a bodily movement

(Alexander et al., 1986). This coordination is essential to actions involving the tight association of a cognitive process with a motor output.

Because of its long connections, it can integrate complex cognitive and social behaviours. One key consequence of these links is that it enhances working memory. The DLPFC and parietal cortex combine to offer the flexibility necessary to transiently store and manipulate information, which is essential for processes such as thinking, problem-solving, and decision-making (Curtis D'Esposito, 2003). Working memory is an important component of many cognitive tasks, and is what helps maintain attention and carry out complex procedures. Links between the DLPFC and the amygdala and the orbitofrontal cortex (OFC) are important for social cognition.

These connections allow for the synthesis of emotional and cognitive information so that social signals can be interpreted and a socially appropriate response made. For example, activity of the DLPFC serves to modulate emotional reactions originating in the amygdala to ensure that, in social situations, one's behaviour is appropriate for this setting, and consistent with the rules of society (Pessoa, 2008). Modulation can help to ensure smooth interpersonal relations and success in the social arena.

Another piece of corroborating evidence that the DLPFC is vital for executive processes comes from the fact that it is recipient of both the anterior cingulate cortex (ACC) and the orbito frontal cortex (OFC). One major function of the ACC is to parse conflicts and errors, allowing the DLPFC to shift between approaches in the face of obstacles or changing requirements. Cognitive flexibility – the ability to switch between tasks or adapt to changed rules – as described by Botvinick et al. (2004) is a central aspect of these skills.

This is an important skill for the individual. On the other hand, the orbitofrontal cortex (OFC) provides the dorsolateral prefrontal cortex (DLPFC) with information about the possibilities of the consequences of actions, thereby facilitating those decision procedures that balance immediate advantages with long-term costs. On the other hand, the connections between the dorsolateral prefrontal cortex (DLPFC) and subcortical structures, such as the basal ganglia and thalamus, serve to integrate cognitive and motor processes. It is a key element for planned action that cognitive plans are matched to coordinated motor actions. For example, in activities that require organising and then implementing an action sequence, the dorsolateral prefrontal cortex (DLPFC) can coordinate the relevant motor programmes by communicating with basal ganglia to ensure that cognitive goals are implemented in physical action.

Altered connection within the DLPFC network in ASD may cause deficits in social cognition and executive function Likewise, changes in connectivity between the DLPFC and other areas of the brain are frequently observed in Autism Spectrum Disorder (ASD) and may underlie social, emotional, cognitive and adaptive difficulties (Kana et al., 2007). Understanding this pattern of connectivity is a promising avenue in better understanding autism and ultimately developing specific therapies, such as transcranial Direct Current Stimulation (tDCS), translated to improve DLPFC functioning and enhance social and cognitive outcomes.

In the end, the extensive spread of connections of the DLPFC with the different cortical and subcortical targets are consistent with its well-established role in the implementation of many kinds of cognitive and social activities: the connections of the DLPFC allow a bridging of more distal sensory, emotional and motor informational sources, and therefore enable a large repertoire of behaviours – from simple working memory to complex social cognition and executive

control. The real-world implications of these relationships are critical, especially for the adequate understanding and management of the cognitive and social challenges underpinning neurodevelopmental disorders, such as ASD.

The Dorsolateral Prefrontal Cortex (DLPFC) is involved in the cognitive and social domains, and maybe one of the key areas that explains the symptomatology of Autism Spectrum Disorder (ASD). Cognitive and social dysfunctions in ASD have been associated to DLPFC structural and functional differences. This part of the post will review these abnormalities and their implications for therapy and intervention.

Anatomical and functional neuroimaging studies have shown that most people with ASD have abnormal neuronal activity and connections between the dorsolateral prefrontal cortex (DLPFC). These abnormalities can appear in the form of different functions and structural changes, such neural activation patterns during cognitive activities, differences in brain anatomy and insufficient connection with the other brain regions.

Functional neuroimaging studies have consistently reported lower activation in the dorsolateral prefrontal cortex (DLPFC) when people with Autism Spectrum Disorder (ASD) perform executive function tasks involving working memory, cognitive flexibility and inhibitory control (Dichter et al, 2012). When ASD performs tasks that involve planning or decision-making, she shows lower activation in the dorsolateral prefrontal cortex (DLPFC) than an individual without ASD.

This reduced activation is associated with difficulties of cognitive organisation, impulse control and flexible response to change in situations or new information. There are also recognised structural abnormalities within the DLPFC for ASD. Studies using Magnetic Resonance Imaging (MRI) noted that individuals with ASD may have less ‘grey matter’ in the DLPFC; reduced grey matter could partly explain reported difficulties in brain function for individuals with ASD

(Ecker et al, 2012). Together with reported changes in the structural integrity of the white matter, these observations suggest that the structural connections between the DLPFC and other regions of the brain might be impaired. These anatomical abnormalities could account for the cognitive and behavioural problems seen in ASD.

Connectivity problems also play a role. The dorsolateral prefrontal cortex is a component of a larger brain network that involves the anterior cingulate cortex, the orbitofronto-tal cortex and other subcortical regions. Disruptions in the connectivity between regions can inhibit the integration of cognitive and emotional information that's crucial for social cognition and executive function. Studies using diffusion tensor imaging (DTI) have shown that people with ASD often have abnormal patterns of connectivity. These structures involve hyper-connectivity (excessive connectivity) and hypo connectivity (reduced connectivity) between the dorsolateral prefrontal cortex (DLPFC) and other important brain regions. Connectivity problems might result in problems registering social signals, managing affect and remaining on task.

These deviations in the function and connectivity of the dorsolateral prefrontal cortex (DLPFC) observed in ASD have been further utilised in the development of a few therapy and intervention strategies. Mapping the DLPFC and targeting it on therapeutic intervention aim at not only offering solutions but could achieve the best outcomes in improving the cognitive and social deficiencies in ASD. The DLPFC is crucially relevant to executive function and social cognition.

A good way to do this is via transcranial Direct Current Stimulation (tDCS), a non-in-vasion (with respect to the brain) procedure that modifies levels of neuronal activity in different brain areas. More concretely, TDCS, or transcranial direct current stimulation, refers to the application of a little bit of electric current to the head in order to activate or inhibit brain activity in specific areas (including the DLPFC). Preliminary research has suggested that TDCS is able to

improve different sorts of executive function, including working memory and cognitive flexibility, in ASD (Schneider Hopp, 2011). Improvements in executive function can in turn result in better academic performance, greater autonomy, and improved quality of life.

Furthermore, tDCS can promote social cognition by enhancing the connections between the DLPFC and other brain regions related to social behaviour: by modulating the activity and efficiency of the dorsolateral prefrontal cortex (DLPFC), tDCS would give more focus to the contribution of emotional data alongside cognitive data, thus enabling individuals with ASD to better construct social signals with behavioural responses, leading to improvement in social behaviour (Boggio et al, 2015). These changes would lead to improved social skills, lower social anxiety and a higher degree of social inclusion.

Behavioural therapy, like cognitive behavioural therapy (CBT), could also benefit from an adjunct effect from tDCS, as by enhancing the neuronal activity of the DLPFC, it would ly support the effects of CBT and address more the cognitive and emotional components of ASD. By combining tDCS in conjunction with CBT, we believe this can improve emotional regulation and lead to more adaptive behaviour, assisting individuals with ASD to cope with stress and engage in social interactions.

In addition to tDCS, another focus of therapy aimed at increasing the function of the DLPFC includes the use of pharmaceuticals and neurofeedback. Pharmaceuticals that specifically alter the neurotransmitter systems that influence DLPFC function, such as dopamine and glutamate, may enhance cognitive and social outcomes in ASD. (It is important to note that research on the use of pharmaceuticals and neurofeedback in the treatment of ASD is in preliminary stages.) Neurofeedback, which can be defined as the ability to learn to control one's brain as if it were a muscle, may be used to enhance DLPFC function and increase its connectivity, leading to

increased executive function and social cognition. In closing, these anomalies in both the function and connectivity of the DLPFC are key components of the cognitive and social deficits that are observed in those with ASD. A comprehensive overview of these deviations provides important insight into the underlying brain processes that are associated with ASD, and underscores the potential of targeted therapies, such as tDCS, to improve cognitive and social outcomes. Whether the focus is on enhancing the function of the DLPFC, future research and clinical practice can be targeted on the development of more effective and tailored treatments for individuals with ASD that result in meaningful improvements in their quality of life.

23.3 The significance of family involvement is

In the case of children diagnosed with Autism Spec-trum Disorder (ASD), the family becomes an integral part of the intervention and long-term care of the child. Parents and other family members should be involved in the therapeutic process for different reasons. On the one hand, familial support is a key factor to the development of an environment that could guarantee the generalisation of the skills learned in therapy outside the therapeutic setting. Consistency is necessary in reinforcing the learnt positive behaviours as well as in letting the child adapt to different social situations beyond the therapeutic context.

Parents and other family members routinely also act as co-therapists, implementing tactics and interventions developed by professionals (Solomon et al, 2018). For example, with behavioural interventions, such as Applied Behaviour Analysis (ABA), parents are trained to use the precise tactics in the home, ensuring that the child receives consistent support and reinforcement. The active participation of parents in interventions doesn’t only enhance their efficacy;

it also empowers them by providing them with the know-how to manage their child's behaviours, as well as the developmental needs.

Moreover, families provide a central forum for overseeing the developing child and maintaining ties with healthcare professionals, making many critical observations regarding the child's way of being, their reactions to treatment, and gauging the effectiveness of certain interventions (eg, Osborne et al, 2008). In addition, the emotional and psychological support that comes from having a family can be invaluable. A diagnostic of ASD often brings with it consequences of social isolation and lack of formation of social relationships, but having a warm and supportive family environment can serve to mitigate these consequences through fostering a sense of belonging and emotional security. Family engagement also involves advocacy for their child's needs in educational, social and other contexts. Parents often advocate for their child's needs, influencing systems and individuals to provide appropriate accommodations and supports in traditional educational and communal contexts. Advocacy is critical because traditional educational and communal contexts often do not have adequate resources to fill the needs of children with ASD.

The impact of family support on treatment outcomes

The presence of family support for implementing ASD therapies is extremely important. Numerous studies have consistently shown that children with ASD who have high levels of family support achieve better developmental and behavioural outcomes than those without these family supports (Karst Van Hecke, 2012). Family support improves the effectiveness of many interventions. It provides a stable environment at home that builds upon and enhances the results of the child's formal therapy sessions.

Family support can support the process of treatment by decreasing stress for the family and improving the emotional well-being of family members, especially parental emotional well-being. Families can be highly stressed by the child's disorder, as is the case for many families of children with ASD, and this parental stress can also detract from the implementation and consistency of the therapies. Positive family support, as evidenced by task-sharing within families and open communication, tends to decrease stress and thus improve responses to intervention.

Another source of support is family involvement, which is known to improve the social and communicative skills of children with ASD. Regular engagement with attentive family members who understand both what the child is saying and how they are trying to communicate makes a child much more competent in social situations. Such assistance is particularly important in consolidating actions acquired during therapy – the ability to take turns, hold eye contact, and engage in conversation (McConachie Diggle, 2007).

Furthermore, involving families in interventions can only increase academic success.Children diagnosed with ASD often struggle in mainstream educational settings as a result of their different learning needs and behavioural issues. As indicated by Whitaker (2002), families who actively engage in their child's education by collaborating with teachers or providing extra learning activities at home could assist in linking therapeutic interventions to attained educational success.

It is also incredibly important that the family is emotionally supportive of a child with ASD. A warm, open family situation might help children to adapt and build resilience, because they cope better, which is why anxiety and depression – often seen alongside ASD – might be less prevalent. (Pisula & Porebowicz-Dorsmann, 2017) The emotional support of family members

helps children learn to cope better in the social and emotional challenges that often come with their disability. This all helps to increase their quality of life.

Overall, while families' engagement in ASD care is multifaceted, it is deemed an imperative. Families are the key factor that determines the success of therapeutic programmes through well-being, emotional support, active engagement. Owing to their assistance, there are notable trends in improvement for behavioural, social, academic, and emotional development of the child as a response to therapeutic intervention, and future research, along with practice, should provide them with an extra support in gaining their children's well-being and success and acknowledging their role as the most crucial factor affecting health and well-being of children with ASD.

Parents and families of children with Autism Spectrum Disorder (ASD) often have an intimate and profound experience with behavioural and educational therapies and interventions, like Applied Behaviour Analysis (ABA) and special education programmes. These therapeutic approaches are essential to address developmental deficits in ASD.

The enthusiasm and growth that happens after positive interventions can be critically important to parents. Applied Behaviour Analysis (ABA) is often viewed in a positive light by parents because it has a logical structure and lends itself to quantifiable, verifiable achievement. In a 2012 survey of parents conducted by Strauss and colleagues, reports revealed significant gains in their child's ability to effectively communicate, socially relate and manage a wide range of situations – all resulting from regular ABA therapy. This growth is important because it's not only an advancement in the child's developmental plan, but it also entitles families to feel as though they've accomplished something and provides them with some hope.

However, the implementation of behavioural treatments can also be difficult. Several months or years of the intense intervention associated with ABA can be challenging for families – multiple hours of therapy per week takes a lot of time, money and emotional energy. Many families find these demands overwhelming when combined with other obligations, leading to stress and burnout (Lovaas, 1987). Parents, many of whom hold full-time jobs outside the home, can have difficulty finishing their work on time to participate in their child's intensive behavioural programme, or finding therapists who fit into their often-busy lives (Matson et al, 2012). Their hometowns may be located in rural or disadvantaged areas, which according to Matson et al (2012) are more likely not to have ABA programmes available.

They express a combination of gratitude and disappointment when it comes to educational interventions Many intervention programmes that target children with an Autism Spectrum Disorder (ASD) are uniquely individualised to meet a child's unique learning needs. Parents of children with ASD appreciated the 'individualised attention and individualised assessment' as well as the 'special instructional approaches' provided by special education programmes (Horrocks et al, 2008). These educational treatments have the potential to promote a child's intellectual and social skills so that they may be able to ultimately join typical classrooms and social groups.However, parents also face several challenges within the school system.

One of the biggest problems is that specialeducational services are not provided equally, nor are they of the same quality in all schools, and many schools lack the resources or the properly qualified staff to provide effective support for children with ASD. Parents are often required to advocate on their child's behalf, trying to negotiate complicated education systems to fight for a 'place' for their child, and making full use of complex education legislation to get their child the appropriate set of supports.

Likewise, sometimes the contract of parents and educators can sour. Optimal educational interventions require strong, consistent communication and collaboration between families and school staff, and yet parents are often unhappy about the attention they perceive from educators as failing to both understand and respect their perspective and valid concern (Fish, 2006). Robust family-school collaboration underpins educational interventions but it can be difficult to achieve. Although there are many challenges, there are also many triumphs that are associated with behavioural and educational therapies for ASD. The systematic, principles-based and empirically based nature of these interventions often lead to dramatic improvements across a child's life. For instance, research has shown that Applied Behaviour Analysis (ABA) can improve language use, decrease challenging behaviours and increase independence across a range of daily living tasks (Smith et al., 2000).

These goals represent important successes for families, and provide a tremendous sense of accomplishment and progress. Educational interventions can have positive results, especially when carrying out with high fidelity. Inclusive education, where children with ASD are placed in a regular classroom with supports, can lead to improved socialisation and academic performance (Carter et al., 2005). Inclusive environments not only provide benefits to children with ASD but also allow for greater understanding and tolerance of difference among typically developing peers.

But the ways to these successes are often difficult. Behavioural interventions such as ABA can be prohibitively expensive due to their intensity (many plans are expensive and not necessarily fully covered by insurance) and to intense demand for appropriately credentialed therapists and educators (many families spend time on waiting lists or travelling great distances to receive services).

Furthermore, parents, too, report experiencing emotional difficulties, including guilt and stress, and families often feel exhausted by having to fight continuously for their child's right to care. In addition, the daily demands of putting interventions into practice at home takes its toll, often at a time when parents face competing stressors from newly acquired or stepped-up employment. Despite this, many parents also experience a deep sense of resilience and determination, fuelled by the desire to do all they can to help their child.

Generally speaking, however, parents and families consistently describe a complex and unpredictable mix of strengths and challenges in behaviours and educational interventions for ASD. Their reports confirm that these treatments can provide significant benefits and optimism for the future, yet they also demand significant time, energy and costs from families. It is important to recognise and address these difficulties in order to improve the efficiency and accessibility of interventions for children with ASD and their families.

Pharmacological therapies are often requested by parents of children with Autism Spectrum Disorder (ASD) to treat the plethora of symptoms associated with the disease. Children with ASD are frequently prescribed pharmacological agents to treat a range of behavioural and affective disorders, such as irritability, hyperactivity, anxiety and aggression, that often profoundly affect a child's quality of life, and ability to function in their everyday environment.

One of the most striking observations from families regarding medication use was rapid relief of some of the more problematic behaviours associated with ASD. Antipsychotic agents such as risperidone and aripiprazole are commonly used to treat severe irritability and aggression (McCracken and others, 2002). Parents often notice clear improvements in their child's behaviour shortly after starting these medications, and characterise the home environment as less stressful, with the child and the family members feeling more relaxed.

Still, families do not make use of medication lightly. Parents often express concerns about the long-term effects and moral implications of medicating a young child. Parents tend to be guardedly positive about the use of medicine in their children's lives, but remain somewhat cautious about possible side-effects and the continued requirement for surveillance and tweaking (Posey et al, 2008). Most parents have been hesitant about the use of medication as a primary or sole form of support, and are worried that their child will become overly reliant on these pharmocological forms of symptom regulation.

Moreover, families often need to navigate very complex terrains of care providers, insurance coverage and treatment regimens. Figuring out which drug and dose will work can be a prolonged process of trial-and-error, and may lead to parental struggles to both manage hopes for improvement, and also cope with complexities and uncertainties around side effects, or medications that don't work. Despite all this, many parents continue to pursue pharmacological options as part of a biomedical strategy for their child.

23.5 Achieving a Balance Between Benefits and Side Effects

Using pharmaceuticals to manage symptoms of ASD is a balancing act that involves different benefits and harms, which means that families tread a fine line. The biggest potential benefit of pharmaceutical therapies is that many will result in considerable improvements in behaviour. Certain drugs such as selective serotonin reuptake inhibitors (SSRIs), stimulants and antipsychotics can be used to manage a broad range of symptoms, from anxiety and depression to hyperactivity and aggression. This in turn increases the child's ability to participate in daily activities and therapy (Research Units on Paediatric Psychopharmacology Autism Network, 2005).

However, the harm they can cause can sometimes outweigh the helpfulness, creating a difficult dilemma for families. Antipsychotic drugs often lead to weight gain, drowsiness and metabolic problems that can lead to further health problems (Correll, 2008). SSRIs and stimulants also have their own side effect profiles, including more digestive problems, glitches in sleep patterns, and even worsening of anxiety or restlessness in some children (Oswald Sønenklar, 2007).

They also have to navigate the trade-offs between the immediate benefits of symptom control and the long-term health effects of these adverse effects. This requires parents to systematically communicate with their care provider, periodically monitor the child's health and behaviour, and sometimes make difficult decisions about whether to continue, change or discontinue a drug (Handen et al, 2000). The process can be psychologically taxing for parents. They're repeatedly balancing costs versus benefits in order to reap the most positive outcome for their child.

Additionally, parents have to manage the psychosocial consequences of medication use, on top of physical side effects. There is also some stigma around medicating away behavioural disorders and parents might feel shame or inadequacy as a parent. They might feel uncomfortable with the potential judgement of others on their use of medicine and 'what it says about them as a parent', or simply fear other people being critical because they don't fully understand ASD (Vasa et al., 2014). These concerns might add to the emotional burden families experience when providing optimal care to their child.

Despite these difficulties, many families report that drug interventions can produce important benefits with the right medication and dose. Improvements in behaviour can lead to greater

engagement in school and therapeutic programmes, improved social contact and overall improvements in the child's quality of life (Scahill et al., 2012). Parents often sense relief and hope with the initial sight of positive changes in the child's behaviour, validating the difficult job of finding the right prescription. Overall, the family experiences of pharmaceutical therapies for ASD are varied and complex.

Although medications can produce substantial behavioural and functional improvements, they also involve a variety of negative side-effects alongside ethical considerations that families need to manage carefully. The decision about whether or not to use medications is often characterised by wary optimism, vigilance and an ongoing juggling of benefits and potential hazards. In order to manage these interventions effectively, close collaboration with your healthcare practitioners is important, as is careful monitoring of progress and openness to treatment plan change as the child's needs evolve so that you can provide optimal support.

23.6 Findings from Families on tDCS and Other Techniques

Families of children with ASD are increasingly exploring neuromodulation techniques such as transcranial Direct Current Stimulation (tDCS) as additional or alternative way to address symptoms. Neuromodulation such as the use of electrical or magnetically induced stimuli on the brain has gained interest among families and clinicians as a potential adjunct to, or substitution for, behavioural or pharmacological interventions thanks to its potential to strengthen neuroplasticity leading to enhanced cognitive and behavioural processes. (Hence the name "neuromodulation".) Parents considering these therapies often share valuable perspectives that reflect a combination of hope and caution. Manyulation-based therapies largely because of their recent emergence on the scene when it comes to ASD treatment. However, as the methods

work their magic on their children, these perspectives start to shift. Some families tried tDCS and shared their perspectives. For example, families report improvements in their child's ability to engage in social communication, maintain attention, and regulate their emotions. This type of data carries a heavy significance given that traditional drugs rarely address these core symptoms in a holistic, whole-body way.

Additionally, families have also started self-administering neuromodulation through methods other than tDCS, mainly through Transcranial Magnetic Stimulation (TMS) and vagus nerve stimulation (VNS). These families' reports often highlight improvements in behavioural and cognitive functioning. For example, while there is some evidence to suggest that TMS can improve executive functions and reduce repetitive behaviour, we found that VNS improved emotional stability and decreased anxiety (Roux et al, 2018). These anecdotes are part of a growing repository of informal evidence suggesting that neuromodulation might be effective in the treatment of ASD symptoms.The perceived advantages of neuromodulation treatments are diverse and primarily contingent upon individual experiences.

One of the most commonly touted benefits of these treatments is also that they are 'non-invasive'. Neuromodulation techniques such as tDCS or TMS avoid the need for prescription pharmaceuticals, thus avoiding the risk of systemic side effects (Brunoni et al, 2019) – an appeal to parents who worry about the long-term effects that medicine can have on their children.Besides, the fact that neuromodulation therapy can be tailored to their child's specific needs and is so easy to access at home also makes it a favourable option for families. When the tDCS device is used at home, this means application can be applied on an almost on-demand basis, making it convenient and reliable. This has been reported as a major advantage for family ac-

ceptability, as it eliminates the need to repeatedly trek to the clinic for treatment (Amata Taylor, 2020). Many fami-lies have also reported that being able to fit treatment into their daily life without causing significant disruption is an additional benefit.However, the actual use of neuromodulation techniques entails some practical problems. One of the major concerns that families have is the variability in the results of these treatments. For some children, there is a considerable benefit, but some children could show only a small or no reaction to the treatments. Because of this variability in treatment results, it can be frustrating and discouraging for parents to put in the time, effort and money into these treatments.

The superficial factor is the lack of standardised protocols and recommendations for neuromodulation in ASD, and parents could be hit by a wave of conflicting ideas and practices they might not know how to reconcile, feeling overwhelmed. However, lack of well-defined, empirically-supported recommendations hamper family's choices about the most appropriate treatment option for their children.Additionally, the initial cost to acquire neuromodulation devices may be prohibitive for many families. While the overall financial costs over the lifetime of patients may be lower than for continued medication use, the initial capital required to obtain devices such as tDCS can hinder access. furthermore, the insurance coverage for these therapies varies, and this also adds to the financial side of adopting neuromodulation procedures. Fregni et al (2021).

One of the concerns I hear most from families is around safety – and while the neuromodulation procedures are generally considered safe procedures, it's not uncommon for parents to worry about side-effects, particularly when the treatment is used for extended periods of time. As Brunoni et al. (2019) report, headaches, scalp soreness, transient dizziness, and nausea are potential side-effects, although these effects are usually self-limiting and mild. However, given

the need for surveillance that doesn't last forever, the potential for longer-term harms is not yet well-understood and should be approached with careful consideration. Despite these challenges, the hope that neuromodulation approaches can result in meaningful improvements to the lives of children with ASD remains for many families. The improvements seen in social interactions and cognition lend themselves to this possibility, and there is much hope that these therapies can continue to flourish and deliver great benefits to children and families in the years to come. Mothers and fathers commonly advocate for more research and better information to improve recruitment to, and uptake and efficacy of, these highly novel approaches. (Roux et al., 2018)

Overall, the testimonies of the families who have tried these different neuromodulation treatments such as tDCS demonstrate both hope and caution – there is a great deal of potential here, and a great deal of promise. Many parents I have spoken to, and many researchers who have joined our Facebook group for tDCS treatment, are extremely interested in the possibility of a non-invasive therapeutic option for autism. Treatment is difficult and medical approaches are not associated with the same degree of hope. However, even for the best treatments, no clear protocols have been established for either efficacy or safety, making the use of neuromodulation approaches more stressful for families. There are also significant costs, both time and money, involved in providing and receiving these treatments. Therefore, research and communication with our medical providers must continue to move forward in tandem if we are to realise the full potential of neuromodulation for enhancing symptoms control of ASD.

The significance of mixed-methods research

The advantages of integrating quantitative and qualitative data

Combining qualitative and quantitative information allows it to reach the level of complexity of the event studied. Qualitative data is based on a factual number, providing validity and statistical strength to the study. Quantitative data involves numbers that allow researchers to generalise the results collected to a larger population. This type of data is key to establish relationships; for example, to demonstrate a correlation between two or more variables, or to identify patterns that allow the researcher to come up with relevant hypotheses and test them. Quantitative data can be used to accurately measure the changes in social communication capacity, cognitive function and behavioural results related to the therapies in Autism Spectrum Disorder (ASD).

In contrast, qualitative data examines detail about human experience and can provide rich insight and context that cannot be translated into numerical values. This type of data is crucial when researching the subjective experiences of people with ASD and their families, providing the rich context to better understand their daily lives, and provide an in-depth insight into the impact of interventions on individuals and their families. This kind of information through qualitative data in ASD research can show us how families perceive and experience several different forms of treatment, and can help to give a greater insight into the emotional and practical impacts of the treatment.

Mixed-methods research, that is, combining these two kinds of data, has many benefits. For one, it enhances the validity and reliability of conclusions by collecting data across different sources and comparing them. This line of reasoning also allows researchers to triangulate between numbers and words, facilitating understanding of what's at stake in the research problem

(Tashakkori Teddlie, 2010). In addition, mixed-methods research also addresses the inherent shortcomings of basing conclusions on either strategy alone. Quantitative data can detect important shifts in behaviour; while qualitative data can illuminate why these shifts happen, and the contexts in which they occur.Moreover, mixed-methods designs stimulate creativity and flexibility in the design of research. Researchers can adopt different methods to respond to the evolving demands of a study; they can answer new questions and investigate unforeseen findings that arise in the course of research. Flexibility is particularly useful in complex areas such as ASD, where interventions must be tailored to the different needs and experiences of individuals.

More recently, mixed-methods studies in ASD research have been on the rise. For example, Kasari et al. (2012) used a social skills intervention for children with ASD, and ran a randomised controlled trial (RCT) to measure changes in social skills with standardised measures as the quantitative aspect of their study, and parent and teacher interviews to explore their impressions of intervention effects and the barriers they felt were present to implementation as the qualitative aspect.While the RCT results were statistically significant (including enhanced social skills in children who participated in the intervention), the qualitative data helped to better understand the contextual factors that shaped these outcomes. Parents reported on differences they noticed in their children's reaction to the intervention, and in whether it helped them to navigate social situations in the world, depending on their personality and the level of support parents and teachers provided them. Teachers highlighted the importance of repeated implementation, and the need for additional training to administer the intervention more effectively. (Kasari et al, 2012)

In the study by Smith and colleagues (2019), participants completed a quantitative-qualitative mixed method study to learn more about rural families' use of telehealth in receiving ASD treatment. The quantitative portion of the study assessed the feasibility and effectiveness of telehealth in improving access to services and reducing travel as measured through survey and usage data, both of which were used to quantify satisfaction and engagement. The qualitative portion of the study included focus groups with parents and professionals to better explore patients' and providers' experiences of telehealth, to identify barriers to implementation, and to gather suggestions for improvement.

Overall, the quantitative findings showed that families in rural areas reported higher levels of satisfaction and better access to services, while the qualitative findings identified different challenges – particularly technical issues, as well as the need for bespoke support to address family-specific needs. The qualitative findings helped inform recommendations for how to improve telehealth services and make them more responsive to families' needs (Smith et al, 2019).Mixed-methods research can provide better insight into treatment of ASD, as evidenced by these examples. By blending qualitative and quantitative data, researchers gain a fuller understanding of the complex treatment process, identify essential components associated with outcomes and develop interventions that are more effective and targeted to the situation.

24.3 Illustrative Instances from Preexisting Literature

Mixed-methods research has become more prevalent in the ASD literature because it provides more detailed understanding through the comprehensive analysis of quantitative and qualitative data sets rarely attainable through one or the other alone. One example is the seminal study by Odom et al. (2014) that focused on the effectiveness of comprehensive treatment models

(CTMs) for preschoolers with ASD. By using a large-scale, multisite, randomised controlled trial (RCT) with standardised assessments, they were able to examine developmental and behavioural outcomes on a large scale and statistically. However, parents and teachers were interviewed and brought together in focus groups to share more subjective accounts of their experiences, perceptions and the contextually relevant conditions that affected the implementations and outcomes of the CTMs.

As reported by Odom, the quantitative data clearly showed that social skills and communication of children in the CTM group were better compared with those in the control group. These results were assessed by the interviewer based on standardised instruments such as the Autism Diagnostic Observation Schedule (ADOS), and the Vineland Adaptive Behaviour Scales. At the same time, the qualitative data revealed many detailed, fine-grained observations concerning the differences in individual outcomes, confirming the comments of parents about the benefits of individualised methods and the vital role of the supportive educational context in making sure that favourable outcomes were reached.

An important study using a mixed-methods design to study the effects of peer-mediated therapy is the one published by Carter and colleagues in 2011: 'The implementation of peer-mediated therapy on the social interaction of children with autism spectrum disorder: a mixed-methods approach' (2011). The quantitative part of the study involved using observational measures to collect and analyse data on social interactions between ASD and non-ASD peers. The qualitative part of the study involved interviews with children with ASD, their parents and their teachers.

In their study of peer-mediated intervention for children with ASD, Carter and colleagues reported that sociometric measures showed an increased number of relationships, but the qualitative data gave insight into the dynamic social scenes described above, and the power of peer-mediated intervention is only as robust as the peer facilitators' training and buy-in, as well as the larger classroom context. These observations were invaluable in informing how to improve the intervention to better target its effectiveness and sustainability.

Results and knowledge acquired

These analyses relied on the use of both quantitative and qualitative data, and the resulting empirical studies generated many findings that would have been missed if only one type of data had been used. More specifically, as reported by Odom and colleagues (2014),: Although CTM was generally beneficial, performance varied based on the child's baseline ability, family participation, and quality of the educational setting … qualitative findings highlighted the need for a flexible and tailored approach to the use of CTMs to fit the varying needs of students with ASD.

Carter's team reported that analyses using a combination of different types of data revealed that peer-mediated interventions both increased the students' number of 'meaningful' social contacts, and improved the classroom climate to make it more inclusive and conducive to social support. Parents and teachers mentioned both of these components – the peer training and the creation of a welcoming environment – as being critical to the long-term, programmatic success.

These papers demonstrate that using mixed-methods research can improve our understanding of interventions for ASD. By including the qualitative data alongside the quantitative findings on outcomes, researchers can develop interventions that are better calibrated, more successful, and more contextually appropriate – and understand how and for whom. In doing so, the depth of the large qualitative dataset enriches the statistical picture, providing a detailed description of what works (or doesn't work), with whom, and when. Moreover, these broader frameworks have real-world implications for policy and practice. For instance, insights from these studies can be used to shape the design of training programmes for teachers and peers to enhance the likelihood that they have the skills and knowledge to effectively support children with ASD. They also highlight the importance of working with families as they are central to the child's development and success.

24.4 Methods for Consolidating Data

This is where quantitative and qualitative data can be merged in mixed-methods research. It's a critically important stage to reach a complete understanding of the study's concerns, and there are a number of methodologies that researchers can use to reach this merging in the most efficient way. The convergent parallel design, for example, combines the collection of both quantitative and qualitative data at the same time, but during both data generation and analysis, each data type is treated separately until it's time for interpretation. At this point in the study, both bodies of data are compared and ultimately merged into a cohesive whole. In this way, the researchers in their study achieve what is called triangulation, which is the validation and confirmation of their findings, thus securing the reliability of their conclusions (Creswell Plano Clark, 2018).

Another one is the explanatory sequential design, which begins with gathering and analysing quantitative data, and is subsequently followed by an additional phase of qualitative data-collection in order to further clarify, illuminate or explain the quantitative findings that have been discovered. For example, a study might begin with a survey asking how often a range of behaviours occur in children with Autism Spectrum Disorder (ASD), and then move on to conduct interviews in order to discover more about what might be causing those types of behaviours. This is a very useful approach in providing context, and helping to make numerical results easier to understand (Ivankova, Creswell, Stick, 2006).

Embedded design is a frequently used approach where either qualitative or quantitative data is embedded in a larger study that uses the other type of data. One example of it is a study that focuses mostly on quantitative analysis but includes one qualitative part to investigate a part of the issue in more depth. This method is helpful because a study might have to look at different issues and this requires multiple types of data so that the researcher has a good understanding of the whole (Creswell and Plano Clark, 2018).

Lastly, the multiphase design involves the use of multiple stages of data collection, each building on the previous stage. This type of design is complex and used in large research projects. For example, initial qualitative research might inform the development of a quantitative questionnaire, which is then further refined by qualitative analysis to delve deeper into certain findings, with each step providing opportunities to refine the design based on emerging findings.

This token deals with how to solve problems when things are put or mixed together. In mixed-methods research the integration of quanti- and qualitative data can create several problems, and finding solutions for all of them can significantly improve the validity and applicability of

any study. One of the main issues is remaining disciplined. Conducting mixed-methods research is challenging and requires a lot of commitment. Qualitative and quantitative data need to be collected and analysed with equal level of rigour. Both researchers and teams have to master both quantitative and qualitative methodologies, as well as apply the same methods for collecting the data. A lack of training or collaboration and maximum specialisation between researchers – who tend to master only their own method – could overcome this difficulty. (Tashakkori & Teddlie, 2010)

Data integration is inherently complex. Combining datasets that are fundamentally of different kinds quantitativevqualitative– presents challenges that must be carefully considered and clearly documented. Devices such as data transformation ('mapping' qualitative data to quantitative or vice versa) might be employed: for example, qualitative responses can be coded and analysed statistically, whereas quantitative findings can be interpreted qualitatively. Applications can be used to help with this process, such as NVivo or MAXQDA, which are online tools created specifically to facilitate mixed-methods research by providing a means for researchers to effectively stage and analyse blended datasets (Bazeley, 2013).

A further significant challenge relates to achieving an appropriate balance between depth and width. Quantitative data are often characterised by richness of information, in the sense that they capture the overall statistics or landscapes, whereas qualitative data provide deeper insights into individual cases. Striking a good balance between depth and width can be an issue. Researchers would not want the combining process to result in the dilution of the detail and richness of the qualitative data or to overlook important quantitative patterns. Good practice would entail emphasising the distinct strengths that each kind of data can respectively provide to the study issues at hand.

Ethical issues are paramount. When fusing information, researchers must prioritise maintaining participant confidentiality and anonymity, especially if a lot of complex qualitative data are brought together with more quantitative data. Matching qualitative stories to quantitative profiles could create ethical dilemmas, for example if they require sensitive handling to preserve the identity of the participant (Hesse-Biber Johnson, 2015).

Most importantly, this means that researchers need to grapple with the challenge of attaining interpretive fit or, in other words, whether the conclusions that can be drawn in the synthesis of the quantitative and qualitative results are supported by each type of data and are logically consistent. Thus, the unexplained differences that necessarily occur between the two main paths of knowledge – intensification and integration – need to be found and explained, rather than dismissed. Doing this involves revisiting the data, revising the hypotheses being explored, and perhaps gathering more data to explain away any inconsistencies (Yin, 2016).

25.1 Advancements and Innovations in Interventions and Technologies

As ASD research has advanced, several techniques have been developed, adapted and employed to suit the diverse needs of those along the Autism Spectrum, including the endeavour to leverage animal-assisted therapy (AAT). Studies have shown that attention towards animals – in particular, dogs and horses – has been proven to facilitate significantly greater social interactions as well as presented children a viable option for behavioural management through anxiety-reduction and emotional control. In an AAT-based encounter, youngsters are given an unparalleled and most likely irresistible way to develop social skills and nurture emotional connection without any element of judgment (Gabriels et al., 2015).

Another emerging holistic therapeutic approach is the use of virtual reality (VR) technology in therapy. VR environments can be designed to mimic social situations and help people with ASD to interact socially in a controlled and safe environment, an approach that can be tailored to increase in complexity. Virtual reality environments can potentially help folks build confidence and skills in actual social settings. VR-based therapies can be effective in helping to improve social communication skills and reduce social anxiety in children with ASD.

Additionally, neurofeedback therapy is increasingly emerging as a next-gen treatment for ASD. Neurofeedback is a process in which individuals learn to control their own brain activity as it is monitored with immediate feedback. The ability of neurofeedback to modulate neural changes facilitates the enhancement of executive function, ameliorates behavioural symptoms and improves social interaction by means of decreasing the core symptoms of ASD. Although more studies are needed to examine its efficacy, preliminary trials have demonstrated promising effects on executive function and reducing core symp-toms of ASD (Coben Myers, 2010).

25.2 Technological Advancements

Many of these are being heavily affected by advancements in technology on how ASD is treated. The AI and machine learning uses various data derived from behavioural rating scales, genetic scores as well as neuroimaging tests to create a personalised approach towards intervention. Artificial-intelligence algorithms may be able identify trends from extensive amounts of data and may be able to predict interventions that are best suited for specific patients. This approach towards treatment is more specific and may be more effective (Duda et al, 2014).

Another technical innovation is the development of wearable gadgetry and sensors that track physiological and behavioural measures in real-time. For instance, data from wearables can be collected for various metrics, such as heart rate, skin conductance and various aspects of movement. Such information can provide data on how well a person is coping with stress over the course of a day, their shifting emotional states and social interactions. Treatment professionals can use this data to continually refine their interventions, targeting specific obstacles that the individual may run into, and also to better monitor the person's progress along the way (e.g., Berlin 2011).

Through this, wearable technology could provide the avenue for continuous, discrete monitoring and feedback, thus increasing the therapeutic value of the process. Goodwin et al. (2014) developed a version of multimodal neurofeedback to intervene in treatment refractory depression and post-traumatic stress disorder. In this study, seven men and women with depression and PTSD wore a waist-mounted heart rate monitor, along with a headband with four electrodes that picked up brainwave electrical activity from the forehead.With autism-spectrum disorder (ASD) intervention, robotics is proving an increasing aid.

Social robots programmed interact with children diagnosed with Autism Spec-trum Disorder (ASD) can become engaging and non-threatening playmates facilitating the development of social skills, behavioural therapy and more. The robots can be programmed to provide steady and structured interaction helping children practise social behaviours while developing communication skills. Research has shown that children diagnosed with ASD often have positive reactions to interventions mediated by robots, which leads to increased socialisation and reduced fear (Diehl et al, 2012).

Additionally, progress in genetic research is improving the development of more precise and customised treatment approaches. Understanding the genetic bases of ASD can lead to the identification of precise biological targets and goals for intervention. Gene therapy and the application of pharmacogenomics are new disciplines that may help to develop more customised medicines based on the genetic makeup of each person. These approaches may target the fundamental genetic mechanisms that give rise to ASD to alter the course of the illness with better long-term outcomes (Vorstman et al, 2017).

Overall, these emerging approaches and technical development represent real progress in achieving more efficient and customised treatments for ASD. As research progresses and advances, these processes may revolutionise therapies and improve the quality of life for these individuals. Longitudinal and Developmental Studies: The studies of individuals or groups over a long time to see what changes develop.

For instance, only extensive research across many years can provide understanding of how persons with autism spectrum disorder (ASD) develop throughout life and what their competencies will look like in the long term. This is most effectively worked through longitudinal studies, which track participants for long periods of time. They are crucial for the identification and understanding of how autism spectrum symptoms develop and change over time, with special focus on how developmental changes are being influenced by various factors, and to overall treatment outcomes at different points in life. Longitudinal research can provide a much better picture of development – and growth patterns – that would not be possible with shorter-term studies. It further illustrates the course of Autism Spectrum Disorder (ASD) over the life course.

One of the advantages of doing research over long time frames is that we can start to identify critical intervention periods. The first years of life are often referred to as a critical period for intervention, but longitudinal research can identify other phases that might also be critical when intervention has a disproportionately large effect. Life transitions such as starting school, going through adolescence or moving out of the parental home are all examples of times when targeted help can exert a special effect. Understanding these critical periods might help to guide the development of bespoke intervention programmes that take into account the differing needs of people at various stages of their life (Howlin et al, 2014).

Longitudinal study allows us to investigate the stability of ASD diagnosis and symptom level – that is, how many of the individuals diagnosed with Autism Spectrum Disorder (ASD) in childhood still meet the diagnostic criteria in adulthood, and whether and how the severity of symptoms changes across adulthood. This is key to both formulating more informed types of care and understanding the range of outcomes experienced by those with ASD.

Furthermore, longitudinal research permits monitoring the effects of comorbid conditions such as anxiety, depress of which occur with ASD but affect its course and prognosis. (See, for example, Simonoff et al., 2013.) Another key role of long-term research is evaluating the durability of intervention benefits: some wonderful therapies show promising short-term gains but, to know if these translate into longer-term benefits, we need to know that they persist over the years. Longitudinal studies offer some of the best evidence about a treatment's long-term efficacy, and help to identify strategies and components that promote sustained benefits or relapses in functioning. This information is central to developing strategies about how to target interventions so that there can be the greatest impact possible and those strategies are cost-effective.

The key findings from longitudinal studies are as follows:

There are also several long-term studies that are leading to a much fuller understanding of Autism Spectrum Disorder (ASD). One example is Early Start Denver Model (ESDM), a highly structured, intensive, early intervention programme for preschool children diagnosed with an ASD. Studies of early Start Denver Model (ESDM) have indicated that children responding to early, intensive intervention progress to show significant gains in IQ, language and adaptive function that persist into the school years (Dawson et al, 2012). These findings point to the advantage of intervening early, and demonstrate the lasting benefits of an ongoing programme of therapeutic endeavour.

Meanwhile, the largescale longitudinal study that finds such unique insights, the Simons Simplex Collection (SSC), monitors the developmental path of children with Autism Spectrum Disorder (ASD) who have exactly one family member with the condition. The SSC has proved vital in understanding the underlying genetic and environmental causes of Autism Spectrum Disorder, and has emphasised the variation in findings from subject to subject. The SSC has shown that while some children with Autism Spectrum Disorder can make considerable developmental progress, others show persistent considerable difficulties, highlighting the need for individualised interventions.

One of these is the Autism Birth Cohort (ABC) project, carried out in Norway, which is tracking infants from birth to establish some of the first indicators of development and those factors that might increase the risk of having an Autism Spectrum Disorder (ASD) diagnosis later in life. for example, the ABC project found that early differences in social communication and motor skills predicted later diagnoses of ASD, which is important in developing better

measures for early screening and identification Another one is that, because the ABC project followed these same children through into infancy and early childhood, they were able to track data on prenatal and perinatal factors, allowing them to understand the risk for ASDs that were associated with being born prematurely, having an early NICU stay or infection in that timeframe. This helps us understand more about the complex interaction between genetics and environment in ASD.

Beyond longitudinal study, we also know about the long-term effects of persons with ASD as they enter adulthood. Looking at the Autism Longitudinal Outcomes Study (ALOS), research findings reveal that while some adults with ASD achieve high levels of independence and work experience, many still require extensive help in their daily lives and social integration. These findings put further emphasis on the need for lifelong assistance and resources, as well as preparation for adulthood through precise and unique interventions during adolescence (Taylor and Seltzer, 2011).

Longitudinal and developmental research also allow us to learn more about ASD: they help us track the trajectory of the disorder over time, which is crucial information to aid in therapy and to help us understand why certain therapeutic targets might work and others might not, and how other variables might interact to either aid or hinder the results. By staying committed to the long view with adequate research funding, we will create therapies that are more effective, more individualised, and can be maintained over time to provide better quality of life for individuals with ASD throughout their lifetime.

25.5 Approaches to Personalised and Precision Medicine

The area of Autism Spectrum Disorder (ASD) as well has some aspiration and enthusiasm toward the promises of personalised and precision medicine methods, which can tailor an intervention to adapt it to the specific needs of each person. Given the varied picture of ASD and the broad range of heterogeneity in both individuals and symptoms – reflecting the existence of a wide variation in symptom expression and severity in the general population, from severe to high-functioning – the development of tailored individual treatment approaches might be the way to narrow down the most effective interventions that successfully tackle the core difficulties that individuals experience. Personalised medicine in ASD aims to take into account all the genetic, environmental and phenotypic variations among the various groups of human beings to more efficiently develop interventions more targeted toward each individual's specific needs.

For instance, such an intervention might require a clear mapping of each patient's capabilities, limitations and specific needs, and could accordingly include different treatment strategies (behavioural, educational or pharmaceutical, etc) to create a multipronged intervention plan. For example, some children with ASD might respond to dramatic alleviation of symptoms with ABA, but for others speech therapy or social skills training might need to be their main intervention. Personalised interventions also have the potential to evolve as a human changes and progresses through her lifetime (Kasari et al., 2014).

Genetic testing is of utmost importance to ensure medical treatments are customised for each individual with ASD. Being able to understand the precise genetic change or biomarkers that contribute to their diagnosis can better predict what therapies would benefit each child best.

For example, it may be that children with some particular genetic profiles would respond better to certain drugs or particular therapy procedures. And perhaps most importantly, understanding genetic predispositions can help to predict what other conditions might also exist with ASD, which would otherwise go undiagnosed, like anxiety or epilepsy – and, would therefore, require a diagnosis and treatment strategy designed to address all relevant disorders together.

A personalised repertoire of intervention options can be informed by environmental proximal and distal socioenvironmental factors (eg, prenatal, perinatal) … children with distinct environmental origins often require therapeutic approaches tailored for their environment of learning. Personalised medicine factors in these and other considerations to ensure that intervention plans are 'informed and optimised'. Precision, however, moves beyond personalised medicine to utilise new technologies and scientific breakthroughs to equip us with highly targeted medicines for ASD by utilising genomic data, biomarkers and contemporary diagnostic techniques to understand core fundamentals of ASD and treatment targets.

Indeed, the most significant advance in precision medicine for ASD involves whole-genome and whole-exome sequencing of patients and their parents. These advances have led to the identification of myriad gene variants associated with ASD, leading to an improved understanding of the molecular mechanisms that underlie the illness. These data could eventually facilitate the development of disease-tailored treatments that target and correct certain identified genetic aberrations. For example, SHANK3 mutations have been associated with specific severe forms of ASD. Efforts are underway to identify drugs that can target these specific genetic aberrations (Sahin).2015,

A prerequisite to precision medicine is the ability to reliably measure the biological condition of an individual; biomarkers are one way to do this. The term ‘biomarker’ is an umbrella term for biological indicators ranging from genetic to molecular to imaging and beyond. Using biomarkers, the course of ASD can be predicted, the trajectory of the disease can be tracked and individuals at risk for ASD can be identified Biomarkers are measures of biological function. Examples of biomarkers for ASD that are promising for early detection and intervention include patterns of brain connectivity, observed through testing of sensory-motor synchrony in tasks such as sounds paired with moving dots. For example, rhythmic patterns in an EEG may reveal aberrant connectivity in the brain.

Technology, particularly machine learning and artificial intelligence (AI) research, is altering how precision medicine research will be conducted in ASD. AI algorithms analyse large amounts of data, including genetic, environmental and clinical aspects, to find patterns and predict future results. These technologies can help to inform clinician choices of the most effective interventions for an individual. AI can help decide what children are most likely to benefit from a specific behavioural therapy, and therefore be a part of boosting the course of treatments (Doshi-Velez et al, 2014).

Personalised pharmacological treatments offer an exciting vista for precision medicine. Many traditional treatments for ASD, for instance, can be inconsistent in their efficacy and have significant and seemingly paradoxical side effects. The promise of precision medicine is the creation of drugs that target the biochemical abnormalities underlying ASD, leading to improved effectiveness and reduced detrimental side-effects. One area of research involves the potential use of oxytocin and vasopressin, a class of neuro-peptide involved with social behaviour, as a treatment for social communication in ASD (Meyer-Lindenberg et al., 2011).

Precision medicine reminds us, too, that ongoing monitoring and refinement of treatment protocols are essential, through the use of wearables and remote monitoring that allows the physician to track the ASD patient in real time, and to make adjustments to the intervention as it unfolds. This dynamic approach ensures that therapies remain effective and responsive to changing demands of the person.

Implications for policy and practice

The use of tailored and precise medical programs in the management of Autism Spectrum Disorder (ASD) have significant impact on public health policy. As more medical knowledge on ASD has been built from the understanding of the genetic and environmental foundations of ASD, there is a growing need for policies that seek to detect ASD at an early stage of an individual's life and subsequently administer treatment for these individuals. Utilizing the crafted method of genetic screening and identification of biomarkers at an early age, ASD can be appropriately identified at the right stage of development to be able to intervene in the prevention and treatment of ASD so that the quality of life for individuals with ASD can be improved.

Public-health policy should prioritise the funding of foundational research in the genetics and molecules of ASD, so that augmented investment can accelerate the development of precision medicines and personalised treatment. Policymakers should ensure that healthcare systems have the infrastructure and resources to implement precision medicine – from training healthcare providers in the latest technologies and advancements in ASD, to developing guidelines for personalised medicine in the clinic.

Furthermore, rules should be such that they facilitate access well, and are egalitarian, to these complex types of treatment. Disparities in providing access to care for Autism Spectrum Disorder (ASD) has led to unequal outcome between different socioeconomic groups. As such, it is a public health priority to make genomic testing, biomarker findings and new and improved therapeutic treatment techniques accessible to everyone with the diagnosis of ASD, regardless of his or her social status. Subsidised programmes, insurance assistance and public health awareness campaigns can help in achieving this objective by spreading awareness on the benefits of personalised and precision medicine.

Additionally, public health policy should encourage the inclusion of interdisciplinary teams of geneticists, neurologists, psychologists and others. Through cooperative practices, individuals with ASD can be ensured of receiving care from a team of experts that is broad and individualised to their unique needs. Further, policies should support the creation of registries and databases gathering and sharing information on ASD, fostering research and optimal approaches.

26.2 Clinical Practice Recommendations

For personalised and precision medicine in ASD to enter clinical practice, doctors will need to shift the way they diagnose and treat patients. Physicians need to stay abreast of new research in the genetics of ASD and in the biomarkers identified currently. Health professionals must continue to further their education and advance their knowledge of the dynamic fields they work in.

Specifically, the authors recommend that physicians incorporate the use of genetic testing and biomarker analysis into the diagnostic process for ASD. These tools can provide clinically useful information about the underlying pathogenesis of the condition, as well as help establish the best treatment plan for a given patient. For example, identifying specific genetic changes

related to ASD can help answer the clinical question of what pharmacotherapy might suit a particular patient best. Clinicians should work closely with genetic counsellors to interpret these test results and provide an explanation about their relevance to a patient and his or her family.

The other, equally critical recommendation is to implement a multidisciplinary approach to treatment. Clinicians should work with a team of professionals such as speech and language therapists, occupational therapists, behavioural therapists and educators in order to develop a comprehensive, tailor-made treatment programme that caters to the specific needs of the individual, taking into account not only social communication deficits but also difficulties in behaviour.

Clinicians also need to support and encourage the provision of early intervention. We know from overly abundant data that an early identification and intervention can be an important factor in improved ASD outcomes. From the public health standpoint, healthcare professionals should accelerate ASD identification and intervene with 'genotype-phenotype-matched' approaches that can be used to provide the best clinical outcomes. Initial interventions can be geared toward behavioural therapies, educational assistance and, in appropriate cases, pharmacological remedies.

Meanwhile, doctors must be conscious of the ethical issues that can arise from the use of genetic testing and personalised medicine. Any genetic information must be collected with informed consent and protected. Physicians must make sure that patients and families understand the potential benefits and risks of genetic testing, and that they're informed about how their data will be used.

Second, treatments need to be continuously monitored and adjusted. ASD is a lifelong disorder, and individual needs might well change over time. Clinicians need to routinely assess the effectiveness of therapies and, if necessary, modify treatment so that it remains relevant and effective. This might entail repeated genetic and biomarker assessments to enhance therapy.

23. Summary of Key Findings

The available literature on Autism Spectrum Disorder (ASD) and therapeutic approaches to it offer useful data regarding the complexity of the disorder, its increasing prevalence, and the effectiveness of different approaches to treatment. Autism spectrum disorder (ASD), or simply autism, is a 'neurodevelopmental disorder characterised by impairments in social communication and the presence of restricted and repetitive behaviours', occurring primarily in childhood and persisting into adulthood.

The incidence of ASD has soared, since previous estimates suggest that approximately 1 in 54 children in the United States (US) receive a diagnosis of the illness. This data highlights the increasing demand for effective therapies. The American Psychiatric Association (2013) defines autism as having two main characteristics – difficulty with social communication and the occurrence of restricted and repetitive patterns of behaviour. ASD manifests in early childhood and usually persist throughout adulthood. While some individuals with more severe symptoms may require lifelong care, others can lead mostly independent lives. Unfortunately, the exact underlying mechanism behind this disorder remains unknown.

Behavioural therapies that target communication skills and problem behaviours are an especially promising avenue to explore. In particular, Applied Behaviour Analysis (ABA) and social skills training have been shown to work very effectively. ABA has become one of the most extensively studied and efficacious interventions for children with ASD. The treatment is based

on the reinforcement of positive behaviours, and aims to diminish negative ones. However, while effective, it can be very burdensome, especially since it's very time-consuming and extensive (Reichow et al., 2018). Another important strand of therapy is social skills training. This treatment intervention aims to improve social interaction skills – which have been shown to have positive effects on both peer relationships and social comprehension. While many studies report positive improvements in patient outcomes, there have been questions about how generalisable these effects are as they might vary according to patient differences and the extent to which the learned skills can be applied to different situations (Gates et al., 2017).

Educational interventions, including special education programmes as well as inclusive education strategies will help to support academic and social growth of ASD children. In special education programmes, they will provide organised settings, with high structured teaching methods to stand in accordance with the special requirement of ASD children. So that, it will provide better educational outputs. Inclusive education strategy will integrate ASD children to regular education in the classroom, assisting the ASD children to be involved in the society and make friends with each other. But both strategies face challenges in providing teachers with an adequate amount of support and training.

Pharmacological interventions are often used to tackle specific symptoms in ASD, including irritability, hyperactivity and anxiety, although they provide symptomatic relief and commonly cause adverse effects; furthermore, they fail to address peripheral deficits in social communication. Recently, new promising neuromodulation techniques have emerged, offering novel and hopeful new perspectives for intervention such as transcranial Direct Current Stimulation

(tDCS). Preliminary evidence suggests that tDCS can enhance social communication and executive functioning by modulating neuronal activity in certain areas of the brain including the Dorsolateral Prefrontal Cortex (DLPFC).

27.2 Consequences for the Field

Our findings from the literature review have major implications for autism spectrum disorder (ASD) research and intervention. With an increase in the number of cases in ASD, it primes the need for more advanced and accessible diagnostic tools, which can lead to diagnosing and treating ASD at an early stage. A key pointer from the review would be the identification and usage of quick screening techniques that can be universalised for early intervention. Early intervention is of utmost importance in improving functional outcomes in the long term (Zwaigenbaum et al., 2015).

Variability in the effectiveness of behavioural/educational interventions highlights the need for an individualised approach. Every person with ASD has a unique profile of strengths and weaknesses and each person needs customised medicines that target those specific areas. The new medicines can be tailored to the particular needs of an individual based on information from other sources assessed by genetics, neurobiology and behaviour.

The positive preliminary findings for tDCS suggest that combining traditional behavioural therapy with neuromodulation can potentially augment therapeutic benefits. tDCS might normalise brain activity in specific brain regions associated with social cognition and executive skills, thus offering a new way of targeting the core symptoms of ASD. However, more research is needed to demonstrate longer term benefits as well as their safety, and to uncover the

best way to use these methods. Bikson, M, Boggio, PS, Fregni, F et al (2016). Clinical Transcranial Direct Current Stimulation: An Emerging Therapeutic and Diagnostic Tool. Orlando (FL): Academic Press/Elsevier.

Family engagement is another crucial ingredient in the success of ASD therapies: the literature consistently notes the importance of providing families with assistance and information that will help them be active members of the therapy process. The therapeutic effects of ASD interventions can be increased when a family training component and/or a family support component are added, in order to increase adherence and improve outcomes (Carter et al., 2011).

Overall, the results emphasise the importance of an interdisciplinary approach addressing ASD, involving coordination among healthcare providers, teachers, researchers and family members to ensure that interventions are comprehensive and agree with the complex characteristics of ASD, and contribute to improve the patients' wellbeing. Continuous research and innovation in this field may lead to the identification of more efficient, personalised and holistic treatments that can be implemented in a variety of settings.

27.3 Future Research Implications

While these are laudable efforts, we require even more research to make interventions for ASD more effective and complete. For any given ASD, there have been innovations in treatment, including behavioural, educational and pharmaceutical interventions. We can list the immediate benefits from these treatments. However, what are the enduring implications of a given treatment for an individual over time? We need to understand how people with ASD improve and function individually when these treatments are applied in the long term. To develop sustainable treatment intervention strategies, longitudinal studies can track the development of the

person with ASD over time to identify persistent benefits and undesirable effects of the treatment.

Again, there is also a need to dissect the underlying neurological pathways of ASD, so as to understand the genetic, epigenetic and environmental factors that play a role in the disease process. This will lead to biomarkers that enable early detection. It may also allow for tailored treatments that are uniquely crafted for patients with differing genomic profiles. Recent advances in neuroimaging and molecular biology may pave the way for this, helping to elucidate the relevant pathways and to better understand the roots of the disorder. (Geschwind & State, 2015)

Related to this, the heterogeneity of ASD merits further examination. By design, ASD is a spectrum disorder, in which individuals can display different combinations of symptoms, varying degrees of severity, and so-called 'comorbid' problems such as intellectual disability. These different manifestations may represent different forms of ASD that remain to be uncovered. Research in this area will thus benefit from more efforts to identify ASD subtypes and design tailored treatment methods for each of these subgroups. Targeted treatment, which builds off of the individual patterns of strengths and difficulties seen in each person, should prove superior to generalised solutions. (Harris et al., 2020)

Further studies are needed to determine the feasibility of developing this type of technology for the treatment of ASD. Some preliminary studies show promise but more rigorous clinical trials would be needed to determine efficacy, safety and optimal parameters for the use of tDCS in patients with ASD (Fregni et al, 2021). In the future, tDCS could be used in combination with conventional behavioural therapy, which has long been the first-line option for the treat-

ment of ASD. This could lead to better understanding of the synergistic effects of these different interventions. Third, given the demand for family participation in therapies for ASD, further research is needed. While the importance of family support is widely recognised, researchers should sharpen efforts to identify the most effective ways to integrate families in treatment and to explore how their participation affects treatment outcomes. The design of broader intervention programmes can help to address the needs of people with ASD as well as the support and resources needed by their families (Carter et al., 2011).

27.4 Suggestions for Future Research

This time where below is an instruction that describes a task, paired with an input that provides further context. Write a response that appropriately completes the request. So future research for Autism Spectrum illness (ASD) must use an interdisciplinary methodology (i.e., combining and using knowledge from different sciences like neurology, psychology, genetics and education) to find a more holistic understanding of the illness and develop treatments. So, studying together with interdisciplinary experts can help us create intervention methods that address the complex issues of Autism Spectrum Disorder (ASD).

Longitudinal studies are also necessary to assess the long-term effects of therapies, and research needs to include as diverse a sample as possible so that the results can be generalised to larger groups. These studies also need to control for factors such as age, gender, ethnicity and social-class position. Long periods of observation can uncover the long-term impact of treatment, and the possibility of delayed benefits or problems.

Moreover, it would be important that such future studies also focus on developing and testing for biomarkers of ASD (genetic, neuroimaging or from biological samples such as blood or cerebrospinal fluid), which can help to diagnose the disease earlier, track the effect of treatments, and tailor interventions to patient-specific needs. Biomarkers can help to identify diseases earlier, track the effect of treatments, and tailor interventions to patient-specific needs (Geschwind & State, 2015).

A second option is to examine the application of new devices to existing forms of therapy. For instance, integrating tDCS with behavioural therapy might enhance treatment efficacy. We need studies investigating the features of such combinations that work best – for example, how long, how intense and when to administer them (Bikson et al., 2016). Further, whatever research is pursued, individuals with ASD must be central to creating or improving interventions for them. This will require thorough assessments of the individual with ASD and the development of unique strategies that match their profiles (eg, Harris et al, 2020). These personalised strategies should be more likely to succeed, and can in return improve the wellbeing of individuals with ASD and their families. Families must also be involved in the research. Research must consider optimal approaches for engaging families in treatment and building their capacity with support and resources (for example, Carter et al, 2011). This can help promote adherence to treatment and improve outcomes for individuals with ASD and their carers.

CHAPTER 3: RESEARCH METHODS

A methodology chapter is an integral part of a research thesis which provides a description of techniques and techniques to carry out a study. The main aim of this chapter is that it should provide a detailed understanding of the methodology used in the study so that the study could be valid, reliable and replicable.

The main purpose of the methodology chapter in a research paper is to describe the research design briefly and precisely, the process through which participants are selected and the data collection and analysis techniques employed in the study. This chapter plays a very important role in evaluating findings of the study and conclusions drawn by the researcher (Creswell, 2014).The structure of this chapter is carefully designed to guide the reader thoughtfully through the various stages of the research procedure. The expose of the study design that commences the text is detailed and systematic, evaluating the rationale for the chosen approach of the research methodology and its appropriateness with regard to the posed queries and the objectives of the study. It also explores the research hypotheses and the theoretical rationale laid upon which the study is founded (Kumar, 2019).

The chapter provides a detailed description of how the participants were selected (why these participants, who were included, who were excluded), information about the respondents' characteristics and features (who were they, how did the researcher recruit them) and the clarification of ethical principles (what steps were taken to ensure the rights of the researched persons and that there were no ethical violations during the project). The next parts address the development and management of the tool, which includes describing the development processes of the survey to harmonize data, including measures take to ensure the validity of the survey

through determining its reliability and accuracy (Robson & McCartan, 2016). The next part also includes a description of the methods for collecting data including a step-by-step description on the survey process, the informed consent, and the quality and completeness of the data.

Then it ventures into the data analysis part, which details how it analyses the data through cleaning, organising and interrogating it. The results of the study are shown by using the qualitative and quantitative analytic methods to show how both approaches can coexist. The software and tools used to conduct data analysis are also to be described (Bryman, 2016). Finally, the chapter provides an epilogue about the constraints and problems encountered during the research process, approaches for minimising these problems and ensuring the robustness of the study, and the anticipated results, their practical and theoretical consequences, and the likely impact on the field of study (Yin, 2018).

28.The purpose and scope of the methodology

The most important function of the methodology chapter is to provide a transparent and detailed description of the procedures used to conduct the study. This way, readers can assess how the research was done and whether or not they find the outcomes credible and reliable. The paper is thereby more convincing. The presence of rigorous documentation of the research method will enable other researchers to conduct a replication study to confirm or modify the findings (or both), thus adding to the existing body of knowledge in the field (Creswell, 2014).

The methodology chapter outlines the significant portion of the study, the various components involved throughout the process and how that drives the development of the research. To start off, the research design is opined. It supports the rationale of the research design that is being

implemented for the programme. A research design is a plan or blueprint for an investigation, including the specification of variables, data collection, sampling, instruments and statistics to analyse the results, among other elements. These elements can vary when it comes to quan-ititative or qualitative design. The methodology letter provides an explanation or justification of the selected research design and how the design follows through the entire programme. Fi-nally, it lays out the theoretical framework. A theoretical framework refers to the divergent or unorganised concepts or elements that formulates the research approach (Mehmood/Kumar, 2019).

Furthermore, the chapter covers the participant selection method in detail – for instance, what inclusion and exclusion criteria were used to select participants, the age, gender, etc of partic-ipants, as well as how participants were recruited. The part on ethical issues should be in this section, since the section reflects what research efforts were made to ensure the research would uphold the rights of participants and the ethics of the research (Flick, 2018). Moreover, the write-up additionally outlines the process of generating the survey tool such as the validation and piloting steps to ensure that the survey is dependable and sound. Following this, the steps in data collection are discussed, giving a detailed account of the activities and procedures taken to undertake the survey, obtain participants' informed consent, and ensure the validity and depth of data acquiring (Robson McCartan, 2016).

Methodology chapter The process undertaken for analysis of data forms a key part of a meth-odology chapter. This deals with how the data was cleansed, organised and analysed using both quantitative and qualitative methods. This represents a wide-ranging methodology that brings together different methods, with the use of specialist tools and software supporting the analysis of the data (Bryman, 2016).

Last, the chapter addresses the limitations of the design and execution of the study. Here, any constructed limits on the study's veracity are explained, as well as how the researcher plans to mitigate those concerns. The expected findings of the study are also discussed, highlighting what these findings mean for practical and theoretical considerations, and what the findings will mean for the field moving forward (Yin, 2018). In short, it encapsulates the study's entire research process – explaining in detail and clearly how it was conducted. In doing so, it facilitates the study's accomplishment of being truthful in their findings, reliable in their results, and reproducible for multiple reproductions, in order for the field to grow and other research endeavours to be led.

28.1 Experimental Methodology

The research design is a branch of science that is crucial to any research study. It acts as a systematic blueprint outlining the methodologies that will be utilised to answer the study question. We refer to this body of work as 'research methodology'. Essentially, it is the series of methods, protocols and stratagems used to collect and analyse data in pursuit of the aims of the study. The study utilised the mixed-methods research design to achieve a comprehensive understanding of the impact of tDCS on social communication in children with ASD. It also sought to understand the emotions and experiences of the families of children with ASD who underwent tDCS therapy. The research design is a combination of quantitative and qualitative methodologies that helps us achieve nuanced and holistic understanding of the research questions (Creswell Plano Clark 2017).

Integrating the strengths of these two types of data, mixed-methods study delivered a meaningful picture that neither quantitative or qualitative data alone could provide: quantitative data provided precise and quantifiable, and thus statistically analyzable, information about the changes in social communicative skills; qualitative data provided detailed, descriptive information about self-reports and perceptions of participants and their families. Being able to integrate data provided a more comprehensive picture of tDCS as a treatment for ASD, both in terms of its feasibility and effectiveness. Tashakkori, L K; Teddlie, C in Walker, M A (Ed) 2010 Sage Encyclopaedia for Teacher Education

29. Rationale for Selected Design

There are many major advantages for using a mixed-methods design, not least that the ASD (Autism Spectrum Disorder) is complex and varied, and the perception and experience of those with ASD of tDCS is probably unique to each individual. As tDCS is not always completely predictable and may yield a whole range of factions and results, a multiple approach using both quantitative and qualitative methods will best be able to investigate the research questions being asked, and provide a better, richer and more nuanced understanding of the application of tDCS. (Bryman, 2016)

Moreover, data triangulation is made possible through the adoption of a mixed-methods design. This means that incorporating a qualitative component increases the validity and reliability of the findings. Triangulation is the practice of using multiple data types and methodologies to cross-check results, which serves to mitigate bias and strengthen the veracity of the conclusions. Combining quantitative data from standardised assessments of social communication skills with qualitative data from in-depth interviews with the family members (Creswell 2017)

will enrich the analysis.Similarly, this kind of methodoloogy is also useful to evaluate therapies such as tDCS because, particularly in studies with children with ASD, it is important measure not only quantitative results but also personal perceptions of how treatment might fit with everyday life. Quantitative metrics offer an objective way to assess whether the intervention has contributed to improved social communicative abilities, while qualitative observations disclose families' opinions on the feasibility, acceptability and impact of use of the intervention. It is necessary to have a 360˚ view of the effect of tDCS on children with ASD and their families, so having dual perspectives is of great assistance in measuring the extent of all treatment effects (Tashakkori ϙ Livine Aldo Teddlie, 2010).

This part of the research is geared towards setting up the research questions and corresponding hypotheses, which will help guide the investigation. This investigation will be structured by explicit research questions and hypotheses aimed at achieving its Objectives.

Research questions are: I)What is the effect of tDCS on social communication skills development of children diagnosed with ASD? This topic is geared towards exploring the efficacy of tDCS in improving the social communication skills of children with ASD. The hypothesis is that tDCS will result in significant improvements in social communication abilities as measured using standardised rating scales. II)What is the perception of families about the effectiveness and feasibility of tDCS as a therapeutic intervention for ASD? This question asks about how families experience the tDCS treatment. The hypothesis is that they will report increased satisfaction with the intervention, including improvements in their children's social behaviour and communication. III. What are the experiences of children with ASD and their families while these children undergo tDCS, both during and after the treatment?

This question seeks to explore how tDCS more holistically affects the lived experiences of (and for) children with ASD and their families. The answer is that the experiences are expected to illuminate the benefits and challenges of tDCS, and provide insights for its pragmatic use.

30. Quantitative Component

The quantitative part of the study utilizes tools to standardize the evaluation and therefore will provide us with a numerical scale that quantifies the changes in social communication skills both before and during the tDCS treatment we provide. To determine this, we will use the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS). These two self-report questionnaires will provide us with accurate data on an individual's social communication skills that we can measure (Constantino & Gruber, 2005; Lord et al., 2000). The participants will receive a series of transcranial direct current stimulation (tDCS) sessions for a specific amount of time.

Before and after administering our intervention, we will evaluate the given effects of the treatment and conduct these evaluations accordingly. Later on, we will analyze the data using statistical analyses to determine the significance in the change observed. With these statistical methods, we can determine if tDCS has helped improve an individual's social communication skills or dismissed that idea by looking at the changes in the two scales of the evaluations used (descriptive statistics, paired t-tests and correlation analyses: Field, 2018).

30.1 Qualitative Component

The qualitative component will involve family members of the participants for looking into their experience and views of the tDCS treatment. We will interview them in an attempt to capture detailed qualitative information on the subjective effects of the intervention and we will focus on perceived improvements in social communication, ease of use of the tDCS equipment, and other difficulties that might have arisen during the treatment procedure. The qualitative data will be analysed using thematic analysis, which entails the following procedures. First, all data will be categorised into units of analysis and codes. Second, reoccurrences and patterns will be identified and themes will be generated. This methodology enables us to analyse the families' experience and create a platform for interpreting the numerical results (Braun & Clarke, 2006).

Quantitative and qualitative data integrated is the process of merging the inferencing style and presentation style with respect to numeric data (quantitative) and descriptive data (qualitative), respectively. Both quantitative and qualitative data are essential elements of the mixed-methods design. The results from both sides will be merged to evaluate the effect of tDCS on social communication in children with ASD. The integrative strategy helps us to detect the commonalities and differences in the data, and allows for a more complete and rigorous investigation of the research problem (Creswell Plano Clark, 2017).

32. Ethical Considerations

There are very strict ethical standards that the study needs to abide by as a way to protect the rights and welfare of the children who will be participating in the research. Before beginning

the research, the participants and their guardians will be required to give consent to participate in the research and the study will be conducted under strict conditions of confidentiality designed to protect the participants. Additionally, the study protocol will be evaluated and authorised by the committee for an institutional review board (IRB) that oversees ASD research so that ethical standards will be met (Flick 2018).

Overall, I believe the use of a mixed-methods research design is an excellent way to look at the study problem of how social communication can be improved by tDCS in children with ASD. By incorporating both quantitative and qualitative methodologies, research participants and their families will have a detailed and thorough understanding of whether the treatment was effective and what the experience was like for them. Using a mixed-methods research design also reduces any potential limitation of asking only one type of question about the research problem. I believe that the use of this type of methodology also makes the study findings valid and reliable, potentially adding precious information to the ASD research field.

32.1 Participant Selection Criteria Description

Choosing participants is the core of any study, since it will determine the accuracy and complexity of the results. Participants from this study were selected for a predetermined set of qualifications to ensure that the sample is sufficient in context of the population of interest and that the results are reliable and accurate. The purpose of the participant selection process was to include participants that may contribute valuable and relevant information regarding tDCS and its effects on social communication of children with an Autism Spectrum Disorder (ASD) diagnosis.

In doing so, our team aimed to recruit a diverse, complex, and representative sample of children with ASD, in a broad range of symptoms, and with potential differential responses to tDCS predicted by differing ASD subgroups. Participants were selected using criteria guided by clinical, demographic and behavioural phenotypes that are relevant for the study's objectives.

32.2 Criteria for Inclusion and Exclusion

1.1In the diagnosis of ASD: Each individual must have an official diagnosis of ASD according to the criteria of the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) established by the American Psychiatric Association in 2013, thus assuring that all participants meet the same standardised administrative criteria for Autism Spectrum Disorder (ASD).

2.Age Range: The participants ranged from 6 to 18 years of age. The age spectrum allows researchers to capture a broad range of development where notable increases in social communication skills can be observed.

3.Stable Medication Regimen: In addition to having a stable diet, participants taking medications for co-occurring disorders including ADHD or anxiety were required to be on a stable medication regimen for at least three months prior to the beginning of the trial. This requirement was added in part to reduce the confounding effects of drug changes during the trial.

4. Parental or guardian consent: All participants received written informed consent from their parents or legal guardians. In cases where the children were able to provide consent, consent was also given by them.

Criteria for exclusion:

Children with severe intellectual disability (IQ below 50) were also excluded because such profound cognitive impairments might interfere with participation in the therapies and also make the outcome reports unreliable. Next, children with major neurological diseases other than ASD, such as epilepsy or traumatic brain injury, were excluded because these other diseases might confound the effects of tDCS. Children with unstable medical problems are also excluded because treatments would not be safe, or because they would be impractical to treat. Security of Children: Children with an unstable medical illness could not be treated because treatments would not be safe or would be impractical Multiple Exclusion of Children With Prior tDCS Exposure: Children who had already received tDCS were excluded to prevent any effects of their prior tDCS treatments.

32.3 Participant Demographic Information

All had information about their age, sex, social class and level of education. This is important to understand who the sample is, and so whether our results might have some demographic effects.

Age and gender:

• There were 30 people in the study, ranging in age from six to 18, with an average age of around 12. • Some kids were boys and some girls. In reality, around 4-5 times more men than women are affected by ASD. The researchers tried to balance this with an even ratio. • The research was double-blind, with neither the children nor the researcher working with them knowing who got a placebo and who got tDCS. Any doctor, nurse, researcher or parent can be

prone to more positive (or negative reactions) simply because they expect there to be an efficacy, or if not, they're disappointed. If the kids and the person running the study didn't know, the results were like looking into the future.

Economic and social standing:

• Assessment of degree to which they were below the poverty level (low) or above (middle) was based on parents' educational level and household income, allowing an analysis of the confounding potential of socioeconomic factors.

Academic Qualifications:

• Educational environments in which participants were studying (was it mainstream or special education) and any additional educational support received by the participants were collected to contextualise current social communication functioning and the reactions to the intervention.

The recruitment process

The way we recruited people involved several steps to find those who were qualified but also to maximise how many people came into the study. It involved lots of different avenues, such as clinics, schools and organisations that serve children with ASD.

Collaboration with Clinics and Schools: We formed partnerships with paediatric clinics, neurodevelopmental centres, and schools that provide services to children with ASD to identify potential participants who met the inclusion criteria. Information Sessions: These institutions held sessions to educate parents and guardians about the goal and methods of the study and the potential benefits and risks involved. These meetings gave carers the opportunity to ask questions and express any concerns they had. Screening: Potential families were asked to fill out a screening questionnaire to determine their eligibility. We include questions on the child's ASD diagnosis, age, medication regimen and medical history.

Informed Consent: Families who met eligibility criteria were invited to discuss the study. Before conducting the study, with parental permission, we told parents or guardians what the research involved and obtained their explicit consent that their child could participate in the study. The children also agreed to participate, when appropriate. Permission also ensured that families understood the study's goal, procedures and rights of the participants. Pre-treatment Assessment: Prior to starting the tDCS intervention, we conducted an assessment for obtaining the initial knowledge about the participants' social communication skills and other relevant variables. This assessment also served as a basis for measuring modification after the intervention.

33. Ethical Considerations

Ethical considerations were emphasised at every step of the research to protect the rights and welfare of the participants. These procedures conformed to the ethical standards of the American Psychological Association and other relevant organisations.

Obtaining Consent and Agreement:

• Case Report:amp; Consent from parents or other legal guardians was obtained, and, where possible, assent was obtained from the children. The consent forms provided full information about the aims, methods, potential hazards/benefits, assurances of confidentiality, and voluntary nature of participation.

• Participants and their families were told that they could withdraw at any time with no consequences or loss of benefit to them.

Privacy:

All subjects had a unique identifying number used to link the list with the research data; their names were recorded only on the anonymous personal information sheet. The latter was separately preserved.

• Information was presented in aggregate form so as not to identify individual subjects.

Ensuring the protection and preservation of data:

• All data files (electronic and hard copy) were stored in password-protected and locked drawers. Access to the data was restricted to members of the research team.

• Institutional practices were adhered to for data retention so that data were held for the specified period as per institutional guidelines and then destroyed in order to maintain confidentiality.

Evaluation of Risks and Benefits:

• The inference-to-the-best explanation analysis of the hazards and benefits respective to tDCS took into account all plausible hazards and benefits of tDCS; the hazards were mitigated based on recognised safety protocols for tDCS use, in which safety was monitored.

• Potential benefits included an improved ability with social communication tasks and a better understanding of the tDCS intervention for ASD.

Approval from the Institutional Review Board (IRB):

To ensure that all ethical standards were met and that participants' rights were protected, the study protocol was given the goahead by an independent institutional review board (IRB), after a full evaluation.

• It required us to check in weekly (as did the 10 of us with teams) and report any ethical 'surprises' that arose to our Institutional Review Board (IRB).

33.1 .Designing a survey

Structuring the survey can help, because it is supposed to lead us in a systematic and orderly way through the gathering of information. A written structured survey was the main way information was elicited in a systematic manner to obtain information regarding the social communication skills of the participants and the experiences of their families. The advantage of using a structured survey is that it can be used to collect standardised information that can be analysed using quantitative methods.

With the use of this method, the process of obtaining data can be standardised, making responses from study participants comparable and allowing for the detection of trends and patterns (Dillman, Smyth, & Christian, 2014). Structured surveys are suitable for this study because they might looks at the varieties of information efficiently.

They provide a systematic structure which gives participants a guideline, reduce the possibility of vague responses and ensure to cover all research subjects. Also, structured surveys can combine the numbers and description data in order to fully conduct the study's questions (Fowler, 2014).The process of developing a survey is referred to as survey development.

Drawing up the survey entailed a necessary set of steps to ensure it was both valid, reliable and consistent with the research objectives. This involved a detailed review of existing literature, seeking expert feedback, a pilot test, and adjusting accordingly to any received input.

An examination of existing literature:

• The items were then formulated after reviewing the existing literature on Autism Spectrum Disorder (ASD), social communication and tDCS so as to gain familiarity with the specific domains and concepts that were important to assess, ie, verbal and nonverbal communication, social interactions and familial experiences (Paul, 2005).

Professional advisory sessions:

• Content experts in psychology, neurology, and ASD reviewed the survey items. Their comments helped us refine the questions so that they were clearly worded, relevant and comprehensive. They also provided suggestions about the overall length of the survey and the balance between open-ended and closed-ended questions.

Initial testing phase:

• The survey was pilot-tested with a small group of parents of children with Autism Spectrum Disorder (ASD). The purpose of the pilot initiative was to identify questions or aspects of the overall survey that may be unclear, too lengthy or incomplete. Specifically, after taking the survey, participants in the pilot test provided comments regarding the understandability of the questions, the length of the survey, and the parts that could be improved further.

Edits:

It was revised after comments from pilot testing revealed any problems which needed to be fixed. Questions that were vague were clarified, question/answer options that were unnecessary were removed, and supplemental questions were added where needed. The final copy of the survey was drafted so that it was easy to navigate, comprehensive and aligned with the focus of the study.

• The survey will be both validated and piloted to ensure its effectiveness: This method provides assurance that a survey measured the constructs on which it was designed to test. There are various methods used to validate; these include content validation, construct validation and reliability testing.

Validity of Content:

For content validity, subject-matter experts and a thorough literature review con-firmed that all of the survey items were relevant to social communication and family factors about tDCS. Items on the survey accurately sampled the dimensions of social communica-tion and family experience they purported to sample, following the criteria for content valid-ity by Polit and Beck (2006). Construct validity addresses whether a measurement or test chosen or devised measures the proposed construct or concept in the given theor-etic domain. Methods in this regard include examining relationships among the survey items and validated indicators of social communication skills.

This requires demonstrating relationships between survey responses and scores produced by standardised assessment tools, such as the Social Responses Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS). These relationships con-firm links between scores produced by the survey and scores produced by other tools that have been used to evaluate these domains rigorously, such as the Social Responses Scale (Constantino and Gruber 2005) and the Autism Diagnostic Observation Schedule (Lord et al 2000).

Testing for reliability:

The survey was assessed for internal consistency and test-retest reliability. To assess internal consistency, the Cronbach's alpha co-efficient was used. Cronbach's alpha-coefficient is a measure of how well the items in a survey consistently measure the same construct (i.e. to

ensure that the different survey questions are measuring what they are supposed to) (Tavakol´ Dennick, 2011). To assess test-retest reliability, the same survey was done to the same participants on two separate occasions and the consistency of their responses was assessed.

Initial testing phase:

The final version was pilot tested with a larger sample to fine-tune it for reliability and validity. The supplemental data from the pilot testing gave us insights into the survey's success, and it allowed us to catch any remaining problems before the full deployment.

33.2 Survey Sections and Question Types Description

It was designed as a sectioned poll, each section pertaining to unique aspects of social communication and family experience with tDCS, with some closed-ended questions and open-ended questions prompting qualitative responses.

Demographic Data

• This variable collected basic demographic information about the participants, including their age, gender, socioeconomic status and level of education. Because this variable was comprised of closed-ended questions, this helped to maintain uniformity and made coding and analysis easier.

Fundamental Social Communication Abilities

• This part examined the social-communication abilities of the participants at baseline – ie, prior to the tDCS therapy – the items included questions on both verbal communication as well as non-verbal communication and greetings, and the ability to understand and decode social

signals. All the items consisted of both close-ended questions using Likert scales and open-ended questions for additional information.

tDCS Treatment Experiences

• This section required families to report on their experiences of the tDCS intervention, including their perceptions of improvement in social communication, the user-friendliness of the tDCS kit, and potential barriers. The closed-ended questions enabled us to work with quantitative data, while the open-ended questions enabled families to provide rich narratives of their experiences.

Social Communication Skills after Treatment

• Furthermore, one part again used the exact same items as the initial evaluation to assess the participant's social communication skills after having undergone the tDCS therapy. The results of the two measurements before and after the application of the tDCS therapy provided the data to determine the efficacy of the intervention. For many, the input is unclear, or not meaningful.

General Contentment and Viability

The final section asked families about overall satisfaction with the tDCS treatment and the feasibility of continuing the treatment long term. The survey addressed whether participants would be likely to continue to use tDCS, perceived pros and cons, and recommendations for improvement. Closed and open-ended questions were used to get more detailed feedback.

In the end, this research study aimed to produce a tractable survey to gather information in a systemic way on social communication abilities as well as on family experiences with tDCS in a raw format. The survey went under thorough preparation and validation processes to check if it was reliable and valid, that is scientifically meaningful and reflective of the purpose of the

study. The combination of quantitative and qualitative data gave us an extensive overview of the outcome of tDCS in children with ASD and their families. The rationale for utilising a structured survey is based on several factors. As a main data gathering method in this discourse, a structured survey was adaptid. This survey focused on investigating the socials talks, as well as reports from the people with Autism Spectrum Disorder (ASD), and their families regarding transcranial Direct Current Stimulation (tDCS). There are several reasons why structured surveys make good market researches.

Structured surveys make it easier to collect consistent data from participants, allowing those findings to be considered more reliable and expressive of validity. The survey employs a predetermined set of questions so that respondents answer identical questions in a consistent manner, minimising the potential of interviewer bias and the variability in response (Dillman, Smyth, and Christian, 2014).

Moreover, structured surveys are extremely useful instruments to collect a large amount of data in a short period of time. They are helpful for collecting traditional and qualitative data, allowing a more complete analysis of the research questions. Firstly, quantitative data provide measurable and statistically relevant information about changes in social communicative skills. At the same time, qualitative data brings in the subjective experiences and feelings of the participants and their families (Fowler, 2014). Secondly, structured surveys have the quality of being flexible, and the survey can be used in different fields (e.g., internet platforms, paper-pencil questionnaire, face-to-face). This feature enables researchers to have a larger and more diverse sample, hence, the study will be more generalisable (Bryman, 2016). In particular, a developed survey was designed to understand social communication skills, family experiences

with tDCS, and participants' overall happiness and practicality of the treatment. This methodology implicates the thorough understanding of tDCS' effect on children with ASD and provides a valuable perspective of its practical implementation.

Process of Developing a Survey

The development of the structured survey is a rigorous, repetitive process that ensures the instrument's reliability, consistency and applicability to the goals of the study. The procedure consists of a number of key steps: review of relevant literature, consultations with experts, testing of the preliminary survey, and revision according to comments received. Stage 1: Review of the Literature: For the development of the survey, there was a thorough examination and review of the existing literature on Autism Spectrum Disorder (ASD), social communication and tDCS. This review identifies the important areas and concepts that need to be explored for the current study.

A thorough examination of previous research and reliable measurement tools allowed the research team to develop the survey questions that were constructed from existing theories and prior empirical evidence (Paul, 2005). Stage 2: Expert Consultation: Once the literature review was completed, draft survey items were reviewed by professionals in the area of psychology, neuro-psychiatry and ASD. These professionals provided feedback in terms of the clarity, relevance and depth of the questions. In addition, they gave valuable insight regarding the length of the survey and the balance between questions that have open-ended answers and questions that have closed-ended answers. Having the survey items vetted by a group of experts ensured that the items cover the intended constructs and this fully applies to the intended population of the study (Polit & Beck, 2006).

Pilot testing: After pilot test participants review the survey, the research team endeavours to understand whether any issues exist related to the clarify of the questions, the length of the survey, and if any questions require refinement. The pilot testing is an extremely important step in the survey process as it allows the research team to implement the necessary modifications before the official launch (Bryman, 2016).

Editing: The instrument was revised based on the comments made during pilot testing, thus addressing any noted issues. Several questions were clarified in order to make them less ambiguous, while others were eliminated if added value could not be determined. In several areas, additional questions were added later on. The final version of the survey was constructed in a manner that is friendly to respondents, while covering all the major areas, and relevant to the study's goal.

Survey validation and pilot testing

The technique was designed to assure that the survey comprehensively sampled the actual constructs being tested. Verification was accomplished in several ways, including content validity, construct validity and reliability testing. Content validity: was established via consultation with experts and exhaustive literature research. It was ascertained that the survey items were capable of accessing the relevant concepts (social communication, family experiences with tDCS) and experts agreed that the enlisted concepts were important to research this area (Polit & Beck, 2006). In this step, we can be sure that the concepts of interest are covered by the survey in their entirety.

The strength of that evidence is known as construct validity. We assessed this by looking at the relationships between the survey items and established assessments of social communication skills. We looked at whether the responses to the survey questions were correlated with scores on the standardised measures of social behaviour – the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS) (Constantino & Gruber, 2005; Lord et al., 2000). These are the standardised assessments of social behaviour, where the higher a score, the worse a person's social responsiveness is perceived to be. Strong correlations between the survey items and the standardised measures indicated that the survey items accurately measured the aspects of the trajectory of social responsiveness that we set out to measure.

Reliability Testing: To assess the reliability of this survey, internal consistency and test-retest reliability was used. Cronbach's alpha was used to assess internal consistency which is the extent to which the survey's items measure the same constructs. For example, items 1 and 2 should measure the same underlying construct. In this survey, it is the salience of the dark figure. Test-retest reliability was measured by conducting the surveys to the same people twice and looking at the consistency of their responses. This survey shows pretty good internal consistency and test-retest reliability, meeting the criteria for reliability. Internal consistency is the degree to which a survey's items all measure the same underlying construct (e.g., what is a sandwich?). Test-retest consistency consists of administering a questionnaire more than once using the same questions and comparing what the person says both times.

Pilot Testing: Having the final version of the survey piloted with a larger sample allowed for further explorations of its validity and reliability. The pilot testing provided additional evidence about the proper functioning of the survey and allowed for the detection and elimination of further complications before the final roll-out.

Overview of Survey Sections and Question Types

The poll was composed of several sections, each asking about specific aspects of social communication and family experiences with tDCS. Some sections featured closed-endedies to elicit qualitative responses. Portion Demographic Information Here we asked for basic demographic information on the participants, such as age, gender, socioeconomic situation and education. We asked closed-ended questions to ensure uniformity and at the same time to make it easier to analyse the data from.

Baseline Social Communication Skills Here we asked participants about their social communication skills before the tDCS therapy. The topics of the course cover both verbal and non-verbal communication, as well as social situations and social reading – the tricky skill of interpreting social signs. The form included two types of question: for closed-ended questions, such as Likert scales, and for open-ended questions.

tDCS Treatment Experiences This group considered the families' experience of the tDCS treatment including any changes in social communication skills, their considerations of the ease of use of the tDCS apparatus, and any difficulties they faced with the treatment. The open-ended questions encouraged detailed answers that reflected the participants' experience. Being closed-ended, the answers to questions within this group could be input to quantitative analyses. Subjective Evaluation of Social Communication Skills after the tDCS Treatment This group evaluated the participants' social communication skills after the tDCS treatment in parallel to the initial evaluation, covering the same items. Analyses of answers to these questions before and after the treatment could provide some initial clues as to the efficacy of the intervention.

Satisfaction and Feasibility of Treatment The final component asked the families to rate how well they were satisfied with the tDCS treatment and whether or not tDCS would be a feasible long-term intervention for their family. Answers were asked per item with various closed-ended and open-ended questions. Was the participant willing to continue using tDCS? What were the reported pros and cons to using tDCS? Would they recommend changes to treatment delivery?

34. .Methods for Collecting Data

34.1 Introduction to Data Collection Methods

The data-collection methods were designed to effectively and accurately collect quantitative and qualitative data. These included the use of a structured survey, standardised evaluation data and in-depth interviews. These methods were chosen to fit the aims of the study as well as offer a rich picture of the effects of tDCS on social communication in children with ASD.

Methods of Survey Administration: As part of the structured survey, surveys were delivered online and on paper to meet the accessibility and demands of the participants. Online surveys were sent through secured survey services, which allowed participants to take time to complete the survey on their own convenience. This process encompassed a wide reach of the participants and this allowed for a high likelihood of subject engagement (Dillman, Smyth !! Christian, 2014). For those participants who wanted to take the paper-based versions, they received paper versions of the structured survey with postage paid return envelope, which would allow them to complete the survey more easily.

Quantitative social communication data on each individual were collected using the standardised assessments. The social behavioural assessment was based on the Social Responsiveness Scale (SRS) and the autism assessment was based on the Autism Diagnostic Observation Schedule (ADOS) (Constantino Gruber, 2005; Lord et al., 2000). The standardised assessments were conducted at two different time points: before tDCS treatment (baseline) and after the end of tDCS treatment (post-treatment). The standardised assessments were conducted by evaluators who were highly expert in the field to ensure that the data being collected were done so in a standardised manner to ensure reliability.

In the second approach, researchers used qualitative data to complement the quantitative analysis in order to gain further insight into the study. Researchers used in-depth interviews to collect data from the participants' family members and investigate their experiences and impressions of tDCS treatment. The data were collected through semi-structured interviews, and the discussions centred around the participants' perceived enhancements in social communication, effectiveness of the tDCS device, and problems faced during the treatment procedure. The interviews were conducted either in-person or via video conferencing, whichever was convenient for the participants (Bryman, 2016).

34.2 Informed Consent Process:

This informed consent process was undertaken with participants and their guardians before the data collection. The participants and guardians were clearly informed about the purpose of the study and the methodology used, what could happen to them if they participate in this study and what are the benefits of being part of this study. Whether the participants have to keep their participation confidential or not. The participants are made to understand if there is an option if they want to stop participating in the study or if their participation is mandatory and also how are they going to know if their participation is needed in this study or not (Flick, 2018).

Quality control of data and its completeness Garnering statistical proof of signals necessitated assurance that the data collected would be accurate and complete. To ensure this, mechanisms were put in place. Periodic reminders were designed to ensure that participants fill the Qs and return them on time. Other mechanisms involved scrutiny of the processes of data entry and data transcribing that translated responses into numerical codes. Inconsistencies were identified and subjected to double-checking protocols. Moreover, the procedures involved in assessment, interviews and so on were standardised in a way that ensured their consistency and reliability.

Data were collected over a period of six months. The schedule consisted of the first stage of recruitment and screening, the delivery of the baseline test and baseline questionnaires, the delivery of the six bridging tDCS treatments over a six-week period, the delivery of the last post-intervention test and second set of questionnaires, and, finally, follow-up interviews with parents seeking feedback on their perceptions of the families' experience throughout the course of the whole intervention procedure.

In conclusion, the strategies used for data collection within the framework of this study were deliberately chosen, considering each type of information being assessed, in order to collect rich and robust data about the effect of tDCS on social communication in children with ASD. By using questionnaires, standardised parental evaluations and semi-structured interviews, the researchers employed multiple strands of knowledge-collection, ensuring a complete under-standing about the question being addressed by the research study.

34.3 Summary of Data Collection Methods

The aim of the study is to take information from the families of children who have been diagnosed with Autism Spectrum Disorder (ASD) and have undergone a treatment with transcranial Direct Current Stimulation (tDCS). In order to get the information, a questionnaire is going to be sent thorough different internet sites and by being face to face with the families in the rehabilitation centres where the children have been treated.

Online distribution platforms such as Qualtrics and SurveyMonkey will be used because of their excellent features in supporting advanced survey designs, and their high standards of data security. The benefit of an online survey is that it's more convenient to access and time-saving; it also has the ability to transcend geographical boundaries. Participants can take the survey at their own speed, at their own convenience, which makes it less burdensome and increases their likelihood of participating.

Additionally, after online distribution, the surveys will also be conducted in person at certain centres for rehabilitation. This will mostly help with families who have limited internet access or who prefer speaking with someone in person. Researchers will be present at these centres to guide participants who require help with the survey, making sure all questions are accurately answered. This manner of data collection allows more qualitative information to be gathered through chatting and observation.One key step to ensure the ethical conduct of the data collecting process is the informed consent, which necessitates an consent form being provided prior to the start of the survey. The informed consent form will explain the purpose of the study, any procedures to be taken, the potential risk and reward as well as the duties and rights of participants. Provided in digital form for the participants who have online access, or in hardcopy form

for in person-participants, participants will be able to provide the confirmation of understanding and agreeing to participation by signing the digital document, or physically. The permission form will emphasise the voluntary aspect of the study, by indicating the participants' free will to opt in and out during or before the survey. In order to achieve the intended clarity, the permission form will be written in plain language with no jargons used. As long as there is a necessity for the centre to reach out to those who may not have the native language (English), multiple languages may be provided in order to increase accessibility.

This data will have to be correct and complete if the results of the study are to be credible and reliable. There are several protocols to be uprighted in order to achieve this objective. A pre-testing of the survey will be done before a full implementation takes place with a limited number of families in order to detect and correct ambiguities and technological hiccups if found. Clear instructions will be provided in the beginning of the survey, and in every section, to guide participants in correctly answering questions. The electronic survey platform will automatically have a validation mechanism that will not allow for incomplete submission of data, and will prompt participants to provide answers in questions left unanswered. Researchers will constantly monitor the flow of data collection as it unfolds and will immediately prevent new concerns. Families who fail to complete the survey in the allotted period will be followed up on to be reminded to complete.

Information collecting process will be as organized as possible in order to guarantee it is gathered swiftly and efficiently. Timescale will be split into various essential stages: Planning Phase, Initial Deployment, Follow-up and Reminders, Data Cleansing and Analysis, Final Review and Report.

Counted into Planning Stage, lasting one month, all activities how to build a questionnaire, get ethical permissions, and conduct pilot testing for the questionnaire is included. Initial Deployment Stage lasts for two months in order to distribute the survey in both online and offline forms. Follow-up reminder will also be coined within Initial Deployment Stage, covering for any participants who haven't finished this questionnaire. Meanwhile, data cleansing during this stage will also commence to clear irrelevant information during the first stage. The Follow-up and Reminders stage will occupy one month.

During this period of time, all participants who haven't gone through this questionnaire will be followed up by us, and if they're unable to do it online, more in-person sessions will be facilitated for them to complete. Data Cleansing and Analysis will take up two more months. After all the raw data collected, data cleansing process will follow first to ensure the reliable information before further analysis. During Final Review and Reporting Stage, lasting one month, we have to check whether the findings drawn after data analysis is reliable or not, and then group the findings to be prepared for our presentation. We will also communicate the findings to the stakeholders.

The survey is designed to gather a full understanding of the family experiences regarding tDCS treatment for children who are diagnosed with ASD. Survey questions will be divided into a number of sections related to different aspects of the treatment and its effect.The beginning will record basic demographic information about the child and family such as their age, sex, ethnic background, socioeconomic status and education. Understanding the demographic context is important because it allows one to interpret the data and make sense of any trends or correlations.

The second part will provide full description of ASD diagnosis given to the child, including age at diagnosis, level of symptoms severity and presence of other comobid conditions, as well as previous interventions and treatments administered to the child, which are highly important to be reported as treatment history and baseline condition before tDCS intervention.

The final section will indicate the precise details of the tDCS treatment through a description of the duration and frequency of sessions, the specific brain regions being targeted, and any suspected or observed side effects. Participants will be asked to provide some feedback, rating how satisfied they are with the treatment procedure and the management of that treatment. In the fourth section, parents will be asked to rate the treatment with tDCS on a variety of scales. Questions will concern the child's social communication abilities, the extent to which the child adapters behaviors, and whether development has improved as a result of the intervention. The goal of this phase is to acquire both quantitative (ie, rating scales) and qualitative (ie, open-questions) information in order to have a more holistic view of the outcome of a treatment.

Finally, the fifth section pertains to the family experience with tDCS (what they might have expected or been told, and their experience during and after the treatment). Their questionnaire will ask them how the tDCS treatment impacted their emotions and psychological comfort level; how it altered their relationship with their loved one experiencing the treatment; and whether they felt satisfied with their experience as a whole. Armed with their personal experiences and insights, they will answer these questions and share their perspectives. Their insights and stories will offer researchers invaluable qualitative information that will complement the clinically based quantitative research findings.

Understanding the logistical challenges and barriers faced by families when engaging in tDCS administering can enhance our appreciation for the feasibility of this intervention and its uptake potential. Question 6 will ask about logistical obstacles and financial barriers, in addition to any other barriers that may have stopped them from engaging with the treatment intervention fully. Understanding these barriers thoroughly may inform intervention development efforts and policy prescriptions. In the last portion of the session, participants will be asked to comment on current tDCS therapies for children with ASD and share ideas on how we can improve treatment by: a) strengthening family caregiver support; b) rectifying flaws or inadequacies in the method; and c) better meeting concerns and needs of children and their families in future interventions.

The first portion of the survey will collect demographic information, so that the team knows something about the background of children with a diagnosis of ASD and their families. It will include questions about the age, sex, ethnicity, socioeconomic status and educational history of the child. Why is this important? Although it can be tempting to focus on the quantified aspects of daily life, such as numbers of steps taken, calories burned, children spoken to, the value of age-old judgements will most likely hold true.

Collecting demographic data is invaluable because it can be analysed in conjunction with the rest of the data and a correlation examined – such as whether the quantified aspects of daily life (just to name one possible direction) show a uniform increase in some demographic groups or not, or whether any of the demographic parameters is significantly predictive of the outcomes. An example of such an outcome would be the subjects' evaluations on the efficacy of tDCS. Is its efficacy affected by certain demographic parameters; could this help in honing future therapies with better accuracy? Are particular groups scantily represented in the study,

which might be a call for experienced efforts to expand the recruitment? Demographic data helps one determine the deliberateness of one's analysis.

Experience with Treatment

In the second component of the interview, we will delve deeper into the treatment experience of tDCS. The questions will focus on how long and often the participants received tDCS; in addition, we will ask them which brain regions were targeted, and if they experienced any unwanted side-effects during the treatment. Individuals will also be queried about the practical aspects of attending the treatment sessions such as the amount of time it took them to get to the session, what types of expenses were associated with attending the treatment, and any problems they might have had in maintaining the treatment schedule. Nuanced information about these issues is essential for the evaluation of tDCS as a potential treatment for ASD. Logistical difficulties can detract from an individual's satisfaction with a treatment irrespective of its therapeutic efficacy.

Perceived effectiveness

A final section will assess the perceived effectiveness of the tDCS treatment in the eyes of the family. This section will include both quantitative questions, such as for the parents to complete rating scales to reflect any improvements in social communication and behavioural changes, and qualitative questions in which the families themselves can detail what they feel has been altered in specific ways. This section will provide an overall assessment for the effectiveness of this treatment by collecting two types of data. For example, a numbered rating shows how the family feels their child has improved, but the qualitative response offers explanation and specificity of how the improvements manifest. The dual methodology creates a richer data set, making it more substantial and rich.

Alterations in behaviour

The fourth section will address the behavioural changes that were noted in the children after tDCS. These questions ask about changes in social behaviours, communication, and repetitive behaviours; changes in any of these domains that had good or poor replicability; and any other new behaviours or skills learned by your kid since starting this programme. This section will also ask how long these changes remained after the original treatment. This question will determine whether changes last, or whether they naturally flatten out after the programme is over. Understanding these mechanisms is important for understanding how long a positive change from tDCS will last, and also whether it is a deliverable option for a sustainable solution for ASD.

User-Friendliness

The fifth segment will consist of an assessment of the usability of the tDCS treatment from the point of view of the family (how clear was the treatment regimen, was it easy to follow, was the therapy practical and comfortable, were the technology challenges overcome etc?). People will reflect on their experience, comment on challenges and offer suggestions for changes in the treatment procedure. This may seem very micro, but that's the point: we readily adjust complex tools like cellphones to be better optimised for our day-to-day use, and tDCS is just as complex, with just as much promise, potential and future.

General contentment

The sixth section will ask the families about their overall satisfaction with the tDCS treatment. It will include questions about the general impression of the treatment procedure, the way in which the health professionals assisted with the treatment, the perceived helpfulness of the

treatment, in comparison with other treatments for ASD the family has tried before, their willingness to recommend this treatment to other families who are facing the challenges of ASD, and their likelihood of continuing this treatment if given the opportunity. Assessment of overall satisfaction with the treatment will provide a broad overview of the acceptability of the intervention, and offer valuable insight into the users' view on the treatment's successes and the areas in need of improvement.

Unrestricted Questions

The last section of the questionnaire will include open-ended questions to give families the space and opportunity to further explain their experiences. These questions will ask participants to share any additional benefits or barriers they have found, their suggestions for making the treatment better, or any other comments they would like to share. Open-ended questions are crucial for gathering complex information and narratives that can be missed by structured questions. The qualitative data will provide detailed information that can contextualise the quantitative findings, and provide a richer information about the family's experience with the tDCS treatment.

Methods for Analysing Data

We will execute an extensive data analysis plan to make sure the study results are robust and accurate. This plan will involve many different steps, each aimed at a systematic review and evaluation of the data.

Statistical analysis of numerical data

The quantitative results of the survey will be analysed using statistical software such as SPSS or R. Descriptive statistics will be used to summarise the information pertaining to demographics, treatment history, perceived efficacy, behavioural change, ease of use and overall satisfaction. Measures such as mean, median, standard deviations and frequency distributions

will help us get an initial look at the data. Statistical inference will be employed to explore the relationships between variables and assess the importance of the results. For example, t-tests or ANOVAs can establish whether tDCS is effective across different demographic populations, while regression techniques can help identify potential predictors of treatment efficacy, and factor analysis can also be employed to identify the latent structure of treatment satisfaction or perceived effectiveness.

Analysis of qualitative data

The qualitative data will be analysed through thematic analysis which aims at identifying, exploring and documenting patterns (themes) within the collected data. NVivo, or any other software for conducting qualitative analyses, can organise and facilitate this through coding.

It will start with the phase of familiarisation, where the researcher will read the qualitative answers several times to get an initial understanding. Then, the coding process will begin, where the interesting phrases will be identified and assigned codes that represent more general themes. After this, the codes will be grouped into themes that accurately reflect the core of the answers. Finally, the themes will be reviewed and improved to ensure they are correctly determined by the evidence and coherent.

Integration of mixed methods

Because of using the mixed-methods approach to gather ideas for this study, it is critical to combine the quantitative and qualitative data in order to understand the outcome. The merging will be accomplished by using the triangulation method, that is, the evaluation of and comparing of data with multiple sources in order to discover patters that are consistent or not.For instance, qualitative observations about perceived effectiveness of tDCS treatment might be compared with quantitative scores. The disparities between these two kinds of data would be investigated to understand the reasons for the discrepancies. In this way, the findings could be

substantiated and participants' experiences of using tDCS treatment would be thoroughly understood.

Methods for Guaranteeing the Accuracy and Wholeness of Data

Achievement of accuracy and completeness are the primary objectives of this research and the approach proposed to implementing an electronic personal questionnaire will realise these objectives. Survey phase - specific information about the methodology involved in the online survey (the questionnaire itself) follows: Multiple protocols will be in place to assure accuracy and completeness of data including conducting a pre-test of the survey with a small sample of families to identify and correct ambiguities and technical faults before fully implementing the survey. Clear instructions will be given at the beginning of the survey and also at the beginning of each section of the enquiry to help respondents accurately answer the question.

An automated validation step will be embedded in the online-survey programme such that the computer will insist that the participant return and fill out any uncompleted questions. Researchers will consistently oversee in real time the progress of the data being collected, addressing any new issues as they arise. Participants who have not completed the survey within a specified period will be contacted so that they can finish.

Data Collection Timeline and Schedule

This process will be conducted in an orderly manner for data submission and collecting. The whole timetable will be divided in several key phases: Preparation Phase, Initial Deployment, Follow-up and Reminders, Data Cleaning and Analysis, Final Review and Reporting.

During The Preparation Phase which will last one month, The survey instrument will be created, ethical clearances will be obtained, and the initial test of the survey will be done. During The Initial Deployment phase which will last two months, the survey will be sent out online and also in person, receiving monitoring and initial data cleansing at the same time.The Follow-up and Reminders phase which will also last one month will send follow-up emails to those who did not complete the survey and arrange for further in-person sessions if necessary After The Data Cleaning and Analysis phase which will last two months, recieve the initial data cleaning process and initial analysis.Finally The Final Review and Reporting phase, which will last one month will examine the data analysis results, prepare the data for presentation for those in positions of leadership, and communicate the results to stakeholders.

35.1 Data Cleaning and Preparation

Before even starting to analyse the data, it is necessary to carry data cleaning and prepration to ensure that the data is sound and trustworthy. Data cleaning is a step of looking for errors and inconsistencies in the data, or dealing with missing information as well as ensuring the validity, correctness and completeness of the data. To avoid any bias and inaccuracy in the future whenever we use the data for study this is important.

Error Identification and Correction

The first stage of data cleansing aims at a manual check of the dataset for major visual anomalies and discrepancies such as outliers for data entry errors, duplicate entries, inconsistencies in data recording methods, and other types of error. The next step in the process is to standardise

any inconsistencies in the recording of the demographics such as age or gender in the dataset to ensure its homogeneity.

Dealing with Missing Values

Missing values are a common issue with survey data, and the study can end up looking very different depending on how it's handled. Three common approaches for handling missing data include this is the tedious option of removing records that are missing data values. If a substantial proportion of your data are incomplete, this could be quite a drastic loss of information.

The value of Imputation denotes replacing missing values in a data set with approximated data derived from other available information. Examples of imputation methods include mean substitution, regression imputation and algorithms such as k-nearest neighbours.Fitting models using statistical approaches that deal with missing data, such as multiple imputation or maximum likelihood estimation.

Ensuring Precision and Thoroughness

Accuracy checks check to see that the information is consistent with predefined standards or criteria; hence, this involves verifying the age values to make sure that they fall within a contextually appropriate range for the study population or checking that all needed fields have been filled out in a way consistent with the survey specifications. Completeness checks check that all relevant data elements have been recorded such that essential information has not been neglected.

35.2 Techniques for Quantitative Analysis

A descriptive statistic refers to any sort of statistic that is used to describe data or summarise observed variations among sample points, such as measures of central tendency (eg, mean, median, mode) and measures of dispersion (such as standard deviation and range).Descriptive statistics summarise the basic properties of a dataset. It provides a simple view of what the sample and measures are like. They incorporate measures of central tendency (eg, mean, median, mode), variability (eg, standard deviation, variance, range) and frequency distributions. This is about understanding the overall properties or trends of the data.For example, the mean of age tells us the average age of the people studied, and the standard deviation describes how spread out the ages are in the sample being studied. A frequency distribution tells us the number of people in various groups, such as reporting different levels of effectiveness of the tDCS treatment.

Analysis of cross-tabulations and correlation

To investigate this relationship, we perform special ‘cross-tabulations’, sometimes referred to as contingency tables. These help us understand how different categorical variables – such as markers of demographic information, or levels of depression or anxiety – interact with and depend on other variables. For example, a cross-tabulation could reveal how likely men are to perceive tDCS as effective versus how likely women are to describe tDCS as effective.

Correlation analysis measures the strength and direction of the relation between two quantitative variables. The process is typically used to assess Pearson’s correlation coefficient. Correlation analysis may be used to explore the relation between duration of tDCS treatment and changes in social communication ability, for example.

35.3 Techniques for Qualitative Analysis

Thematic analysis

Thematic analysis is a rigorous process for systematically identifying, analysing and reporting themes across qualitative data. It relies on the following stages.1. Repeated examine of the qualitative answers to grasp the content.2. Systematically grouping significant characteristics of the data over the entire dataset. 3. Are the codes tentatively classified into possible themes? Are you sure all the data relating to each possible theme have been collected? If not, find them.4.Developing themes: Mapping out the themes that have been identified. Consider how they interconnect, overlap, connect and follow on from each other. Ensure all quotes and evidence relating to each theme are relevant.

Explore these themes in more depth via external websites. Here is a coded extract:‘I've always had a strong belief in my individuality and personal identity. Although I was not brought up in a super-religious environment, I have always had a clear sense of the symbols that identify me as a Jew. We celebrated Jewish festivals and commemorated the Holocaust every year. My grandparents, who were survivors, as well as my two great-uncles, all came to Britain from Poland. So, the identity of being a Jew was always "front and center" in our family.'Jewishness influenced every aspect of martinfilms life: ‘As a Jew, the world you grow up in is vast, but your reference points, especially if you are of Eastern European descent, are all Jewish.’Martinfilms describes the experience of being raised as a Jew in 1970s suburbia as relatively isolated compared with today. ‘It wasn't alienating, although it was isolating,' he recalls. This was mainly because there weren't that many Jews living in his neighborhood.'I was the only Jewish kid in my infant school, and there were only two Jewish children in my class in primary school – a brother and sister who become my best friends. They were of an Eastern Jewish background like myself.'Martinfilms felt quite alone for many years.

Five.Explaining and Labeling Themes: Continued experimentation to deepen the specific details of each topic and the overarching story the work conveys.

Programming and Designing Themes

Coding involves the labelling, or tagging, of parts of text with key terms that denote a specific theme or pattern. Codes can be both descriptive and interpretative depending on the depth of the analysis. Usually, the first step of the coding process involves simply reading the data – which, depending on the size and nature of the text, can be quite an intensive process in itself. After a first round of coding, the codes are grouped together around key themes that encompass key data elements.For example, answers about the perceived benefits of tDCS could be clustered into themes such as ‘improved speech’, ‘behavioural change’ and ‘ease of use’. These themes could then be explored in depth to better understand shared experiences with, and variation in, use of the treatment for families.The combination of quantitative and qualitative data is known as the fusion of quantitative and qualitative data.

The use of quantitative and qualitative data provides a more comprehensive and thorough understanding of the research outcomes when they are combined. With the addition of another source of evidence (qualitative), it is possible to triangulate data by cross-checking results from one method with another.For example, we could improve quantitative statistics that reveal improvements in social communication scores with qualitative reports on how these changes manifest in everyday life. There are many ways to achieve this combination. The convergence model compares and contrasts the quantitative and qualitative data, focusing on places of agreement and disagreement.

dialectical number Explanatory Model: Qualitative data incorporated to explain and contextualise quantitative findings.The answer is Exploratory Model: Harnessing qualitative findings to inform the development of quantitative measures or to investigate new hypotheses.

35.4 Software and Tools for Data Analysis

Different software programmes can help you work with both quantitative and qualitative data. A programme such as SPSS, R or SAS (packages used for statistical software) give you the knowledge of advanced statistical methodologies for quantitative data. They have the ability to handle and analyse large numbers of data points, conduct complex statistical testing procedures, and produce detailed reports and visualisations.Qualitative analysis: NVivo and Atlas.ti are manufacturers of qualitative analysis software – basically, life became much easier. The options for coding, theme-building and even for the visualisation of qualitative data provided by NVivo and Atlas.ti are almost infinite. What's more, the programmes allow for combining quantitative with qualitative data; researchers can connect quantitative results with qualitative observations.

36. .Ethical considerations

During research on humans, such as children with ASD, ethical issues are of supreme importance, as it is essential to guarantee the physical and emotional safety of participants besides respecting and adhering to principles and rules. Addresses the safety of participants and the requirement for informed consent.

Informed consent requires providing prospective participants with detailed information about the study, including its aims, methods, risks and benefits, in a manner that is comprehensible. Prospective participants (or their legal guardians) must give consent to be included, and must do so on the understanding that they can withdraw at any time without detriment. Ensuring the safety of participants involves taking steps to minimise any risks associated with the research – this includes closely monitoring for adverse effects of the tDCS treatment and having procedures in place to correct them if they do happen. Keeping open the privacy and confidentiality around participant data involves data storage in safe ways, and anonymising data where possible.

What we talk about there is how future research might retrieve ethical standards to keep in mind. Second, ongoing future studies must maintain high ethical standards by refining informed consent practices to ensure consistent and optimal levels; developing participant safety measures; and maintaining transparency in study results dissemination. Third, it is essential to integrate ethical considerations by actively including participants and their families from the onset throughout the research and encouraging and addressing their perspectives and concerns.

Obtaining Consent and Maintaining Privacy

Two of the primary normative requirements of research are gaining informed permission and protecting anonymity, ensuring that research participants know precisely what the study is about, the risks and benefits, and their rights. Gaining informed consent means that participants are given detailed information about the study that is clear and easy to understand. This includes information about the study's purpose, the methods, the risks, the benefits, and the level of privacy protection in the research. Participants should be given sufficient time to consider their

participation and give their consent freely, without coercion or deception. For vulnerable populations, such as childhood-onset ASD populations, conditions that impact cognitive faculties, assent should also be given by a legal guardian or parent, but wherever possible, seeking their assent should also be pursued.

As implied, confidentiality does guarantee that participants' information can't be revealed to anyone without their permission. All efforts should be made to guarantee confidence and the privacy of participants. Research data that can identify participants under an assumption must be anonymised. Data, therefore, should be kept secure in appropriate data storage systems, access to which can be limited to certain number of authorised personnel. Participants' identifiable information should also be replaced with pseudonyms or case-number designation, to ensure that participants won't be identified from the research records easily. Confidentiality is especially important in ASD research since the data may contain sensitive information about the health and family dynamics. One of the incentives behind conducting such research is to alleviate stigmatisation of the participants and their families, to safeguard their dignity.

Protection and retention of data

Data security refers to the policies about securing data from unauthorised access, breaches or other types of misuse. When doing research with human subjects, we must develop strong data security practices. This involves using encrypted storage systems, highly secure servers, or password-protected files. One can also conduct the periodical audit and update of the security systems to uncover potential vulnerabilities. To guarantee the physical security of the data, such as the informed consent, for example, questionnaire, we suggest using a safe storage place, typically a locked cabinet, to prevent an unwanted access.

This means electronic data should be saved on encrypted devices, researchers should use protected mechanisms such as encrypted email and secure file transfer protocols to send data to members of the team, and researchers should develop a data management plan detailing the exact procedures for the collection, storage, access and disposal of data that is compatible with the institution's rules as well as data protection legislation such as the General Data Protection Regulation (GDPR) in the European Union or the Health Insurance Portability and Accountability Act (HIPAA) in the United States. Data disposal methods such as physically shredding papers or securely 'wiping' electronic data will need to be used to ensure the data are unrecoverable once your study is complete.

Evaluation of Risks and Benefits

Given that studies on vulnerable populations involve special protections, it is important that the prospective benefits of the study far outweigh the prospective harms to participants. In tDCS research with children with ASD, the prospective harms might include having an unpleasant or modest adverse experience due to the stimulation, while the prospective benefits might include improvements in social communication or behaviour. Researchers must run safety risk assessments with a view to identifying and evaluating threats to physical, psychological and social safety, as well as implementing techniques to minimise risks. This might be done through the development and use of safety procedures, explicit participant instructions, and research staff who could oversee or manage an adverse event. The benefits of participation for participants and their families (such as potential enhancements in quality of life and contributions to the scientific endeavour) also need to be clearly communicated.

The weighing of risks and benefits must be part of an iterative process as the research progresses, with ongoing monitoring allowing early identification and mitigation of new risks. Researchers need to be transparent with participants about the limitations of the study and the extent to which success can be guaranteed. This allows the research to maintain confidence and for participants to have realistic expectations about the outcome of their contributions.

Approval from the Institutional Review Board (IRB)

Before beginning any experiments with human volunteers, investigators must submit their proposal for review to an Institutional Review Board (IRB). The IRB asks questions about the study design, its methods, risks and benefits, its informed consent procedures and more, to ensure the ethics of the research. During the IRB review process, the research proposal is carefully considered in light of the overall goals, the methods to be employed and the potential outcomes for participants. Researchers must provide detailed paperwork, such as informed consent forms, procedures for handling private data, and risk mitigation strategies. The IRB ensures that the research project is conducted within the general principles of the good life, including respect for persons, beneficence and justice – as these concepts are specified in documents such as the Belmont Report.

The researchers are obligated to conduct the study as was approved by the IRB within the confines of the protocol approved, and any substantial modification of study design or protocols during a study must be submitted for review and ap-proval. It is important for investigators to communicate with the IRB on an ongoing basis during the course of the study, whether that be through a status review of the study showing timelines of completion, any adverse events discovered, or re-approval for continuance of the study.

Ethical principles and adherence to regulations

Taking ethically responsible actions and adhering to the relevant rules are core aspects of doing research. As is typical for standards in our society, ethical standards define a systematic route to doing research responsibly and, most importantly, with respect to the well-being of participants and the integrity of the research procedure. The criteria refer to the values of autonomy, non-maleficence, beneficence and justice.Researchers need to know what the ethical rules are for their discipline and follow them during the entire course of a study. At the outset, this includes gaining informed consent, maintaining confidentiality, minimising harm, and making the benefits of a research project outweigh the risks. Following institutional policies as well as national regulations for conducting research with human participants is likewise essential to ethical conduct.All members of a research team should be subject to similar training and education in ethical research methods, so that their remained vigilant and capable of maintaining elevated ethical norms, and whenever the conduct of the research raises ethical issues, that these are dealt with appropriately. Researchers should also work cooperatively with their ethical review boards or other oversight agencies to ensure ongoing monitoring and improvement of their methods.

Reliability and validity

Reliability and validity are closely related, and it is vital that both are maintained throughout to create reliable and generalisable conclusions. Reliability refers to consistency in the study findings, where the results should be reproducible under similar conditions. Validity relates to accuracy in the findings, where research must carefully measure what it is intending to measure.

Standard-ised study protocols and methods were necessary to increase the validity of findings; these include the use of self-administered, previously validated questionnaires; standardised methods for collecting data; and training of data collectors to achieve standardisation. Pilot studies can help to identify and correct methodological discrepancies. Increasing validity demands that study design and procedures are appropriate to the research questions being posed: sampling techniques must be relevant, confounding variables minimised and measurement tools accurate. Triangulation – using more than one type of data source or methodology – can strengthen validity through providing a more informed understanding of the research topic.

Reviewing and reflecting on the research process periodically can help to spot any threats to the reliability and validity of the project. Peer reviews or discussions with experts can give significant feedback and can help to improve the methodology of the study. Making concerted and persistent efforts to maintain high levels of reliability and validity is important to creating strong and reliable findings.

Methods for Guaranteeing Dependability

Dependability refers to how consistent a measure is, or how much it gives the same results when applied over and over again. Ensuring dependability relies on many different techniques at various steps in the study process.

This is why standardisation of the data-gathering techniques is one of the key strategies to maintain the reliability. This means that a researcher has to use the same protocol for each subject. This guarantees that each individual is placed under the same conditions and asked the same questions in the same way. For the study comparing the effects of applying tDCS on individuals with ASD, researchers have to give the stimulation with the same settings and duration to each subject. Finally, conducting surveys or assessments under controlled conditions

can also reduce the influence of confounding variables that might have caused a difference in the results. Yet another fundamental way is to make sure all data-collectors receive thorough training. Training ensures that all workers involved in the data-collection process understand the protocols and are able to apply them consistently. This includes training in how to administer questionnaires, how to record the answers properly, and how to deal with unforeseen eventualities or questions from respondents. Refresher training sessions can help maintain high standards of data collection over the course of the study.

Measurement devices must be reliable. Study designers must use methods and scales that have been field tested and validated, shown to be consistent in their results. For example, pilot testing is often done to see if a new device will be a dependable instrument for collecting data before it is employed in the main research. Statistical techniques, like Cronbach's alpha, can be used to evaluate the internal consistency of a scale – that the items within the test are measuring the same thing.

Another promising approach involves repeated measures or test-retest reliability assessments. In these approaches, the exact same assessment was performed on the exact same group of participants at intervals to determine whether the results would remain consistent. For instance, repeated social communication assessments on children with ASD in the presence of improvements post-tDCS therapy would serve to demonstrate consistency of the results.

Methods for Guaranteeing Accuracy

Validity refers to the extent to which a research study actually measures what it claims to measure. Many strategies for validity go after different elements of a research design and procedure.This is known as having content validity. In other words, do the measures fully capture the necessary dimensions of the domain of interest? This can be accomplished through a thorough review of the literature and consultation with subject matter experts who can verify that all the necessary dimensions are incorporated into the tools. The measures used to evaluate the effects of tDCS for the social communication of ASD children should cover numerous facets of social responsiveness and communication abilities, as well as behavioural changes as caused by ESDM.

Another overlooked important characteristic is construct validity: measuring what you intend to measure. To do this, researchers can use a method such as factor analysis to provide evidence that there is a unique structure to the instrument and that the items are grouped into the correct theoretically defined constructs you are trying to assess. Construct validity can also be addressed by demonstrating a correlation between the measures and other instruments that are recognised to measure related constructs.

Criterion validity involves comparison against external criteria or standards. An aspect of this is concurrent validity – that is, comparison with an existing measure at the same time of the relevant event. Another manner is predictive validity, which involves having the measure predict a future outcome. For instance, if a new social communication assessment is used in your research, you can check your results against what you'd find using an existing social skills inventory.

37. Conducting pilot testing and making necessary revisions

Pilot testing is a necessary step of the research process that helps in determining possible problems and improving the study design before the primary collecting of data occurs. Pilot studies allow researchers to test their instruments, processes and protocols in a smaller and controlled environment, shedding light on problems they might not have foreseen beforehand, and to enhance the overall design of the research.Through the pilot testing of a given study, researchers test the feasibility of the study design, including the recruitment strategy, data collection approaches and intervention protocols. For example, prior to starting a formal investigation on the potential effect of transcranial direct current stimulation [tDCS] in Autism Spectrum Disorder [ASD], a pilot test could help with establishing the feasibility and tolerability of the stimulation protocols among participan-ts, as well as with the efficacy of the evaluation tools in measuring the effects of the intervention.

The feedback from this pilot testing phase is crucial in making the necessary tweaks. The scientists have the chance to modify the equipment to increase the clarity, tweak the data collection protocols to enhance efficiency, and plan the intervention protocols to confirm that the participants are comfortable and compliant. Such an iterative process ensures that the study design is robust and capable of collecting reliable and accurate data. This position is worked out by measuring the angles between the object and two known points – a process known as triangulation.

Triangulation involves the use of multiple data sources, methodologies or perspectives, in order to improve the accuracy and reliability of research findings. Researchers may increase their understanding of a problem under study by applying different methods and comparing their findings.

Triangulation comes in different forms. Methodological triangulation involves the use of more than one type of data-collection method, such as combining quantitative surveys with qualitative interviews. For example, with tDCS, we wanted to measure its impact on ASD, using both standardised evaluations and in-depth interviews with parents to contribute depth of data regarding the outcomes of the intervention. Data triangulation can help to determine patterns and verify findings in different contexts, by collecting data from multiple sources or multiple time points. For example, researchers could collect data from different rehabilitation centres, or follow up with participants at different time points after their initial assessment or treatment.

What is imagined is a sort of investor triangulation, in which data are analysed by multiple researchers independently, and such triangulation can reduce bias and raise the credibility of findings. Multiple perspectives and expertise, if applied to evaluating and interpreting the data, can increase its benefits to the study.

37.1 Constraints and Difficulties

No matter how thoughtful they are about rational measures that promise secure reliability, and quality control, every re-search is limited by constraints and problems. Those constraints have to be dealt with for the sake of the credibility of the process itself. The first fundamental constraint is the threat of bias. Reducing bias – the inadvertent impact of the investigator's own beliefs and expectations on the results of a study – requires rigour in the proper planning of a study design, collection of data and analysis: from conducting a thoughtful assessment of one's own beliefs and biases to putting in place approaches to maximise the objectivity of the processes of the study. For example, blinded assessment procedures (that is, evaluations that are

done without the assessors being aware of the group members to which they are assigned) can be an effective way to minimise bias.

A second is the transferability of the findings to other contexts. Studies tend to focus specifically on a certain kind of group (eg, pensioners in Durban) and place (eg, Durban), distancing the findings from other contexts. Researchers must be rigorous about analysing and reporting the characteristics of the sample and the study context, and about the limits of generalisability of the results. They therefore need to be very clear about the scope and limits of any practical recommendations they might propose for transfer to other contexts. They should explain the limits in any reports and recommend how the findings might be extended through further research in future studies.

Besides academic criteria, pragmatic constraints might also affect the design of the research process. For instance, insufficient budget, time and staff might add a significant limitation to the conduct of the research; researchers need to show their ingenuity in using the available resources and finding smart solutions to overcome the barriers. Given the struggles faced by vulnerable groups, one must be especially careful in selecting and assessing research that involves such populations: attention to ethical issues must be clear, reflecting ethical norms, and applied in practice through informed consent, confidentiality and minimisation of risk to participants. Considerations must be carefully and openly articulated; collaboration with an ethics review board is crucial, and there must be regard for institutional and regulatory regimes.

Possible constraints of the research

Studying the use of transcranial Direct Current Stimulation (tDCS), more specifically the Dorsolat-eral Prefrontal Cortex (DLPFC), on paediatric patients diagnosed with Autism Spectrum Disor-der (ASD) is a daunting task with many possible constraints. These constraints might 'bias' study outcomes and effect results on internal validity, reliability and generalisability, while a responsible practice of science must account for such constraints. One particular limitation is that you could have a small sample size. Being able to find enough subjects who fall exactly into a tDCS study's criteria can be quite challenging. This is especially true in certain populations (for example, children with Autism Spectrum Disorder [ASD]). Low power due to small sample sizes can make it more difficult to spot significant effects, or to generalise your findings from your participants to the population at large.

The third limitation relates to between-participant variability in tDCS reactivity: children diagnosed with ASD come with a wide range of symptoms and severity in each individual case, and can react differently to any given therapy. When there is a high degree of variability, it can make data difficult to analyse and interpret, potentially masking whether the observed effects were due to the tDCS intervention or some other participant-related characteristic.A further limitation is the likelihood that the outcome will be influenced by placebo effects, or the expectations of those participating in the study. In order to attempt to control for this, when you set up a study of neuromodulation (in this case, tDCS), the individuals participating and their families might have heightened expectations of improvement, and these can influence how they perceive the effectiveness of the intervention, generating a placebo effect that might confuse the results, making it difficult to know if and to what extent tDCS is having a real effect.

37.2 Difficulties in Data Collection and Analysis

If you want to study child-ren with ASD, collecting data could also be quite the challenge, due to several factors. First of all, in order to sustain participant engagement and participation in your experiment, you could run into some challenges, especially since children with ASD can have various tolerance levels for the duration of the tDCS sessions and also the tests involved so you might not get enough data in the end.

Also, in research with this device, administering the intervention reliably to all participants is very involved: the study team needs to be trained to deliver it carefully and by the book, and also to ensure that its administration is consistent from one participant to another. Even small deviations from the book can add a confounding element to a study's results. Ensuring that tDCS is administered uniformly, across all participants, is vital for getting replicable results. Mixing quantitative and qualitative data in data-analysis is challenging. Quantitative data reflecting ability, such as standardised measures of social communication abilities, must be recorded with precision and analysed using appropriate statistical methods. Qualitative data from sources such as family input and observations must be coded and thematically analysed in order to produce meaningful findings. Combining these data types in a logical manner can be complex and time-consuming.

37.3 Approaches to Address Limitations and Challenges

If you're bound to a small sample size, you can bring others into the fold by recruiting from multiple sites – essentially partnering with other institutions to sign up more people. This has the dual benefit of not only increasing the number of participants, but also makes it easier to

generalise the results to multiple contexts. With stratified randomisation one can try to level out the impact of individual differences in response to the tDCS, and so make sure people are equally distributed across the various strata of severity and demographics. A full-scale baseline assessment should also help with picking up possible moderators that mask positive effects, or with effect-correcting adjustment of the statistical analyses.

One way of dealing with placebo effects is to have and use a good control. This could be a sham tDCS condition, where we subject participants to placebo stimulation that mimics the real montage and costumes the participant the same way, but they receive a placebo (figure shown above). Blinding the study could be used to mitigate some expectancy effects, which involves having neither the participants, nor families and not the researchers or test administrators know who gets what. This would further improve internal validity. This can be achieved by using engaging, child-friendly assessment measures, and flexible scheduling of sessions to fit participants and their families. Providing feedback to families on the progress of the study can make them feel part of the endeavour.

This thoroughness can be achieved by employing mixed-methods approaches to the analysis of the data that are collected. Advanced statistical techniques for calculations based on quantitative data and strict coding protocols for qualitative data can be employed with assurity that all of the data relating to the intervention are analysed in the level of detail and rigor required. Then, the use of triangulation can substantially increase the validity and richness of the results.

Anticipated Results

While this is unreliable in some respects, any findings are likely to be important, as the study will shed light on tDCS effects on the social communication of children with ASD. Quantitative outcomes could include improvements on certain standardised assessments of social communication abilities, while qualitative data could give vital information from families regarding the potential of the intervention in the real world. When these elements are brought together, they should give a rounded picture of the potential benefits and harms of tDCS as a treatment for ASD, which could help steer future trials and clinical work.

Improved comprehension of familial encounters

A valuable part of this study is a greater understanding of what families think tDCS represents as a treatment for children with ASD. Family input should always be obligatory in the treatment and maintenance of children with autism: their opinion on new therapies can help quantify whether a new treatment is practical and valid. This study employs a variety of tools, from standardised surveys to qualitative interviews, and incorporates many strands of perspective from family members on the acceptability, challenges and overall satisfaction with tDCS as a treatment method.

Parents expressed a range of qualitative enhancements in their children's social communication abilities, with achievements balanced by aspects that required extra attention. This qualitative data demonstrates the richness of perspective from non-clinical settings – including how people perceive tDCS interventions in terms of the ability and effort required to use them, the observed changes in behaviour, and the emotional and psychological changes experienced by the child and their family members. Importantly, the act of gathering family input in assessing and designing therapeutic interventions advances treatment development beyond the clinical

understanding of effectiveness, a necessary step for making treatments feasible and persistent when placed in the complex contexts of real life (Gibbs et al., 2019).

Understanding the effectiveness of treatment

It is a well-designed study showing significant improvements in social communication skills in children with ASD, measured with quantitative metrics. Those positive effects following non-invasive tDCS occurred in participants with increased visual contact, higher participation in social exchanges and improved oral communication. tDCS does improve social abilities (Fregni et al., 2021).

Further, the combination of quantitative and qualitative data allows us to make a broader evaluation of tDCS success. Quantitative data provides evidence of behavioural improvements, while qualitative data from families provide a broader context in terms of understanding the manifestations of these improvements in their everyday interactions and overall quality of life. It is through this dual approach that we can increase the reliability of our findings and gain a broader understanding of the potential benefits that tDCS might have. This supports the continued investigation of tDCS as a potential therapeutic option for ASD (Eack et al, 2018).

37.4 Practical Implications for Clinical Practice

What are the clinical implications of this study? On the one hand, we've demonstrated tangible improvements in social communication that support the use of tDCS along current treatment strategies for treating children with ASD, potentially alongside other behavioural and educational therapies. Health professionals can rely on these findings to fine-tune these intervention strategies (both neurological and psychological) related to social communication in children with ASD.

Additionally, feedback on tDCS application from families provides an opportunity to improve the treatment regimen. The description of user-friendly tDCS was highlighted as a critical consideration, as was the need for family members to be trained in administering the therapy at home. Paying attention to these important considerations helps us improve adherence and the salutary effects of tDCS, contributing to a more viable and effective option for more families (Donaldson et al., 2020). Comprehensive Examination of the Effects of tDCS on Autism Spectrum Disorder (ASD) for a richer understanding of tDCS in ASD from the perspective of both clinical outcomes and familial experiences. Such an expansive approach can greatly assist in appreciating both the heterogeneity of ASD, as well as the diverse myriad of ways in which an intervention might ultimately shape different members of a family.

The paper emphasises the importance of using multidisciplinary approaches that combine neurological, psychological and social elements in treatment, enabling us to ensure that a therapy such as tDCS is evaluated not just for its clinical efficacy, but also its broader impact on the wellbeing of children with ASD and their families. The study plays a role in fostering more coherent and holistic therapeutic interventions that meet the multifaceted needs of people with ASD (Minson et al., 2017). This study contributes a longitudinal and ecological perspective that helps colleagues and families gain a deeper understanding of tDCS as a treatment for ASD, and to understand its usefulness, both in terms of its effectiveness (reduced anxiety and disorganisation), and the practicalities of using it in the real world (eg, treatment feasibility and generalisation, as well as some pitfalls, such as dose escalation and increased OCD symptomatology). This critical appraisal offers a detailed and broad-spectrum assessment of tDCS, illustrating both its potential benefits and the real-world trade-offs. It also underscores the im-

portance of developing a treatment strategy that accounts for the wider effects of an intervention on the individual and their family across the lifespan. ASD treatment strategies should seek to expand the repertoire of functional skills, enhance physical and mental wellbeing and – ultimately – better integrate people with ASD and their families within their communities. More assertive neurodiversity campaigns might consider casting aside the struggle for neuron equivalence and instead pursue a strategy parity perspective for greater impact.

Summary of the Methodological Approach

The methodology behind this study was developed to thoroughly evaluate the combined effects of transcranial Direct Current Stimulation (tDCS) and social communication in children with the neurodevelopmental disorder Autism Spectrum Disorder (ASD), as well as provide insights into the real-world experiences of their families, taking into account their valuations, thoughts and feelings. The researchers employed an experimental hybrid research methodology that incorporated both quantitative and qualitative data collection methods to comprehensively examine the results in terms of clinical outcomes as well as what families thought about them.

38. Quantitative Component: Randomised Controlled Trial (RCT)

The study's quantitative component employed a randomised controlled trial (RCT), generally considered the most efficacious experimental design in clinical research when it comes to evaluating the efficacy of therapies. The primary goal of this intervention was to measure the immediate effects of the tDCS on the social communication skills of children with ASD. Participants were randomly allocated to a tDCS treatment group vs a control group that received a sham intervention, and the participants' social communication abilities were measured before and after the intervention with standardised methods.

39. Qualitative Aspect: Structured Family Survey

The qualitative component consisted of administering a fortified survey to the family members of those children participating in the RCT. This survey was aimed at accessing robust qualitative information related to family experience of the tDCS treatment; perceived benefits, challenges and overall family satisfaction with the programme. It was designed with both closed and open-ended survey questions so as to allow families to articulate their impressions and feelings in their own lingual environment, and operated as a valuable source of context that bolstered the quantitative findings.

40. Methods for Gathering and Analysing Data

A combination of uploading online and engaging in on-site discussions at residential facilities also helped to guarantee wide accessibility and a large sample response. The informed consent process was overseen with precision to ensure participant confidentiality and ethical conduct. The quantitative results were interpreted with descriptive statistics, cross-tabulations and correlation analyses in support of finding important patterns and correlations. Qualitative data were interpreted through thematic analysis – a process of coding and topical development that extracts important meaning.

The significance of the methodology in attaining research objectives

This particular approach was adopted in order to reach the research objectives of the study. This study applied quantitative as well as qualitative methods to assess an intervention for ASD – tDCS – via a robust post-experiment evaluation. Through a randomised controlled trial, tDCS

was found to increase social communication skills. Meanwhile, the structured family survey demonstrated the feasibility and perceived benefits of the treatment.

40.1 Thorough Assessment

This was possible because the use of the mixed-methods approach simultaneously allowed for the assessment of tDCS on both the objective and subjective levels. For true process evaluation to occur, we need to capture the impact of an intervention in its entirety, and for that we must assess outcomes to determine if the intervention is operating as planned and where it can be improved next time.

40.2 Improved Validity and Reliability

The use of both qualitative and quantitative evidence also corrected for biases and helped to make the results more consistent. The quantitative measures served to provide unbiased evidence for the efficacy for the intervention; and most importantly, the qualitative measures provided contextual details and personal accounts of the experiences, which ensured that the findings would reflect both scientific rigour, but also practical usefulness.

40.3 Expert clinical and policy recommendations based on informed knowledge and expertise.

This systematic knowledge is paramount in guiding clinical practise and policymaking – deep understanding of family experiences together with identification and validation of barriers and facilitators of tDCS can be harnessed for designing more efficient and patient-friendly ASD interventions and support services for children and their families.

Thus, the use of this type of study design guaranteed that both depth and breadth of understanding about what and how tDCS changes participants with ASD was documented. This study demonstrates not only the clinical efficacy of tDCS, but also the importance of considering the familial perspective in evaluating novel treatment approaches, through using both qualitative and quantitative data.

CHAPTER 4: FINDINGS

41. Summary of Findings Chapter

The results chapter is a critical section of this dissertation which includes summary of the learnings gained through the administration of the structured survey, standardised assessments and in-depth interviews. The purpose of this chapter is to provide a robust examination of the extent of tDCS effects on the social communication abilities of children with ASD. This chapter contains an organised presentation of the data including response to tDCS, family experience, and overall implications of these findings for the clinical practice and future study.

The first section of the results chapter starts with a description of the demographic characteristics of the participants: age, sex, socioeconomic status, level of education, etc. The demographic description provides the context for understanding the sample, and helps make sense of the results presented from then on. Next is the impairment assessment results, which includes a detailed description of the participants' premorbid social communication abilities and behavioural profiles prior to the tDCS intervention (baseline). Baseline data is fundamental in determining if the changes observed after therapy actually stood out from the range of possible pre-therapy states the same person could have exhibited.

Second, following the initial assessments, the chapter analyses the outcomes of the tests administered after the therapy. It highlights the improvements in social communication abilities and the behavioural outcomes gained following the tDCS sessions. The clinical relevance of the impacts related to social communication skills and behaviour in ASD are further supported by quantitative information drawn from the standardised tests and statistical analyses, and by qualitative information gained through interviews with family members. Using the combined information coming from different types of data, it is possible to understand, in full, the effect induced by the intervention.

41.1 The structure of the Results section

The overall organisation of the results chapter is an efficient, coherent way of presenting the results in an abbreviated format. It is automatically divided into several key areas, explaining the different parts of the questions and hypotheses of the work.The Demographic Characteristics of Participants section is the most detailed summary of who the participants are, including by age category, gender, socioeconomic status, or education. It provides the characteristics needed to contextualise the results of the study and make sure the sample is representative.

Baseline Results: Here are the children's social communication skills and behavioural characteristics before the programme. Using validated measures such as the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS) provides a good benchmark with which to compare the data before and after therapy (Constantino and Gruber, 2005; Lord et al, 2000).

Here present the results of the assessment conducted by the end of the therapy. The main issue is social communication skills and behaviours changes as a result of the tDCS treatment. The quantitative results taken from the post-the treatment assessments were compared with baseline data to determine the positive effect of the intervention. Moreover, qualitative data collected from interviews with families provide a deep understanding of the treatment effect.

Quantitative Data Analysis. In this section different kinds of statistics are used. The description of these statistics can be found on the below mentioned figure, but is not restricted to only that: descriptive statistics (mean, standard deviation, etc) inferential statistics (t-test, ANOVA) correlation analysis regressions analysis (Field, 2018).

Qualitative three-level thematic analysis of the interview data is presented here, highlighting main themes and sub-themes of family experiences with tDCS, and providing participants' quotes to help contextualise the qualitative findings (Braun and Clarke 2006).This section brings together the qualitative and quantitative data and looks at how they agree or disagree with each other where there are areas of overlap. Bringing the two sets of data together offers a more rounded picture of the results and enhances the validity of the findings (Creswell Clark, 2017).

The following section summarises the impact of tDCS with regard to improving social communication abilities and cites case examples that report on individual progress. Statistics are presented examining the more theoretical significance of this, with regards to practical applicability in therapeutic practice.

Number 8. This section contains a summary of family reactions and experiences to the intervention of tDCS. Feedback from families provides a useful source of information about the

acceptability of the intervention from a practical viewpoint. This section presents initial exploratory studies including subgroup analysis (eg, age, gender) or assessing potential moderators and mediators followed by preliminary results to guide future replication studies. The conclusion sums up the main results, discusses the theoretical and practical implications, and spells out the limitations of the study. Such a meticulous approach guarantees that the results will be presented in a thorough and orderly fashion, ready for rigorous scrutiny and analysis of the data.

41.2 provides an overview of the demographic characteristics of the participants.

The demographic profile of the study participants provides valuable contextual background information about the effects of tDCS on social communication in children diagnosed with Autism Spectrum Disorder (ASD). It is important to know the demographic profile in order to ensure that the sample is representative and to explore the possible effects of varying demographics on the outcomes of the study.



Figure 1: Gender Distribution

Age Demographics

The subjects of this study ranged from 3 to 18 years of age. Participants were split into several age groups to facilitate a more thorough analysis and comparison of results across ages. The percentage of the overall sample comprising of subjects aged between 6 to 12 years stood at 57% (Q1 Survey Data). Age group that ranges between the ages of 6 to 12 years is the most important group in this study because there are peak ages of development where intervention would contribute a lot to the improvement of social communication skills.



Figure 2 displays the distribution of participants according to their age.

The allocation was as follows:

• Of the population, children between the ages of 3 and 5 make up 14 per cent.

• Population of children from 6-8 years old forms 29 percent of the whole.

• Ages 9-12 make up 28 per cent of the population

• The percentage of individuals between the ages of 13 and 15 is 18%.

• The percentage of individuals aged 16-18 is 11%.

The choice of age-range was made with the intention of including a good mix of children from different developmental stages, and thereby to isolate some of the underlying reasons for the variability in the effectiveness of tDCS at different periods of development. By using a wide range of ages, the study hopes to capture the diversity of problems and possibilities in intervention for all different age groups.

Gender Allocation

While the majority of subjects were male – which is representative of the epidemiology of ASD – Q2 revealed that 265 (100%) participants out of the 347 subjects were of the male gender, thus accounting for 79.25% of the total, while 61 (17.75%) were of the female gender, and 7 (2%), of the third possible option: 'prefer not to say'. The male/female ratio of around 4:1 is in keeping with the conventional view that ASD is disproportionately represented in males (Baio et al., 2018).

The allocation was as follows:

There were 210 male participants.

There were 55 female contestants.

Being able to recognise the fraction of males and females in the sample is important because it allows you to examine whether there is a difference in response to tDCS based on gender. It's been found that there can be gender-specific differences in symptom presentation and symptom progression for Autism Spectrum Disorder (ASD), which might be related to differential responses to therapeutic therapies (Lai et al., 2015).

Level of economic and social standing in society

Socioeconomic status (SES) can play a decisive role in health and developmental and education, which are critical for child growth and wellbeing. In this study, we used a variety of measures of socioeconomic status including paternal education, paternal occupation, and household income to assess SES exposure.

41.3 Parental Education

Most of the parents had varying educational degrees, a good percentage having been to university:



Figure 3: Level of education achieved in higher education.

• 37.21% of parents possessed a college degree

• 25.19% of those had higher education – five or more college courses completed.

• 22.64% possessed a high school diploma

The result of the survey data shows that 14.96 percent of population were high school educated or lower. Parental education is a key facet of SES since it often overlaps with other SES indicators, including income and occupational status. Higher levels of parental education are usually associated with greater opportunities for access to healthcare and educational resources, which in turn can affect the efficacy of interventions such as tDCS (Bradley Corwyn, 2002).

41.4 Occupational Status

The occupation of parents was categorized in three groups: Professional, skilled and unskilled. Their distribution is given below.

•Professional jobs such as doctors, engineers and teachers comprise 46.42 per cent of the total.

• Highly proficient individuals (e.g., technicians, artisans): 34.72%

• People who were non-skilled workers included labourers and service staff. They accounted for 18.86% of the study population (Survey Data, Q11).

And an individual's employment status provides an important baseline of information about their economic status and social context. While jobs in the professional class often correlate with higher wealth and better access to healthcare, this correlation can have a significant impact on the efficacy of ASD therapies.

41.5 Household Income

Income levels from different households were be in assigned to three different grades, namely low, middle and high-income levels. The allocation was as follows:



• The percentage of individuals with low income is 21.13%.

The percentage of individuals with middle income is 48.30%.

• Percentage of individuals with high income: 30.57% (Based on survey data, question 12)

As such, income is a powerful proxy for SES and a direct measure of the ability to afford any forms of therapies, educational supports or even essential everyday provisions for subsistence that might be required to manage children with ASD. Higher income, all else being equal, means greater access to initial intervention and longer-term support – both of which are necessary for a child with ASD.

42. Socioeconomic Consequences

This socioeconomic-status (SES) distribution of the sample indicates a varied populace with different levels of resource access. The variation allows us to explore the extent to which socioeconomic conditions might influence response to tDCS. For example people from a high socioeconomic-status (SES) family have substantial and reliable access to medical and education services, which may enhance response to tran-scranial direct current stimulation (tDCS). In contrast, low-socioeconomic-status (SES) individuals may have fewer resources and, consequently, the benefits of treatment may be impeded (Durkin et al., 2010).

43. Academic History

This took the form of a questionnaire on the educational background of the participants, because educational environments are important in enabling the development of any social-communicative abilities, particularly in children with ASD. Participants ranged from being in regular schools to specialised education programmes, or were being home-schooled. The split was as follows:



Figure 5: Educational background of the participants.

• The percentage of students enrolled in mainstream schools is 68.05%.

The percentage of special education programmes is 20.30%.

According to survey data from question 7, homeschooling accounts for 11.65%.

Usually, in mainstream schools, children on the spectrum are in mixed (inclusive) education settings with neurotypical classmates who, potentially, could help them engage more socially and hone their communication skills with regular interpersonal exchange. Again, it's complicated: on the one hand, children with ASD might need additional support to deal with the complex social dynamics of a mainstream setting, yet on the other, comprehensive support and assistance could help them navigate the social interactions that might otherwise have been puzzling.

Special educational provisions are tailored to individual needs of children with disability so, for example, for a child with ASD these may include specific needs-led support and interventions which can be very valuable, and can be effective in support of social communication. Homeschooling may mean that children miss out on the chance to practise these skills with typically developing peers, and being able to apply such skills in various contexts. Even though homeschooling may lead to a more personalised, targeted teaching approach, it may also limit opportunities for socialising. Homeschooling parents typically actively make seeking socialising opportunities a priority, joining community groups and participating in clubs, to help their children acquire and practise social skills.

43.1 Supplementary Academic Assistance

Furthermore, in addition to their mainstream schooling, a large number also received educational supplements – speech therapy, occupational therapy and social skills training – aimed at addressing the specific challenges faced by children with ASD.

Speech treatment accounts for 45.28% of the total.

Occupational therapy accounts for 32.08% of the total.

According to survey data from question 8, social skills training accounted for 38.87%.

These add-on supports are important to augment the benefits of tDCS for speech therapy to better address the communication impairments that are often related to individuals with ASD, and for occupational therapy to help with sensory integration and fine motor skills that can be a challenge for children with ASD.

44. Implications for Education

The wide range of educational attainment among the participants emphasises the variety of means by which families assist their children with ASD. Understanding the different types of education environments is key to interpreting the results of this study, because how tDCS works might be affected by whether more or less assistance is given. Children with ASD in special-ed programmes might have distinct response-patterns to tDCS compared with those in regular schools, mainly as a result of differences in the level of assistance and opportunities for social interaction.

To sum up, the study's demographic information provides a vast sample for understanding the impact of tDCS on children with ASD's social communication. The 8-14 year and 15-19 age groups reflect the 2 most prominent periods that have significantly higher incidence rates of ASD. The separation of gender helps distinguish the effectiveness of the treatment for boy v. girl v. mixed samples. The various socio-economic statuses and levels of education diversify the college-going population and ensure sampling of adults that have various living conditions so that the treatment is assessed in a real-world setting.

45. Results of the initial assessment

We conducted the baseline assessment of social communication skills to get insight of the initial abilities of children with Autism Spec-trum Disorder (ASD) prior to the transcranial Direct Current Stimulation (tDCS) treatment. communication skills. That is, they show impairment in both spoken and non-spoken communication, social interactions and cognition of social signals. The baseline assessment of social communication skills afforded us a better understanding of participants' initial abilities. It is notable that the one with ASD often have difficulty not only in spoken communication but also in other communication areas such as non-spoken communication and social interactions.

Before the tDCS intervention, the children varied too in linguistic abilities, 'with many expe-riencing difficulty initiating and maintaining talks'. Common problems included a reduced vocabulary, difficulty in forming sentences, and echolalia (the repetition of phrases or sentences of others). Almost two-thirds had problems with how they expressed themselves verbally (SRS scores). These problems hindered their ability and willingness to communicate their needs, ideas and feelings to others. The inability to communicate led to frustration and behavioural

problems when their needs were not met. Being un-able to communicate, the children had no tools to perform an essential aspect of socialisation.



Figure 6: Enhancements in Verbal Communication

Closer examination of social reciprocity challenges revealed that 75 per cent of the participants exhibited substantial difficulties in engaging in reciprocal interactions. Problems included difficulty responding to social initiations from others, limited sharing of interests and feelings, and a general lack of reciprocity in social interactions. The deficits were further substantiated by low scores on the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS), which measures social awareness, social cognition, social communication and social motivation (Constantino Gruber; 2005 Lord et al; 2000).

Furthermore, many showed deficits in emotional understanding and expressivity, often having difficulty in spotting and understanding others' feelings, as well as adapting their behaviour to social contexts. Their low social insight and knowledge interfered with building and maintaining friendships, as well as with relationships with adults.

The fundamental deficits of social communication were likely to have pervasive and far-reaching impacts on the children and their families on a daily basis. Parents reported frequent incidences of miscommunication, social isolation, and frustration stemming from their child's limited ability to engage appropriately and communicate effectively with others. These factors were likely to bear on social contacts, in addition to the children's capacities to participate in school and recreational pursuits.

There was also an assessment of nonverbal communication, such as visual input (eg, eye contact), expressive and receptive facials (eg, smile, frown) as well as gestures and body movements. Of the 70 per cent of the participants struggling to communicate nonverbally, 60 per cent were rated 'above threshold' on the Autism Diagnostic Observation Schedule (ADOS) (Lord et al., 2000). An example of how this score breaks down is with nonverbal communication; it has often been observed among the youngsters that they are particularly averse to eye contact, with limited facial expressions, as well as challenges when it comes to the appropriate use of gestures.

During interactions, the lack of eye looking created an awkward environment for communication, increasing chances of misunderstandings and miscommunication. The reason that nonverbal communication impacts so heavily is because children rely on such communication in their day-to-day social relationships. Deficits here can seriously hinder the child's ability to engage in a meaningful and positive way with peers and adults. A third critical dimension of assessment was the ability to partake in mutual social interactions. It was observed initially that 75 per cent of the participants had significant difficulties in social reciprocity. These children often experienced problems in turn-taking in conversations, interpreting social cues and understanding the perspectives of others. Q4 score revealed that they were less enthusiastic to initiate social interactions and were more likely to prefer spending their time alone with solitary tasks

rather than participating in group play. Lack of mutual social interactions is a feature presenting in ASD, and social isolation can have implications on the child's emotional and psychological development.

Numerous subjects showed many deficits in deciphering social signals, such as understanding emotional expressions, identifying social demands, and accurately responding to others in social scenarios. The majority of children (about 60 per cent) showed deficits in interpreting social signals, as assessed by their score on the SRS and ADOS. Their poor social cue-detection skills made it difficult for them to form and sustain meaningful relationships with both other children and adults. Misreading social signals usually led to incorrect reactions in the social context, which added to the peer-rejection and led to a vicious circle of social dysfunction.

Baseline assessments measured participants' behavioural, cognitive and social communication profiles, giving us a detailed understanding of each person's strengths and weaknesses prior to starting the tDCS. This allows us to quantify any potential 'benefits' that tDCS might confer.The first behavioural assessments revealed that the sample subjects displayed a significant frequency of repetitive and restricted behaviours, which typically inhibited the child's flexible, adaptive behaviour and interfered with normal functioning. Their high RBS-R scores reflected that these behaviours were wide-ranging and of significant functional impact (Bodfish et al., 1999).

In addition, the majority of the subjects also had problems with emotional regulation. They had high levels of mood lability, uneasiness, and temper outbursts or self-injurious behaviour. The ABC was used to assess these problems with emotional regulation, and produced an overall description of the behavioural problems the children presented with (Aman et al, 1985).

Also, the behavioural problems exerted a lot of influence on family dynamics and on the family stress and burnout. The parents of the participants felt high levels of strain and exhaustion because they needed to constantly cope with their children's behavioural problems, and the siblings also felt emotional duress because ASD family dynamics often centre around the needs of the child with ASD. Children with ASD usually exhibit repetitive, restricted behaviours, interests and are very sensitive to sensory stimuli. During our initial perceptions, we conducted behavioural evaluations using standardised test instruments such as the Repetitive Behaviour Scale-Revised (RBS-R) and the Aberrant Behaviour Checklist (ABC).

RBS-R scores revealed that 72 per cent of the children exhibited unusual behaviours in the categories of stereotyped movements (eg, hand-flapping, rocking), insistence on sameness, and ritualistic behaviours (ie, adhering to specific sequences of activities) that disrupted everyday life (Bodfish et al, 1999). Repetitive behaviours can bring some comfort to children with ASD, but they can also interfere with their ability for flexibility and adaptation, particularly in social and educational settings.

Moreover, about two out of three (68 per cent) showed a marked preference for certain topics or items; a sign of restrictive interest. The focus is typically given over to these interests, invading conversations and activities, so that the young person is unable to participate in broader experiences. These highly focused interests might be utilised for educational purposes; but the extent to which children are often now 'bound' to their specific interests limits their ability to engage in other activities with their peers, which again serves to isolate them from the social world.

In a follow-up parent survey, 65 per cent of parents reported that their children experienced sensory sensitivities, such as to sounds, lights, textures and other stimuli, and these sensitivities led to uncomfortable, anxious or behaviourally 'off-base' responses in the context of their everyday settings (Survey Data, Q5). Given the extent to which this requires modifications to everyday schedules and settings, this can place pressure on the child to minimize his participation in usual social and educational pursuits.

Executive functions were evaluated with the Wechsler Intelligence Scale for Children (WISC), the Behaviour Rating Inventory of Executive Function (BRIEF). The cognitive assessments showed the dysfunctions of executive functioning that the participants were experiencing. These dysfunctions were more evident in the uses of working memory, cognitive flexibility and inhibitory control, for example. The BRIEF scores showed that these impairments of executive function were observed in different settings: family or school situations.

Their deficits of executive function had a direct impact on their scholastic achievements. Many of the children had trouble completing tasks that required sustained concentration, systematic planning and analytical thinking capacities. They also revealed difficulties on academic records, in subjects that called for executive functioning to be at its highest levels.

Third, the ratings assessed adaptive behaviour, which includes practical, social and conceptual abilities needed to navigate through life's daily circumstances. A majority had serious impairments in adaptive behaviour, which limited their ability to function independently in their daily life. The delays were evaluated using scales such as the Vineland Adaptive Behaviour Scales that provide a comprehensive picture of participants' adaptive functioning (Sparrow, Cicchetti & Balla, 2005).

Many children with ASD have impairments in executive processes – including working memory, cognitive flexibility, and inhibitory control. According to Gioia et al (2000), 70 per cent of the individuals evaluated had executive process impairments. For instance, many kids struggled with chores, such as folding laundry, that required the ability to organise, organise and problem-solve. The impact of these deficits can be extensive. Academic performance can suffer and, more importantly, daily life can quickly become overwhelming. Executive functions enable us to manage our activities and respond appropriately to unprecedented circumstances; children with ASD can perform inconsistently in these contexts.

But 65 per cent of the study's subjects singled out working memory, the ability to hold and manipulate information for short spans of time, as especially challenging. The subjects' trouble with working memory hindered their ability to grasp and carry out complex instructions, as well as tasks that require sustained attention, such as reading until completion. This deficit is often exposed in school settings where assignments require the integration of multiple chunks of knowledge and the ability to maintain focus for long periods of time.

About 60 per cent had weak Inflexible cognitive habits Hard to adapt to new Plan changes Hard to manoeuvre around uncertain circumstances This is a manifestation of poor cognitive flexibility, the ability to switch your thinking and shift in response to new information or to changing circumstances. Thus, weakness in cognitive flexibility could hinder a child's ability to change habits or adopt new routines if they aren't well-learned, or to make adjustments on the fly when things shift or become uncertain. The child may end up having a lot of trouble with solving problems, or adapting to changing social situations – which fits with the picture of children with ASD struggling with the very same kinds of tasks.

My other interest was inhibitory control: the ability to suppress inappropriate or impulsive behaviour. About 58 per cent of our subjects showed significant deficits in inhibitory control, which leads to impulsive behaviour and difficulties with emotional and neurological regulation. If a child has inhibitory control problems, it will manifest in various ways, including frequent social and academic interruptions. The child will have a hard time regulating interactions with peers and adults.

46. Evaluation against established standards

These presented the ASD group's baseline assessment results, then compared them with average results from healthy, typically developing children, so as to highlight the areas of weakness it's characteristically Makes it easier to understand To better compare the therapeutic intervention to other types of treatment, we also made pairs of bar graphs that displayed the results for a control group and two different types of tDCS interventions: cathodal stimulation over langual regions, and anodal stimulation over visual regions. As you can see, each presentation had a corresponding legend with terms that highlighted the areas of weakness it's characteristically more difficult to learn, and indicated to what group of individuals it applied.

These subjects' scores, when compared with normative data, indicated much worse deficits in their abilities for social communication. Children (who are TD) are typically in the average range on the SRS, which measures verbal and non-verbal communication, social relationship skills, and social recognition, and on the ADOS, which assesses language and communication, reciprocal social interaction, and imaginative play skills. But on both of these scales, the subjects in the research reported significant deficits in all three of these areas. These comparisons illustrate the substantial challenges in social communication that children with ASD struggle with and provides a clear indication that therapies like tDCS could be very beneficial in aiming

to improve the impairments in these areas. In comparison to TD children, these subjects with ASD scored significantly worse in every way in SRS and ADOS – depicting significant deficits in social communication. The normative data tells us that TD children are more socially aware, communicate and interact at a higher level. Children with ASD struggle with these abilities and a specific treatment to improve these abilities is necessary.

Behavioural assessments revealed differences between children with ASD and typically developing children as measured by increased scores on the Repetitive Behaviour Scale-Revised (RBS-R) and the Autism Behaviour Checklist (ABC). This suggests more frequent and widespread behaviours for the children diagnosed with ASD. Children with typically developing minds had fewer repetitive behaviours, fewer narrow and intense interests, and decreased abuse of objects and self-injurious behaviours than those with ASD, which indicate how behavioural difficulties are different in ASD compared with usually developing children (Bodfish et al., 1999; Aman et al., 1985).

Self-injury represents a range of behaviours from firmly hitting oneself or pulling one's hair to severely restricted alimentation that can lead to illness or death Self-injury includes actions driven by impulses and environmental factors. It is challenging to assess whether this is a typical behaviour because many individuals with intellectual disabilities injure themselves in environments lacking consideration for physical and mental wellbeing. Self-injury represents a continuum of behaviour from stubbornly hitting oneself or pulling one's hair to overlapping with restricted alimentation that can include a lack of eating or defecation, which increases the risk of serious illness or death.

Children with ASD showed greater repetitive behaviours, narrower interests, and heightened sensory sensitivities than typically developing children. In 1999, Bodfish et al found that the general population demonstrated not only fewer symptoms but also less severe symptoms on the RBS-R and ABC assessments. These intensified behavioural phenomena help us to understand that people with ASD have very specific needs, and why targeted therapeutic approaches that target these behaviours are needed.

We observed strong impairments of executive processes in the ASD group relative to the normative data. Normal children outperformed children with ASD on most measures of working memory, cognitive flexibility and inhibitory control, which is associated overall with more effective executive functioning (eg, Gioia et al, 2000). Cognitive deficits in ASD can undermine both children's academic performance and adaptive functioning. Therefore, it is crucial that interventions facilitate both cognitive skills and social communication abilities.

47. Significance of Baseline Comparisons:

Their group scores also highlight the significant challenges that children with ASD face in using social communication in any setting, exhibiting socially inappropriate and challenging behaviours, and performing poorly on every measure of cognitive abilities. Differences on the first three measurements serve to underscore the need for targeted therapies like tDCS allowing us to specifically target those impaired areas.The large differences observed between the ASD and TD groups provide a unique framework for evaluating the potential benefits assumed to be arising from the tDCS intervention. Reinforcing differences noted on the neurocognitive measures were differences observed on the AASEH parental questionnaire for metacognitive abilities On measures of cognitive abilities, we noted differences between the ASD and TD

groups with TD children showing better executive functions as demonstrated by the higher BRIEF scores. They performed better on demonstrative and precise working memory, cognitive flexibility and inhibition, which are hallmark abilities associated with executive functions. These differences demonstrate the fact that children with ASD do experience cognitive impairments in executive functioning, as demonstrated by their poor overall performance on academic and daily life pursuits, as highlighted in an earlier study (Gioia et al., 2000).

Overall, the ASD subjects scored significantly lower than TD children on all adaptive behaviour measures. When comparing the ASD subjects to TD normative data from the Vineland Adaptive Behaviour Scales, the TD children were more independent in daily living skills and had better social and practical skills. The ASD subjects' delays in adaptive behaviour suggested that their difficulties in developing independence and age-appropriate functioning caused them greater problems than the TD children. (Sparrow, Cicchetti, & Balla, 2005).

These measures compared the participant's results with a group of children who had been carefully chosen because their scores fell within an expected range for people without developmental or social impairments. These comparisons with normative data seemed to provide a deeper insight into which areas of social communication showed the most marked differences for the children with ASD. The fact that these initial differences existed also suggested that therapies such as tDCS should be aimed at specific areas of social communication, behaviour and cognition that are influenced by the condition. The way that we analysed DBS in people with Parkinson's disease suggests that tDCS had the potential to shift behaviour in the ASD group in ways that would be targeted to meet the unique needs of their condition.

48. Social communication skills after receiving treatment

The rationale behind this post-treatment approach was to assess the impact of tDCS in improving the social communication abilities of children with ASD, providing an overall assessment of the changes that occurred in their spoken and unspoken communication, social exchanges and ability to grasp social signals following the tDCS intervention.

Post-treatment results indicated considerable improvement in verbal fluency. On the Social Responsiveness Scale (Constantino Gruber, 2005), a full 60 per cent of the children showed significant improvement in abilities to initiate and sustain conversations. Many displayed both an increase in vocabulary and in sentence structure, with much of their talk being more widely logical and eloquent. Children who had been known to produce monosyllabic utterances or very short phrases began to elaborate into multi-clause sentences, allowing them to express more effectively their wishes and ideas.

The research also found a reduction of echolalia. This revealed that the children were showing increased ability to produce spontaneous vocalisations as answers, rather than merely echoing questions or statements spoken to them. Moving away from echolalia to spontaneity is an important step in the refinement of communicative skills. Improvements in nonverbal communication were evident, including sustained eye contact, facial emotions and gestures, as well. Approximately 65 per cent of the children were rated higher, post-treatment, in these behaviours on the Autism Diagnostic Observation Schedule (ADOS; Lord et al., 2000), a standard approach for evaluating autism. These modifications allowed for more robust and engaging interactions with peers and adults. As a result, the children's ability to communicate their feelings and goals without words had more significant impact. Nonverbal communication is thought to play a vital role for establishing rapport and for providing cues to understanding

social interactions. For example, children who would not previously look at people's faces during conversations began to maintain eye contact much longer, thus raising their level of engagement and responsiveness. The increased use of gestures such as pointing to things or nodding their heads also provided additional information to supplement what they were saying and doing, thus offering additional clues to clarify spoken messages and their intentions.

Overall, 70 per cent of the children showed demonstrable increases in social reciprocity. At a follow-up evaluation one year later, children showed increased initiation of social relationships, increased responsivity to social cues, and increased conversational talk. Improvements in social reciprocity were crucial for relationship formation and maintenance with peers and adults as reflected in increased SRS and ADOS scores following treatment. Increased social reciprocity led to greater participation of children in co-operative gross-motor activities as well as more spontaneous play with peers. Social play is the context in which children acquire social skills and their peer relationships develop. For example, children previously described as inactive or avoidant in social situations started requesting and engaging in play with their peers by taking turns and sharing toys. The children showed dramatic improvements in peer acceptance and were no longer regarded as isolated or socially rejected.

As a consequence, their ability to understand social cues improved enormously. Almost 65 per cent of the kids benefited from better capabilities in recognising emotions, social rules, and proper reactions in various social contexts. These improvements all contributed towards better understanding of various social settings and, consequently, lesser misunderstandings and quarrels in social interactions. Understanding social cues more sensefully made it possible for youngsters to adapt their behaviours to respective social contexts – to respond in a more ade-

quate and flexible fashion. For instance, kids that previously used to be incapable of recognising when someone was distressed started showing signs of care and consideration, which promoted the development of better social relationships.

Parents and carers reported meaningful changes in children's day-to-day interactive capacities after tDCS, including heightened engagement and responsiveness during family conversations, measurably increased interest in social activity and measurably improved competencies in forming friendships. Changes of the type outlined above were reported to have had a substantive impact on children's quality of life and social welfare, including, for example, a cessation of children's social isolation and related difficulties forming friendships (where such difficulties had previously been reported), to now being involved in play dates and social occasions with peers, in turn leading to increases in self-confidence and social capabilities. Specific positive changes were also reported to occur with respect to children's changes in social anxiety and irritability, where these arose, including measurable improvements in children's 'social calibration' dynamics involving social interaction competencies and, furthermore, improved modulation of the perception of others' responsiveness and emotional states. Day-to-day changes of the sort reported were supported by qualitative comments from parents detailing measured changes in family dynamics and children's emotional and social functioning.

Further case reports confirm this finding of tDCS effect on social communication and another study showed that these effects can last for 60 minutes following the electrode application. The example of the 10-year-old male shows the severity of pre-existing communication challenges (verbal and nonverbal) before the tDCS therapy, and the significant gains in the SRS scores, particularly in the subdomain of expressive language, and in another subdomain of social motivation. As reported by his parents, he started to use longer sentences, was more proactive

initiating conversations with his family members, and displayed a real desire to socialise with his friends. The report on the 7-year-old child showed that she had little verbal communication and was limited in non-verbal (eye contact) communication before the tDCS; however, after the therapy session, she started communicating with verbal language when making a request, had eye contact regularly, and used gestures for better communicating. These case studies provide another glimpse of the effect of tDCS on social communication skills, all viewed as salient improvements, granting the targeted participants greater ability to engage with others in the surrounding context and participate in daily activities.

The assessment of social communication skills demonstrated that spoken and non-spoken communication, social interactions and decoding of social signals were significantly improved by the treatment in the ASD group. The improved social communication enabled improved social integration and contact, and these in turn improved their quality of life. The findings presented in this paper demonstrate the possibility of tDCS as an intervention to improve social communication functioning in ASD. Given the promising trends, this is a promising direction that warrants further investigation and clinical translation.

49. Alterations in behaviour and cognition

For assessing the missing piece: Is it also working on their behaviours and cognitions such as executive functions and social communication after treatment? This was the missing piece for assessing the broader changes brought by tDCS in a child with ASD.



Figure 7: cognitive functioning levels both before and after the treatment.

Repeated Behaviours: One of the cardinal symptoms of ASD is the repetition of movements, sounds or objects, along with a narrow, restricted interest. Post-treatment RBS-R scores reflect substantial decreases in these repetitive behaviours for 65 per cent of participants (Bodfish et al, 1999). Specifically, those children who repeated movements by, for example, hand-flapping, rocking or otherwise engaged in stereotypic movements either eliminated these behaviours altogether or greatly reduced those movements in frequency and intensity. Lessening the repetitive behaviours allowed these children to more actively participate in many activities and interactions that otherwise would have been challenging; thus, these children became more flexible and engaged in carrying out and performing everyday life activities.

Emotional Regulation: The emotional regulation benefit was very consistent following the tDCS treatment; parents reported that approximately 60 per cent of the children once again found it possible to handle the normal vicissitudes of living, where previously they would have

had an outbreak or a meltdown. As one parent said: ‘Our child can finally regulate himself again.’ She meant that children with autism tend to either get extremely upset or shut down when things don’t go their way, but now, her son dealt with frustrations such as minor schedule changes much more calmly than before. He had learned emotional regulation again.

Enhanced Behavioural Flexibility: There were also improvements in behavioural flexibility. A large number of children showed increased flexibility to varying daily routines and became more willing to try out new experiences. Improved behavioural flexibility was necessary for participating in many social and educational environments. Reduction in the demands for stereotypical and repetitive behaviours led to increased flexibility and responsiveness to new experiences, important for holistic development and for socialisation.

Executive functions – that is, working memory, cognitive flexibility and inhibitory control – are often impaired in children with ASD, and post-treatment assessments on the Behaviour Rating Inventory of Executive Function (BRIEF) (Gioia et al., 2000) indicated significant improvements for ~ 55 per cent of the participants.

Enhanced working memory: These improvements in working memory were particularly noteworthy. Kids improved their ability to temporarily store and manipulate information for short periods of time, a skill important for following complex directions and solving problems. Improved working memory translated into better grades and better ability to complete more complicated activities. For example, kids demonstrated improved ability to follow teacher directions and complete homework with less help.

Another psychological marker that showed improvement was the participant's cognitive flexibility – their braining's powers of shifting set, or the ability to shift how one is thinking; in particular, to shift one's thinking and modify one's responses to novel demands. Once they were in guided-meditation mode, participants became more adept (quickly) modifying the way they were thinking and behaving based on the new information or new conditions that faced them. They demonstrated a greater ability to quickly shift between sets (to shift between tasks) based on a change in context. Improved cognitive flexibility facilitated better problem-solving as well as being more flexible in terms of learning and socialising.

Inhibitory Control: The ability to hold back inappropriate or impulsive responses – known as inhibitory control – was also improved. After treatment, kids exhibited better regulation of impulsive behaviour and more goal-directed decision-making. Improved inhibitory control played a role in reducing disruptive behaviours at school and home. Parents and teachers reported less impulsive behaviours – interrupting during a discussion or not being respectful – leading to a calmer and more productive environment.

Overall Cognitive Abilities: The executive processes development resulted in an overall improvement in cognitive abilities. At post-treatment, many cognitive areas such as verbal comprehension, perceptual reasoning, and processing speed were improved in the majority of subject (in accordance to the Wechsler Intelligence Scale for Children, WISC) (Wechsler, 2003). An improvement in the cognitive abilities played a crucial role in better performance in school and greater ability for functional autonomy.

As a result of the treatment, their general behaviour and ways of thinking seems to have improved considerably, positively affecting the children's daily lives and those of their families.

Improved control of the emotional part of their brain, reduced repetitive behaviours and improved cognitive skills enabled them to participate more purposefully in everyday life and with more enjoyment. Parents reported that their children became considerably more functional, cooperative, involved and competent in dealing with the multiple challenges of everyday life. The beneficial changes in the children's behaviour resulted in better and more enjoyable family experiences and a reduced stress and burden for their families.

The magnitude of the behavioural and cognitive enhancements was documented in several case studies. One of them was an 8-year-old boy who initially had severe problems with emotional regulation and had frequent episodes of angry outbursts. After treatment, the child's ABC scores showed clear reductions in irritability and hyperactivity; moreover, his parents reported he was much calmer and substantially easier to manage than before the tDCS intervention. Another child, a 9-year-old girl, had some challenges with her working memory and cognitive flexibility. Subsequent to tDCS, she showed significant improvements in the relevant subscales of the BRIEF; furthermore, her teachers reported that she was now able to complete more school tasks that required attentional focus and cognitive flexibility than before the tDCS intervention.

Following the intervention, children showed significant improvements in repetitive behaviours, emotional regulation, executive functions and cognitive capacities in general, translating in better functioning at school and in their social lives, and improving their quality of life. These results demonstrate how tDCS can be used as a therapeutic intervention for enhancing behavioural as well as cognitive outcomes in children with ASD. It's a very encouraging finding for future clinical research and use.

50. Results of the assessment conducted after the treatment

In this part, the post-treatment evaluation by children after using tDCS as a supportive treatment for ASD is analysed to show the observed signs of change. This is divided into two main categories: Post-Treatment Social Communication Skills and Behavioural and Cognitive Changes. In each category, a detailed summary of changes that were observed is provided inclusive of both quantitative and qualitative data from parents and carers.

Social communication is a vital component of autistic children's development. This involves the spoken and non-spoken aspects of engagement that are so important for social contact. Post-treatment testing compared the role of tDCS in improving these social capacities (ie: conversational initiation; maintaining eye contact; using gestures and facial expressions; and overall participation in social communication).

Enhancements in Verbal Communication

• The mean baseline score was 3.2 on a scale of 1 to 5.

• The average score after therapy was 4.0, measured on a 5-point scale.

• The resultant data is thus deemed statistically significant (t-value of 8.45 with 264 degrees of freedom and p-value smaller than 0.001)

• The effect size is represented by Cohen's d value of 0.52.

The statistics show an improvement in verbal communication skills. Parents found that youngsters were now more likely to initiate conversations, express their ideas more articulately, and use a larger vocabulary. The small effect size is significant: the changes were both statistically and practically meaningful.

Nonverbal communication

• Nonverbal communication through physical movements and facial expressions.

The average score before therapy was 2.8.

The average score after treatment is 3.6.

The statistical analysis yielded a significant result: $t(264) = 7.89$, $p < 0.001$.

The effect size, as measured by Cohen's d, is 0.48.

There were also major changes in their use of gestures and facial expressions. Children started using their hands and faces more frequently to indicate emotions and intentions, and thus also became more effective communicators nonverbally.

Interpersonal involvement

• Socializing with peers and family members

The average score before therapy was 2.5.

The average score after treatment is 3.3.

A high t-value of 6.78 with 264 degrees of freedom and p-value less than 0.001 confirms the data's statistical significance.

The effect size, as measured by Cohen's d, is 0.42.

More social engagement with peers and family was noted, and an increase in propensity to work together, share interests and respond to social cues was documented. The greater social interactions observed suggest that tDCS can really help with the sociability of children with autism.

Subjective Evaluation

Parents offered insightful qualitative comments, pinpointing particular areas for growth and identifying obstacles.

Beneficial Transformations

My son now says hello to people when they say hello to him, something he didn't do before.

She is using a higher number of gestures and expressions to help indicate what she wants.

Obstacles

My son was initially unwilling to challenge himself in social environments. But then he showed changes. These were subtle at first.

Though there was visible improvement in spoken conversation, we are still not capable of maintaining eye contact. Overall, the qualitative feedback complements the quantitative metrics, showing both the gains and continuing challenges in improving social communication abilities with tDCS.

50.1 Alterations in behaviour and cognition

All behavioural and cognitive changes represent the overall effects of tDCS on autistic children as a whole. The results of this part of the essay analyses the modifications of maladaptive behaviours, self-control skills, concentration and the cognitive abilities after the completion of the procedure.

Enhancements in behaviour

• Decrease in troublesome behaviours

The average score before therapy was 2.5.

The average score after treatment is 3.3.

This finding is confirmed by the statistical significance of the data, which obtains for the t-value of 6.78, 264 degrees of freedom and p-value less than 0.001.

The effect size, as measured by Cohen's d, is 0.42.



Figure 8: At the end of the study there was a significant decrease in behavioural problems.

Further, he notes that prior to the treatment, the children were noted to have tantrums, hostility, and non-compliance, which radically reduced by the end of the study creating a peaceful atmosphere at home.

Emotion regulation

The average score before therapy was 2.9.

The average score after treatment is 3.8.

The stat value of the t value is 9.32 with 264 degree of freedom, p-value <0.001.

The effect size is represented by Cohen's d, which has a value of 0.57.

These results also suggested improved emotional control: children showed better emotion regulation, lower mood variability and better frustration tolerance. Perhaps most noteworthy is that the effect size here was large, demonstrating the meaningful impact of tDCS on emotional stability.

Concentration and attentiveness

• The average score before therapy was 3.0.

The average score after treatment is 3.7.

Is the difference statistically significant?

The answer is a resounding yes: these data have a t-value of 7.45 with 264 degrees of freedom and a p-value of less than 0.001.The effect size is represented by Cohen's d and its value is 0.45. Attention and focus was also boosted and, as a result, children not only improved their capacity to stay on task and follow through with instructions, but also showed improvements in academic performance and day-to-day functioning.

Cognitive functioning encompasses your thinking and remembering. It involves mental processing such as attention, the organisation of thoughts, and consciousness.

• The average score before therapy was 2.9.

The average score after treatment is 3.8.

Figure 2: Typical scenario of the model.This test results show the power of the data which is; t-value = 9.32, Degrees of freedom = 264 and p-value is <0.001 .

The effect size is represented by Cohen's d and has a value of 0.57.

The investigation found large improvements in executive function/problem solving, memory, information processing and speed of cognitive processing. Evidence suggests tDCS promotes cognitive development in autism children. Parents provided detailed descriptions of their children's improved behavioural and cognitive performance.

Positive observations

'There have been drastic decreases in meltdowns, and his level of calmness has increased.

• Her ability to focus on her schoolwork has significantly improved. She is completing homework that previously was difficult for her to do.

Areas need improvement

Yet, despite better conduct, sometimes we still experience those latest, best, prominent and identifiable feelings, in unaccustomed and somewhat untimely circumstances.

Although the focus of attention has become more durable, retaining it for prolonged periods continues to be a challenge. These comments underscore the net-beneficial effects of tDCS, but also reflect additional neural regions that could use some remediation.

Quantitative evaluation of the scores before and after treatment through statistical tests. A hallmark of examining tDCS effects on children with autism is the analysis of pre- and post-treatment test scores and this portion of the report will present the relevant data, followed by interpretations and a thorough review of the results based on the data analysis.The scores of the children participating in the survey before and after receiving tDCS treatment are compared across multiple parameters below, such as verbal communication, social interactions, habits, and behaviours as well as cognitive abilities and social capabilities.The survey collected numerical data on the children regarding their behaviour and capacities prior to and after the administration of tDCS treatment. However, the most important areas under study are:

- Proficiency in spoken communication
- Interpersonal skills
- Alterations in behaviour
- Mental and emotional control and management
- Scholastic achievement

To analyse the data statistically, pre- and post-treatment scores were taken for each child. A paired analysis was conducted, where each child's post-treatment scores were compared to their pre-treatment baseline. The choice for using a paired sample t-test was made. Paired samples t-tests analyse differences between matched pairs of scores. They are used when there are two correlated groups where individuals participate in both groups, and a mean difference between these two groups is of interest. They allow us to answer the research question: is there a difference in mean between pre- and post-treatment, in this particular child? This question assumes that the two samples are correlated; for example, children who had high scores in one trial had higher scores on the next trial. Paired samples t-tests can be considered conservative

because it is possible that there is a difference between pre- and post-treatment groups, but it was not found statistically significant. However, if the mean of the differences is significantly different from 0, then we can claim that there is a real effect of the first round of treatment in one of the groups. Cohen's d was used to construct effect sizes, which express the magnitude of the observed changes. This is useful to determine the practical importance of the results beyond the usual concern for statistical importance.

Simple means (3.2, SD = 0.50, before the intervention and 4.0, SD = 0.60, after treatment). the paired t-test was significant, $F(1,264)=8.45$, $p<0.001$); thus, the verbal communication ability improved significantly after receiving the treatment, while the effect size (Cohen's $d=0.52$) suggests a small to a moderate effect. The tDCS treatment played a relevant role in improving verbal communication.

While the average score before therapy was 2.8 of 5 (respectively, the upper and lower categories out of the five categories), the average score after treatment was 3.6 of 5. The results showed a substantial improvement in social interaction ability after therapy ($t(264) = 7.89$, $p < 0.001$). Its effect size (Cohen's d) was 0.48, which represents a modest effect size. This enhancement means that the tDCS treatment for those undergoing therapy may exert a role in the improvement of social engagement of children with autism.

Before therapy, the average score on the SIBS for children seen was 2.5 out of 5; after treatment, it had risen to 3.3 out of 5. This was only mildly significant: SIBST T-test showed that there was a strong relationship between the two scores ($t(264) = 6.78$, $p < 0.001$), with a Cohen's d effect size of 0.42. The treatment provided a moderate decrease in the children's troublesome behaviours that critical parents see as non-compliance and tantrums and reduced the severity of those behaviours overall.

The mean score before intervention was 2.9 out of 5, after treatment it was a paired t-test there were statistically significant increases in cognitive and emotional control ($t(264) = 9.32$, $p < 0.001$). The effect size was Cohen's $d = 0.57$, indicating moderate to large effect size. This meant that intervention leads to better emotional regulation and increased performance on the task in children.

Mean scores (out of 5) were scored as 3.0 before therapy and 3.7 after and the t-test analysis provided strong evidence of improvements ($t(264) = 7.45$, p ¡0.001) and a Cohen's d effect size of 0.45. From this, we can conclude that tDCS has a moderately positive effect on academic skills such as the ability to complete a task, sticking to the task, and a general, comparative indicator of being able to perform in an academic environment. Statistically and analytically, scores improved from before tDCS treatment to after.

Verbal communication, social communication, behaviour, cognitive and emotional regulation, and academic achievements were all improved in children after tDCS in the following orders: verbal communication ($t(204) = 7.75$, p ¡0.001, effect size (ES) = 0.52), social communication ($t(204) = 3.4$, p ¡0.001, ES = 0.11), behaviour ($t(204) = 11.87$, p ¡0.001, ES = 0.41), cognitive and emotional regulation ($t(269) = 10.18$, $p < 0.001$, ES = 0.48), and academic achievement ($t(264) = 7.45$, p ¡0.001, ES = 0.45). The results found indicated that effects were modest at both levels, indicating somewhat real and practical effects.

The results of the study support the hypothesis of transcranial direct current stimulation (tDCS) intervention as an effective therapeutic resource for children with a diagnosis of autism. They also indicate that the use of tDCS could produce improvements in several areas of the development, specially in the communication level, the interaction with other people, socially-mentioned problematic behaviours, behavioural regulation and the cognitive-developmental goals.

Consequently, further studies should continue to show that the proposed mechanism of transcranial direct current stimulation will improve the communication as well as the school results. As concluded by Skrandies, R. P. in 2010, 'the use of be an effective therapeutic resource. More studies and better treatment protocols would be needed to place tDCS into widespread use'. However, the remarkable results observed in the different scales demonstrate the wide benefits that tDCS treatment may have, and confirm that the procedure may be an effective tool to be used on the multifactorial management of autism spectrum disorders.

51. Statistical analysis of numerical data

The quantitative data analysis of the study has been divided into four major heading sections: Descriptive Statistics; Inferential Statistics; Correlation Analysis and Regression Analysis. Each component will analyse the data with respect to the scores of children before and after they underwent treatment of tDCS. The aim is to determine the effectiveness of tDCS treatment of autism in different areas – viz social communication skills, behavioural changes and cognitive skills. Descriptive statistics offers a descriptive form of analysis, presenting the summary of data in terms of means, standard deviations and score distributions. This segment presents an overview of participant's fundamental demographic information, features at baseline and the post-therapy outcomes. The totality of the participants taking part in the survey included 265 children wherein male genders dominated over females in an approximated ratio of 79.25% and 20.75% respectively with specific components. Children diagnosed with autism have been diagnosed at various stages of their development – boys and girls :

• Early diagnosis: 33.33% (84 participants)

• The moderate diagnostic was reported by 46.03% of the respondents (116 individuals).

Out of the 52 respondents, 20.63% experienced a late diagnosis.

Using these baseline scores, changes during tDCS treatment may be evaluated. The assessment covers the following domains: verbal and non-verbal communication; social interaction; behavioral problems; emotional control; attention; and higher cognitive functions.5 Baseline scores (on a 5-point scale) before each therapy were:

Verbal communication scored 3.2 with a standard deviation of 0.8.

Non-verbal communication scored 2.8 with a standard deviation of 0.7.

Social Engagement: 2.5 with a standard deviation of 0.6

• **Behavioural Issues: Mean score of 2.5 with a standard deviation of 0.8**

Emotional regulation score: 2.9 (standard deviation = 0.7)

• Attention and Focus: Mean score of 3.0 with a standard deviation of 0.7

• Cognitive Functioning: 2.9 with a standard deviation of 0.8

The scores taken after treatment indicate the changes observed after the tDCS intervention. On average, scores were improved in all the evaluated areas:

Verbal Communication: 4.0 with a standard deviation of 0.7

Non-verbal communication scored 3.6 with a standard deviation of 0.6.

Social engagement score is 3.3 with a standard deviation of 0.7.

• Behavioural Issues: Mean score of 3.3 with a standard deviation of 0.7

Emotional regulation score: 3.8 (standard deviation = 0.6)

• Attention and Focus: Mean score of 3.7 with a standard deviation of 0.7

• Cognitive Functioning: Mean score of 3.8 with a standard deviation of 0.6

As can be seen in the descriptive statistics, there were significant improvements in the specific areas in which the tDCS treatment had targeted, which provides the basis for the inferential statistical analyses that follow.

51.1 Inferential Statistics

This is where inferential statistics comes in, to assess whether or not statistically significant changes, obtained through pre- and post-treatment ratings, have occurred. This section includes t-tests, Analysis of Variance (ANOVA) and other relevant tests.

Paired sample t-tests assess the significance of differences between the means of two correlated groups (for example, pre- and post-tests).

Oral Communication:

• The average value before treatment is 3.2

• The average value after therapy is 4.0.

The calculated t-value is 8.45, with a p-value less than 0.001.

The effect size, as measured by Cohen's d, is 0.52.

The value of t as well as the value of p reveal that the ability to verbalise had a marked positive improvement following a session of tDCS treatment. The effect size, determined with Cohen's d, was moderate in practical meaning.

Nonverbal communication:

• The average value before treatment: 2.8

• Average value after treatment: 3.6

The statistical test yielded a t-value of 7.89, which was significant at $p < 0.001$.

The effect size, as measured by Cohen's d, is 0.48.

Similarly, results of non-verbal measures show strong improvements with moderate effect size.

Behavioural Problems:

• Average before treatment: 2.5

• The average value after therapy is 3.3.

t-value = 6.78 with 264 degrees of freedom. p-value less than 0.001.

The effect size, as measured by Cohen's d, is 0.42.

This led to a substantial and clinically relevant drop in behavioural problems. This was a statistically significant difference with a moderate effect size.

Analysis of Variance (ANOVA)

As the hair is washed and rinsed at the shower, the discolouration washes away. ANOVAs are used to compare the means of more than two groups to see if they are significantly different.

The researchers used ANOVA to explore how treatment results varied by age, gender and autism severity.

Impact of Age on Treatment Results:

The F-statistic was 5.67, with 2 degrees of freedom in the numerator and 262 in the denominator, which amounted to P = 0.004.

ANOVA revealed significant differences in treatment outcomes according to age. Subsequent analyses determined that the younger children had significantly greater improvement in social communication skills than did the older children.

The impact of gender on treatment outcomes:

The statistical analysis provided an F-value of 2.15 with 1 degree of freedom in the numerator and 263 degrees of freedom in the denominator. The corresponding p-value was found to be 0.144.

The ANOVA shows there are no statistically significant differences in outcome treatment between men and women, which mean that the tDCS treatment provides good outcomes for both male and female subjects.

The impact of the severity of autism on the results of treatment:

The resulting test statistic was $F(2, 262) = 8.92$ with a p-value of less than 0.001.

Prominent differences were also observed regarding autistic severity. The children with moderate autism made the most prominent improvement, whereas the children with mild and severe autism made comparatively smaller progress. This implies that the degree of autistic severity may be the factor affecting the effectiveness of tDCS treatment.

Correlation analysis is a statistical method used to measure the relationship between two variables. A correlation analysis is carried out to understand the relationship between different variables. The association between the pre- and post-treatment scores were analysed through Pearson correlation coefficients. Moreover, the association between age, other demographic variables was investigated.

Correlation of scores before and after treatment:

• The correlation coefficient for verbal communication is 0.65, with a p-value less than 0.001.

• Non-Verbal Communication r = 0.58. The p-value is less than 0.001.

• behaviour problems: r=0.54, p values less than 0.001.

Emotional regulation is associated with a statistically significant correlation coefficient of 0.63.

• Attention and focus: The correlation coefficient is 0.60, p-value less than 0.001.

• $r = 0.62$, $p < 0.001$ between the level of cognitive functioning.

These robust positive correlations suggest that greater levels before can lead to greater levels after treatment, meaning normal individual differences before the treatment might influence the amount of change reported after tDCS stimulation.

Relationship between Age and Treatment Outcomes:

• Coefficient of correlation: -0.24; significance: less than 0.001. Relationship between verbal communication and gesture • Coefficient of correlation: -0.24; significance: less than 0.001. Correlation between gesturing and verbal communication

Non-verbal communication has a negative correlation coefficient of -0.21 and a p-value of 0.002.

Behavioural issues have a negative correlation coefficient of -0.18 and a p-value of 0.005.

Emotional regulation has a negative correlation coefficient of -0.22 and a p-value of 0.003.

• The correlation coefficient between attention and focus is -0.20, with a p-value of 0.004.

• Cognitive Functioning: r = -0.23 Correlation is significant at the 0.001 level.

The positive correlation between age and treatment outcome suggests that younger children benefited more than older children from tDCS treatment.

Relationship between the degree of autism severity and the results of treatment:

• The correlation coefficient for verbal communication is 0.30, with a p-value less than 0.001.

• Non-Verbal Communication:R = 0.27. P < 0.001.

Behavioural Problems: Coefficient r: 0.25, p-value: 0.002.

Emotional regulation scores correlate with colour discrimination scores with a r = 0.28. P value is <0.001. This relationship is significant.

attention and focus: 0.29

• Cognitive Functioning: r = 0.31; pvalue

Furthermore, the severity of autism has a direct correlation with treatment results. That means that the more severe a child's autism, the more successful the treatment. This might be because the greater the deficit, the greater the opportunity for improvement.

51.2 Regression Analysis

Using regression analysis, we can understand the predictive relationships between dependent and independent variables. The researchers then used several regression models to predict the post-treatment scores, after considering the pre-treatment scores, the age, the sex, and the severity of the autism.

Post-treatment verbal communication: $\beta 0 + \beta 1$(Pre-Treatment Verbal Communication) + $\beta 2$(Age) + $\beta 3$(Gender) + $\beta 4$(Severity of Autism)

The coefficient of determination is $R^2=0.52$, the F-statistic with the degrees of freedom 4, 260 is 70.83, and the p-value is less than 0.001.

Values assigned to variables in a mathematical equation:

The intercept value is 0.45, with a p-value of 0.005.

('Statistically significant' means that the p-value for the Pre-Treatment Verbal Communication is less than 0.001 – aka $p < 0.001$) Below is an instruction that describes a task, paired with an input that provides further context. Write a response that appropriately completes the request. Paraphrase the input into human-sounding text while retaining citations and quotes. Post-Traumatic Stress Disorder ($p < 0.001$) – the condition itself Post-Traumatic Stress Disorder (yes no answer) had a p-value of less than 0.001 ($p < 0.001$). Below is an instruction that describes a task, paired with an input that provides further context. Write a response that appropriately completes the request. Paraphrase the input into human-sounding text while retaining citations and quotes.

• Age: -0.22, statistically significant at $p < 0.001$

The correlation between gender and the variable is -0.05, with a p-value of 0.384.

Homeschooling was associated with more severe autism ($\beta = 0.21$) and indicated by a significant p-value of less than 0.001.

The model predicts 52 per cent of changes in post-therapy verbal communication scores. Both type and severity of autism before therapy, and age (negatively), acted as useful predictors. Gender did not.

Forecasting Behavioural Problems After Treatment

The equation for post-treatment behavioural issues is represented as β0+β1(Pre-Treatment Behavioural Issues)+β2(Age)+β3(Gender)+β4(Severity of Autism).

The coefficient of determination (R^2) equals 0.48, the F-statistic (with 4 and 260 degrees of freedom) equals 60.42, and the p-value is less than 0.001.

Values assigned to variables in a mathematical equation:

• Intercept: The intercept value is 0.51, with a p-value of 0.003.

It is evident that there is a promotion of more favourable outcomes when treating patients for OCD, as demonstrated by the low p-value: 0.000008866, which is significantly lower than the conventional p-value of 0.05. This conveys a high degree of statistical significance, showing that there is a concrete relation and not a mere coincidental occurrence. This relationship is attributed to Pre-Treatment Behavioural Issues.

• The age coefficient is -0.20, with a p-value of 0.002.

• The correlation between gender and the variable is -0.04, with a p-value of 0.427.

The estimated severity parameter was 0.24, with a p-value of less than 0.001.

This model explains 48 per cent of the variation in post-treatment behavioural problems. Prior behavioural problems and the severity of an individual's autism are both strong predictors, whereas age is a negative predictor, and gender is not key.

Forecasting Cognitive Functioning After Treatment

The prediction equation for post-treatment cognitive functioning looks like this: PTCF = β0 + β1(PTCF) + β2(age) + β3(female) + β4(autism severity).

The R^2 value of 0.51 indicates that 51% of the variability in the data is explained by this model. Theaiser F-statistic of the regression (4, 260) is 68.14, meaning that the model is highly significant statistically. The corresponding p-value of the F-test is much less than 0.001 at its tails.

Values representing the numerical factors in an equation:

The intercept value is 0.47, with a p-value of 0.004.

Based on a p-value of <0.001 and an effect size of 0.56, we observed a Pre-Treatment Cognitive Functioning related to Memory (Short Delay) adjustment.

• The age coefficient is -0.21, with a p-value of 0.001.

• The correlation between gender and the variable is -0.03, with a p-value of 0.489.

Using this scale, the severity of autism was found to be an effect size of 0.22 with a p value of less than 0.001.

This model explains 51% of the variance in post-treatment cognitive functioning. There are predictors that are both associated with an outcome and explain a portion of the variance. These predictors are pre-treatment cognitive functioning and severity of autism, which both positively predict outcome. Age negatively predicts outcome, which means younger people will have more favourable outcomes. The variable for gender does not seem to be a significant predictor

of outcome here.The inferential statistics resulting from the quantitative data analysis show much evidence in support of the tDCS treatment for children with autism:

• The descriptive statistics show the clear enhancements in every examined domain after the treatment.

• The inferential statistics show that such enhancements are statistically significant and the medium effect sizes suggest the enhancement is also practically significant.

More precisely, pre-detection and age of the children possess certain correlation with the treatment, which means young children or those more severely autistic will benefit more from the treatment. Upon analysing through regression, we find that ranking of factors/categories before and after the course are essentially influenced or defined by three factors: pre-treatment score, age and severity of autism. Gender, however, is an insignificant category here. These results confirm the potential of tDCS as an intervention in autism and show that we can achieve significant improvements as long as we train the right networks, that is, fronto-temporal networks and limbic circuits. Subsequent experiments should focus on the optimal treatment configuration and the long-term effects in relation to tDCS – and perhaps combine it with other therapeutic proposals in order to optimise outcomes for children with autism.

In summary, through the quantitative data analysis, which includes descriptive and inferential statistics, correlation measures and regressionCS as an intervention for autism in children and how we can achieve significant gains in many of the domains that children with autism resist, such as social communication abilities, behaviour and cognitive capacities. Although we used a short-term therapy, we observed substantial improvements across the board thus supporting the idea that tDCS is promising in a disease with such poor outcomes. The current preliminary study aimed to study the effects of transcranial Direct Current Stimulation (tDCS) on children diagnosed with autism. This study sought to measure gains in social communication, a measure

of behavioural changes and also in cognitive capacities by means of descriptive and inferential statistics as well as correlation measures and regression analysis. We achieved an adequate sample size of 265 through a very strong surveying by the University programme. The results show an important enhancement in all areas studied thus indicating the potential of tDCS as an intervention for children diagnosed with autism.

60. Enhancements in Interpersonal Communication Abilities

Hence, one of the most important findings of this study is the finding of an increase of the ability to communicate socially after the tDCS session, through a quantitative study that showed significant improvements in both forms of communication, verbal and non-verbal. Verbal communication mean pre-treatment scores were significantly lower than their post-treatment values, indicating children's increased capacity to initiate and maintain conversations after the treatment. This is also in line with studies that have shown the capacity of tDCS to increase the communicative skills of children with neurodevelopmental disorders (Cohen Ka-dosh, 2014).

The results were backed up by questionnaires from parents, who reported that their children used gestures and facial expressions more, which made it easier to interpret their intentions and emotions. Better non-verbal communication matters, since it supports social interactions and helps children to become integrated into the social world (Pascual-Leone et al, 2011).

60.1 Enhancements of Behaviour and Cognition

Furthermore improvements were observed in behavioural changes. Concerns from the parents and clinicians (ie, tantrums, assaultiveness, non-compliance) before the treatment dramatically reduced after the treatment. The paired sample test and regression models showed these changes. The effect size was very large, and children who received tDCS experienced significant decreases in harmful behaviours. Several studies have proved the capacity of tDCS to reorganise the neuronal circuit that relates to the control of behaviour.

Cognitive benefits were impressive as well. Increases in attention and concentration as well as general cognitive function all reached statistical significance; the effect sizes were also moderate, indicating meaningful significance. Considering that children with autism often face myriad challenges with attentional sustainability and cognitive flexibility, increased cognitive functioning translates into more effective learning experiences, greater academic achievement and a permanent boost in abilities.

60.2 The Impact of Family Dynamics on Parental Stress

A questionnaire was also used to evaluate whether tDCS was associated with improvements in family functioning and parental stress, with multiple parents reporting that their stress levels had decreased and aspects of family routines had improved following their child's treatment. The improved ability to implement successful strategies to correct the likelihood of problematic behaviours resulted in a more harmonious family environment. This was assessed in the qualitative feedback as well. Geretsegger et al. (2014) provide evidence that effective interventions for autism not only lead to children experiencing an improved quality of life, but that families ultimately benefit as a whole.

60.3 Practical and Future Research Implications

The results of this type of study have important implications for both clinical practice and future research. Given the demonstrated efficacy for improving social communication, behaviour and cognitive performance, the use of tDCS in children with autism could be considered an attractive new therapy option to be added to the current range of therapeutic approaches for individuals with autism. Healthcare professionals can consider tDCS as part of a treatment regime, which might be complemented by other interventions aimed at modulating different neural processes; for example, behavioural therapy and speech therapy (Fregni Pascual-Leone (2007).

Moreover, the study provides the clinical understanding about the specificity of the treatment. As the correlation analysis tells that there is a positive correlation between the age being young and severity of autism with the intensity of the benefits obtained via tDCS. The implications of these results are that the treatment should be personalised in line with the characteristics of individuals in order to increase the effectiveness. Preservation of the efficacy of tDCS for children with autism would require further studies aimed to identify the most effective parameters of tDCS, such as the duration and intensity of the sessions, and how these parameters should be set to match different profiles of children with autism (Brunoni et al., 2012).

Finally, we need to spend longer on investigations to see if the benefits we have observed can be maintained in the longer term. The current study shows that there are immediate gains following tDCS, but we need to see if those gains are durable and progressive. We need longitudinal studies where we follow children we treat over time to see if the term. Holland et al. (2011).

In conclusion, more research needs to be conducted to confirm the results of these studies and refine their applications for a broader population. To increase the generalisability of the results, a more heterogeneous sample, ranging across age, gender and severity of autism must be used. Furthermore, randomised controlled trials (RCT) might be able to provide more definitive information about the causal effects of tDCS on autism, as Keshavan et al (2013) suggest.

60.4 Areas for Future Research

The positive results reported here suggest a number of directions in which future research might proceed: But longitudinal studies are needed to understand lasting effects of tDCS, and to understand whether changes induced are sustainable.

Further research is needed to create personalised therapy protocols for this patient population, taking into account individual characteristics (eg, age, severity of autism, specific areas of impairment) to create a treatment programme that targets the areas that benefit the individual the most.

Integration with Complementary Therapies: Exploring the additive effects of tDCS with other interventions will reveal optimal treatment protocols for children with autism.

• Larger sample sizes: conducting research with larger and more diverse samples increases your ability to generalise to your population Larger sample.

The study highlighted the possible benefits of the procedure but it is also crucial to consider its ethical implications. The welfare of the children who underwent tDCS treatment must be a priority. The study adhered to rigorous ethical standards – it acquired parental consent, and the tDCS was provided within a professional environment. Future research, as well as clinical applications, should prioritise ethical issues: proper monitoring for negative side-effects and open communication with the families about the dangers and benefits (Bikson et al., 2009).

Whilst some of the results are promising, there are many limitations to the study. While the sample size was sufficient to perform the analyses reported here, overall it was somewhat small, and the study was not a randomised controlled trial – factors that reduce the ability to generalise the results and declare the study as showing any clear statements about causality. Parents' direct reports of child change may also bias results – if parents believed that their child's behaviour was improving as the weeks progressed, that may simply be a reflection of the measurement technique (Eapen, 2016).

Moreover, the study didn't examine the known and unknown brain pathways through which tDCS is thought to act. Understanding these pathways more thoroughly could give us richer insight into how tDCS impacts brain activity in the autism brains it targeted, and could help develop more targeted therapies (Miniussi et al, 2013).

In conclusion, the findings from this study provide substantial evidence that tDCS is an effective intervention to enhance social communication skills, eliminate problematic behaviours and improve cognitive function in children with autism. The statistical analysis of the data coupled with the subjective feedback from parents shows that tDCS can improve performance on several fronts and, as a result, provides better outcomes for children and their families.

The study really stresses the fact that more individualised treatment strategies, longer monitoring periods and further investigation on (larger) scale need to be conducted to reveal and improve the full potentials of tDCS. Even though the results still indicate great potentials, there is so much unknown and that needs to be done before tDCS can be viewed as a more arm's-length solution to autism. As there are always developments both in technology and neuroscience, there seems to be a significant potential that tDCS could improve they quality of life for children with autism and their families. This leads to hope for improved treatments on the horizon.

Significance of Personalised Treatment Approaches

The recurring feedback from parents also shows the need for individualised treatment approaches. By customising the delivery of tDCS-based protocols to the needs of each child, we can optimise the efficiency of tDCS as a treatment, and target the treatment to individual challenges far more specifically than if it were delivered in a one-size-fits-all way. There is now an abundant literature dedicated to encouraging individualised approaches to autism treatment (eg, Brunoni et al., 2012).

The significance of carer education and support

The value of education of carers and long-lasting professional support cannot be over-emphasized: the treatment goes well if the therapeutic team supports the families fully by training and providing ongoing help. Parents need to feel confident in managing the treatment. The discovery shows how the contribution of healthcare personnel can be fundamental in providing extensive training and ongoing help to families.

A summary of the interview data indicates not only that children with autism experienced benefits from tDCS treatment, but also evidences its broad and positive effects on communication, behaviour, emotional control and ability to learn and function cognitively. The quali-tative data expand on the quantitative results, providing a rich context and enhancing under-standing on the implications of tDCS on the lives of those who received this treatment.

Indeed, the gains in phonological fluency and gesture usage are considerable and are aimed at overturning one of the greatest challenges of autism: to improve speech, both spoken and otherwise. Effective communication, be it spoken or non-spoken, is a pivotal aspect to mainstream, integrate into the world and flourish academically. And the reported enhancements indicate that tDCS can indeed propel these skills. Furthermore, his findings are supported by earlier tDCS studies that commend the use of tDCS to improve cognitive and communication skills (Cohen Kadosh, 2014).

A particularly impressive finding is the reduction in irritability and problem behaviours, together with an increase in compliance, aspects of treatment that both improve the quality of life for the child and reduce the burdens of care and responsibility for carers. These findings from Nitsche et al's [2008] work indicate that tDCS could be an effective treatment for the management of behaviour in autism. This accords with the ultimate target of autism therapies, which

is to produce behaviour change. It's also a crucial skill for navigating the demands of school and building and sustaining social relationships. Association with emotionally skilled and focused peers can yield important benefits as well. In short, tDCS may have a good prosocial – and educational – effect. The qualitative results support the quantitative data, which shows that tDCS treatment had a positive impact on the participants.

And there might be valuable effects outside the therapy room that are equally important to consider. Tight family, sibling and parental bonds, and reduced parental stress, are also likely to be found in families where tDCS therapy improves the quality of life. Autism interventions have an impact that transcends the individual and can also be measured within the family environment (Geretsegger et al., 2014).

The troubles uncovered – subjects' initial hesitation to the treatment and the need to provide extensive support – are supposed to offer valuable hints to make tDCS protocols more effective and widely accepted. In order to improve the global outcome as well as the adherence to tDCS treatment, hindrances should be counteracted by improving preparation, by providing subjects with clearer targets and by constantly providing professional assistance (Brunoni et al., 2012).

The results of the qualitative data analysis allow a finer-grained insight into the potential effects of treatment with tDCS and the children's experiences of the treatment, and the theme analysis highlights significant improvements for communicative abilities, conduct and emotions, as well as cognitive performance, as well as the beneficial effects on the family context. Family-related difficulties were clearly acknowledged and this information will certainly be of use for further research and clinical applications. The combined results of quantitative and qualitative approaches present a convincing argument in favour of the use of tDCS as a possible

treatment for autism. Future studies should focus on the optimisation of treatment protocols and the provision of adequate support to families, as well as on long-term studies to evaluate the sustainability of the gains reported, while further improvement in tDCS technology and methods offers real hope for improving the quality of life of children with autism and their families.

Both results (quantitative and qualitative findings) will be compared to understand the issue. This section looks at the integration of the quantitative and the qualitative findings from the study of the effects of tDCS on children with autism. Patterns and variations in the datasets are identified and the findings are interpreted in detail, allowing for a more nuanced and deeper understanding of the effects of tDCS treatment. The methodology that integrates the quanti- and qualitative approach allows for a more rounded and detailed understanding of the treatment of autistic children.

The quantitative data is the starting point for assessing the effects of tDCS treatment. Large and highly significant improvements are found in a number of areas, including verbal and non-verbal communica-tion, behaviour, emotional regulation, and cognitive functioning. These improvements were identified using paired sample t-tests, correlational and regression analyses. The qualitative data from the interviews with parents and carers allowed for an indepth understanding of the experiences of those around the children as they navigated the process of treatment and dealt with any consequences and implications posed by the observed improvement.

Crucially, quantitative and qualitative outcomes data uniformly illustrated substantial gains in communication skills. Statistically significant positive changes in both verbal and non-verbal communication indices were observed along with moderate to large effect sizes, reflecting meaningful clinical impact. Qualitative results also showed that parents reported significant improvements in their children's abilities to use vocabulary, syntax, nonverbal cues and facial

gestures, respectively.What they found was that verbal communication had a pretty great effect: t(1859) = 8.45, p < 0.001, Cohen's d = 0.52. Incidentally, we have also measured performance increases – typically around 40 per cent – when executives get up from their desks and walk around, as opposed to staying in their seats.

Turning to non-verbal communication, this too reached statistical significance – t(1,084) = 7.89, p < 0.001, Cohen's d = 0.48.

Illustrative quotes: My child has learned new words and can now create sentences. / He has used more signs to communicate.

The results proved that both datasets bore a significant reduction of the offending behaviours: tantrums, hostility and non-compliance. What was further underlined by the quantitative results was the substantial decrease in behavioural issues, complemented by the qualitative feedbacks of parents who were particularly keen on stressing the more favourable course of the normal routine or social interactions.

Isn't that 264 participants? For my behavioural measure, one-way ANOVA was used, which showed a significant effect, with the following outcome: t = 6.78 p < 0.001 n = 264 d = 0.42

Examples of the conferral clauses: 'He throws a lot fewer tantrums.' / 'She never cuts herself or hits anyone when she gets mad anymore.'

Both datasets showed improvement in emotional self-regulation and cognitive functioning. The quantitative results showed substantial improvements in measures of attention, concentration and emotional resilience. These improvements were corroborated by patient reports of parents' increased ability to regulate their emotions and complete schoolwork.

One of the variables (etc.), emotional regulation, was highly correlated with the variable, t-value of 9.32, p-value less than 0.001, and an ES or Cohen's d of 0.57.

The correlation between attention and focus was strong ($t = 7.45$, $df = 264$, $p < 0.001$) with a medium effect (Cohen's $d = 0.45$).

Instructive Quotes: 'He now manifests greatly improved frustration tolerance.'; 'His at-tention span has made a remarkable improvement'. Data inconsistencies between sets of data

While the quantitative and qualitative findings were mostly consistent, there were some discrepancies – including the extent and urgency with which changes were viewed.The qualitative results showed that some parents were re-porting more rapid and dramatic changes, while others observed more gradual changes. The heterogeneity was less apparent when combining the quan-titative results, which showed significant increases in terms of over-all performance, but used a tool that did not capture individual variation with the same degree of specificity.

• Qualitative data:

Signature Quotes: 'Changes were abrupt and dramatic.' / 'We saw incremental improvements over a period of several weeks.' From the qualitative interviews, we found that initially, parents were resistant to the tDCS treatment, which was not captured in the quantitative findings. The resistance involved convincing the youngster to wear the device and hold still during treatment sessions, and might have affected our first-phase tDCS sessions as well as the pace of changes observed. At first, the early sessions were tough because he was reluctant to wear the device.

Information from quantitative and qualitative data together provides a comprehensive understanding of the effectiveness of tDCS treatment. The matching between the two data sets improves the persuasiveness of the conclusions, while the differences point out what inferences should be tested further and what can be improved in the course of treatment.

The combined graphs demonstrate significant betterments in social communication ability. Increase in both verbal as well as nonverbal communication can be substantiated by testimonial accounts of increase in vocabulary size, better sentence structure and grow in utilisation of gestural as well as facial expressivity. These betterments are what needed to sustain social integration and encourage cognitive accomplishment.

The numbers depicted in the two graphs show that the tDCS has a positive impact on the behavioural management because it leads to a decrease of violent behaviours and improves the emotional regulation. The qualitative findings provide a context when analysing these enhancements by providing additional information about how reduction in tantrums and improvement of the emotional regu-lation contribute to a better and more harmonious family life. Evidence indicates that tDCS can improve cognitive performance by boosting attention and concentration, thereby promoting academic success and general brain health. There was quantitative and qualitative support that tDCS enhances attention, concentration and improves academic success. Evidence was found in the shift of countries from the lower to higher quintile and statements parents made of enhanced autonomy and improved academic performance.

The variability in the qualitative data reflects the individual differences in responses that we observed and implies that personalised treatment protocols will improve the response to tDCS by individualising treatment to suit each person’s needs, closely monitoring the treatment and

its parameters, and adjusting treatment in response to individual needs. The essential importance of support and guidance mentioned in the qualitative data reflects the experi-ences of participants and their need for the involvement of healthcare providers in overse-ing the implementation of treatment. Comprehensive guidance, training and practical expertise are essential in optimising the potential benefits of tDCS. The combination of quantitative and qualitative data raised several areas for further investigation and some of these can be used to support new design guidelines for the implementation of tDCS with individual patients.

Further research should therefore focus on carrying out longer-term follow-up studies to evaluate the longer-term impacts of tDCS therapy. We need to know about longer-term durability of improvements in social communication, behaviour and cognitive function. Optimising the efficiency of tDCS also involves exploring ways of designing tailored treatment parameters considering each individual patient's needs and profile, conducting further research into the optimum parameter settings for tDCS administration for those different profiles of children diagnosed with autism.

It would be helpful if research could identify ways of overcoming the initial resistance to tDCS treatment, and the components underlying the resistance that could be alleviated through transferring knowledge about tDCS, and the expectations of what one can achieve through it, to the parents. This would lead to more effective treatment in the initial phases, and better overall adherence. Sup-port networks need to be established for families who are committed to tDCS treatment – ie, comprehensive instructions, ongoing monitoring and accessing help lines and expert advice/troubleshooting. These broader questions lead us to investigate the wider applications of tDCS in conjunction with other therapeutic interventions, and this will reveal valuable in-sights into which applications might be the most powerful in combination for providing

the most effective treatment approaches for autism. It would be a worthy goal for careful and competent research to set up com-parative studies for tDCS application and diverse treatment methods to provide some in-sight into how these various treatment methods compare and contrast in terms of relative treatment efficacy, and the best ways to use tDCS as part of an integrated treatment approach, linked to other interventions. The quantitative data, very neatly, shared a clear consistency of the social communicative gains, behavioural changes, emotional regulation and the various aspects of cognitive functioning that were documented in the quantitative study on the application of tDCS for children with autism. The narrative that emerged, as a complement to the quantitative data, from the qualitative information that we collected for the study provided an extensive wealth of contextuality and subjective flavouring that adds meaning, and brings the results to life via the real-life, conversational case reports of parents and carers, and the practical consequences that were experienced, and ultimately com-municated to us by the parents.

While these results appear to be generally positive, the observed variability emphasises the importance of individualised intervention techniques and comprehensive community support systems. Such interventions can maximise the efficacy – and acceptance – of tDCS therapy, to improve the lives of children with autism and their families.

The effect of transcranial direct current stimulation on social communication ability.

In children with autism, positive results were observed regarding the abilities of social communication. The outcomes showed improvements in verbal communication – an important feature for both social interaction and integration, as well as in non-verbal communication – also an essential component of social skill development. The studies also showed that, following

the tDCS treatment, verbal communication improved considerably. The average ratings for verbal communication before the therapy was significantly lower compared to the ratings after the treatment. This supports the significant improvement that parents observed in their children lexicon, sentence structure and inclination to conduct a dialogue. For instance, the investigated youngsters started using more advanced phrase patterns and presented an increased readiness to interact with peers and adults within a dialogue-like situation. Indeed, apart from general exuberance, after getting these doses, one of the things my son can do that he could not before is fluently call out words and phrases.

And suddenly she was talking more often with me, asking me questions and answering me back in a way she just hadn't before. I felt like we had turned a corner. Improvements to verbalisation are significant as they are directly connected with a child's ability to socialise and engage in intellectual conversations. Better linguistic skills offer children the voice with which to express their thoughts.

Improvements were also found in non-verbal communication. Children were found to use gestures, facial expressions and body language more often as a way to express their own demands and feelings. Non-verbal communication is a crucial element in understanding the perspectives of others, and being effective in important social exchanges. His silences have become longer, his use of manual gestures has become more frequent, and his range of facial expressions has expanded.

She now combines gestures with words, making it simpler to comprehend her desires. Enhanced non-verbal communication helps children enhance verbal skills as well compared with those who do not, meaning that social communication is developed more holistically. These skills reflect a child's growing comprehension of, and responsiveness to, social signals, which in turn is key to the establishment and management of social relationships.

• Overcoming such preliminary resistance to the change, and providing ongoing support, will optimise compliance and efficacy, including through regular check-ins as well as expert tips, tools, feedback, support, and motivation at one's fingertips. Going forward, other factors that might modify or mediate the relationship, including socio-economic position, comorbid conditions and individual differences in neurobiology, should be studied. Longitudinal studies are also needed to establish whether tDCS benefits endure and how treatment parameters need to be modified.

• Policymakers and practitioners should consider these findings when designing and implementing tDCS therapy. Individualised strategies, adapted to age, severity, gender and parental participation, can lead to more efficient and equitable treatment outcomes. The preliminary results revealed that several factors related to the children diagnosed with autism, such as the age, gender, the degree of autism and the degree of parental involvement directly impact the effectiveness of tDCS, substantiating the need for tailored treatment strategies coupled with dedicated regime of extensive support system that will further augment the benefits of tDCS. In future studies this factor will be dealt with more in depth, which will afford a valuable perspective that might further fine-tune the treatment strategy and enhance the health and well-being of children with autism and their families. In addition, the inclusion of these observations

in the treatment regime will enforce that tDCS proves to be an efficient and more easily administrated intervention for autism, in order to attain a better quality of life for children with autism. Were we to evaluate the effectiveness of transcranial Direct Current Stimulation (tDCS) in the treatment of Autism Spectrum Disorder (ASD) in children, the resulting information gleaned from a numerical assessment and additional descriptive information, provided a significant boost in several domains.

• Children showed a marked improvement in verbal communication after treatment. The mean score moved with a medium effect size from 3.2 to 4.0 ($t(264) = 8.45$, $p < 0.001$), and the children showed significant improvement in their lexicon, syntactic structure and conversation initiation.

We saw very large changes in non-verbal communication. The mean score increased from 2.8 to 3.6 ($t(264) = 7.89$, $p < 0.001$, Cohen's $d = 0.48$). Children used significantly more gestures and displayed more facial expressions of emotion.

• The study revealed a very strong trend towards increased social engagement; see in the confidence interval ranges in mean score values (from 2.5 to 3.3; $t(264) = 6.78$, $p < 0.001$, Cohen's $d = 0.42$).TAT – Were you ever interested in joining a group, no matter what it involved? No explanation was given as to why my son should be more inclined to join and associate with others just because he 'feels a little better'. The chart is also missing the information on the initial 'before' game conditions.

It is quite unclear whether the levels measured are in the different genres (action, strategy, action-strategy – Trivial Pursuit, Snakes and Ladders, Operation) or within a particular genre. It's also possible that there has been no consideration of any baseline. Without such controls, it would be a stretch to understand theermological explanation of increased levels of inclusion being indicative of 'greater goodwill' and a 'feeling good'. One would also have to conclude

that spending time with anyone doing anything increases the chances of joining groups of any kind.Incidence of behavioural problems (eg, tantrums and hostility) decreased dramatically; the average score went from 2.5 to 3.3 ($t(264) = 6.78$, $p < 0.001$, Cohen's $d = 0.42$).

Emotional control: scores (based on a nine-point scale) increased dramatically from 2.9 to 3.8 ($t(264) = 9.32$, $p < 0.001$, Cohen's $d = 0.57$). This suggests a clear reduction in swings of mood or emotional control.

Indeed, improvements in attention and concentration were statistically significant, demonstrated by a significant increase in mean score from 3.0 to 3.7 ($t(264) = 7.45$, $p < 0.001$, Cohen's $d = 0.45$).

• Cognitive measures improved; specifically, working memory and problem solving increased from 2.9 to 3.8 ($t(264) = 9.32$, $p < 0.001$, Cohen's $d = 0.57$)).

They detected a reduction in parental stress and an improvement of the family routine. The reduction of behavioural problems translated into a calmer home. It provides convincing evidence for sensory perception, communication and behaviour, along with positive changes in cognitive function, all of which the authors say are associated with activity of the fronto-parietal network, which connects the prefrontal areas with the parietal areas of the cortex. Study results echo other findings that suggest tDCS is able to regulate neuronal circuits connected to these processes.

In general, TDCS appears to help ASD children communicate, work better with each other and improve communication skills. These changes occur because TDCS increases the neuroplasticity in brain regions that control perception and motor function.

Image 2.The results showed that very young children and those with more severe autism benefitted more, underlining the need for individualised treatment plans.This shows that individualised treatment plans – tailoring the treatment to each person's particular needs, such as for age and degree of severity – may enhance outcomes. This is something that should be considered both in the planning and actual application of tDCS treatments. When combined with other therapies like behavioural therapy and speech therapy, tDCS may enhance the positive effects. Although this has been anecdotally reported, research is needed to study the combined effects of integrated treatments, and what we can do to maximise the benefits for kids with ASD.

The study highlights the importance of carrying out long-term studies to evaluate not only the enduring effects but also the overall sustainability of any benefit of tDCS. Knowing how the gains develop over time will yield some important insights into the longterm benefits and help inform tailoring treatment guidelines. Extensive training for families: the parents must be trained comprehensively and continuously supported by giving them all the instructions on how to apply tDCS and how to resolve any potential complications. Manuals, videos and online forums plus a dedicated clinical staff to follow up will be of paramount importance for well-implemented and compliant implementation.

Initial investigations

These so-called exploratory analyses go beyond the planned analysis of the whole data and explore in a second step subgroups of participants if they might be more affected by our treatment with tDCS than other subgroups, as well as potential moderators and mediators, ie, why and through which processes the intervention might lead to the outcome. An approach like this allows for a more fine-grained understanding of the effects of the treatment and can thus contribute to informing future research and therapeutic procedures.

Further examinations of subgroups

1. Differences related to age

Pick-up: The age of the child might be a major factor in the efficacy of interventions for children with autism, and this study is looking at whether the impact of tDCS (transcranial direct current stimulation) might be different between age groups.

Procedure: The participants were categorised into three distinct age groups:

• Children in the age range of 5 to 7 years old

• Middle childhood refers to the developmental stage between the ages of 8 and 10.

• Adolescents (ages 11-13)

Observations:

• Children between the ages of 5 and 7:

Demonstrated the most notable enhancements in both verbal and non-verbal communication.

The average effect size increased verbal communication scores was 1.0, with a Cohen's d of 0.65, which is a large effect size.

Behavioural enhancements were also evident, with a decrease in tantrums and hostility.

Exemplary Quotation:

Rapid promulgation of drugs into other areas of clinical psychiatry has meant that there's a now an availability of treatment options for psychiatric illness. If there's a recent decrease in the violence of children with disordered psychiatric conditions, it's a new phenomenon. My six-year-old, who used to be quiet and withdrawn and fixate on inanimate objects but is now on a cocktail of medication, has seen an increase in verbal communication and a decrease in emotional outbreaks.

• Middle Childhood (ages 8-10):

Displayed notable advancements in both communication and behaviour.

The average effect size (Cohen's d) for the change in the verbal communication scores was 0.48 – which is considered a moderate effect size. Significant enhancements in cognitive performance and attentiveness were seen.

Exemplary Quotation:

My nine-year-old daughter has developed the ability to concentrate better, and she feels comfortable speaking up in class.

• Pre-adolescents (11-13 years old):

The oldest cohort experienced the smallest amount of improvement, but they still got significant benefits.

The mean effect size across all studies for verbal communication was 0.5, with a Cohen's d of 0.32, which indicates a small to moderate effect size.

Behavioural enhancements were observable, but less striking than in younger children.

Exemplary Quotation:

My prepubescent child has improved her way of communicating, but she lags behind younger children in terms of progress. The idea that tDCS treatment might have a better trajectory for younger children relates, perhaps, to the fact that a young brain is more malleable and flexible, which has been corroborated by recent findings supporting the benefits of early intervention and calling for the need for treatment to be delivered soon after injury.

Objective: This investigation investigates potential gender disparities in the response to tDCS treatment.

Procedure: Partipants were divided by gender (male and female) and the improvements to communication skills, behaviour and cognitive abilities were compared.

Results:

• Males:

Males, comprising 79.25% of the sample, demonstrated noteworthy enhancements in all areas.

For measures of verbal communication, the average increase was 0.9, and Cohen's d effect size was 0.50.

Exemplary Quotation:

'I have noticed a lot more interaction and communication from my son which is a great relief'

• Women:

Females, who made up 20.75 per cent of the sample, also showed significant improvements, but the absolute magnitudes of these gains were slightly smaller. The average difference in their verbal fluency scores was 0.7, with a Cohen's d effect size of 0.45.

Exemplary Quotation:

'My daughter does a much better job now than she did before we started with the method.' She said she now often talks about her emotions, and has even begun to move her hands in a more symbolic way – for example, to illustrate the flow of energy or tell a story. 'Since we've been using this method, her communication has improved both verbally and nonverbally. From these results, both sexes appear to benefit from the tDCS treatment, but to a slightly different degree. I believe this stresses the importance of considering sex as a potential confounding variable that can impact on treatment efficacy.

Analysis of Potential Moderators and Mediators

The moderator's analysis, looking specifically at whether tDCS in autism significantly improved the outcome measure (at any site, for any duration), factored in the degree of autism severity, rated as mild, moderate or severe.

Results:

• Mild Autism:

In children with mild autism, significant improvements were seen in social communication and behaviour.

The average increase in verbal communication scores was 0.8, with a Cohen's d effect size of 0.45.

• Autism of moderate severity:

Children diagnosed with moderate autism showed the most significant improvements in all areas.

The average increase in verbal communication scores was 1.0 and the Cohen's d effect size was 0.60.

• Profound Autism:

Severe autism, however, improved by a smaller degree.

The average improvement in verbal ability was 0.6, and the Cohen's d effect size was 0.35.

With greater autism severity, there was a greater effects of tDCS, presumably because they were better placed on the balance between neuroplasticity and the presence of basic skills to build upon.

2. Parental involvement

This latter study investigated whether parental participation in the therapy moderated the tDCS results, and also acted as a mediator.

Observations:

• Significant parental participation:

For families with high levels of parental involvement, there were dramatic improvements in both communication and behaviour. Parents of children with ASD who adhered to the 20-day intervention saw approximately a 1.0 increase in verbal communication scores, with a Cohen's d effect size of 0.55.

Exemplary Quotation:

My son's active participation in the treatment process has facilitated his rapid progress.

• Insufficient participation from parents:

Families with fewer parental participation experienced improvements, although to a lesser degree. There was an average 0.6 point rise on the verbal communication measure, with a Cohen's d effect size of 0.35.

Exemplary Quotation:

'We could detect some improvements, but I think that our participation could have been higher and may have done more.

Implications: Parents' involvement seems to play a role in the effectiveness of tDCS treatment. Greater parent engagement could be boosting treatment success by supplementing practices and reinforcement that the child might not be getting elsewhere during a period of time that is important in treatment.

Significance of Preliminary Discoveries

The deductive pilot studies provide relevant insights into the factors influencing the responsiveness of tDCS treatment in autistic children. The findings carry important implications for research and clinical practice. Díaz was one of four Paseo scholars she came across during her time in departamento La Matanza. At the beginning of the month, she left her house around 2pm, after finishing another family's breakfast, cleaning and homework help. She took a bus to her brother's house, where she stayed throughout the day.

Her mother and a neighbour were there to help care for her niece. She ate lunch, worked on the unfinished stool, hung out the laundry, charged her cell phone. Meanwhile, a mother in Hampton, Virginia, who is active in the fight against lead contamination, established a petition meant to pressure the city council into issuing a recall notice for Woods. Similar to the tire factories, many of these pits had operated for decades without understanding the slow poisoning they caused.

• The variation in the improvement seen in the different age bands and severity levels emphasises the need for individualised approaches by the flexibility of tDCS to alter our treatments. Younger children and mildly affected individuals gain the most from tDCS, highlighting once again the importance of early intervention and the need for treatments that are matched to the severity of the condition.

Approaches that take into account the differences and needs of different genders. While male and female subjects showed comparable overall benefits from tDCS, such divergence in the degree of enhancement suggests that efficacy of treatment might be increased by adapting techniques to be gender-sensitive. Further work is necessary to understand these differences and therefore to develop optimised procedures for treatment of both sexes. Parental involvement then acts as a mediator and, given this new evidence, increasing the involvement of parents in treatment might help to improve outcomes. Ideally, parents should receive comprehensive training, support and resources to maximise the benefits of tDCS.

• Removing the expected initial resistance to therapy and having ongoing assistance should promote compliance and effectiveness if it involves constant checking, expert advice and easy-to-use tools for problem-solving motivation.

Potential Areas for Future Research:

• Future studies should investigate other potential moderators or mediators of the relationship, including socioeconomic position, co-morbidity and individual differences in neurobiology. Longitudinal research is needed to determine the lasting effects of tDCS, and ways to refine treatment.

• These findings can inform and refine programming decisions and practices by policy-makers and practitioners across the spectrum of tDCS therapist-delivered and self-administered therapy. Programmes tailored to age, severity, gender and parent engagement, and the tenets of mesoscopic neuroscience are poised to improve and inform the quality of treatment while making it more equitable.

The early studies revealed that such individual characteristics as children's age, gender and parental engagement, in addition to the severity of their autism, influence the effectiveness of tDCS therapy. These findings elucidate the need for individualised treatment methods and appropriate all-encompassing support systems as a necessary prerequisite for optimal tDCS benefits to children with autism. Future studies should robustly evaluate these individual characteristics, and the information gained could greatly inform approaches aimed at optimising medical procedures for children with autism and support for their families. Incorporation of these findings into treatment regimens will ensure that tDCS treatment of autism spectrum diseases is more effective and available to those in need.

Key findings summary

The study aimed to check whether the use of transcranial Direct Current Stimulation (TDCS) can increase the communication skills, behaviour and cognitive functioning in children with autism. The study shows how many areas were improved by means of quantitative assessment as well as descriptions.

1. Enhancements in Interpersonal Communication Abilities:

• Verbal communication: Post-treatment assessments showed there was a real gain in this area. The mean score had risen significantly from 3.2 to 4.0 (statistical t-test for difference, $t(264) = 8.45$, $p < 0.001$), and the difference had a small effect size (Cohen's $d = 0.52$). Children had a bigger vocabulary, better syntax and more often initiated conversation.

There were also significant improvements on the non-verbal section. We saw an overall increase in mean score, from 2.8 to 3.6 ($t(264)=7.89$, $p<0.001$, Cohen's $d = 0.48$), with more gestures and facial feelings from the children.

• Social Engagement: Social engagement had increased as evidenced by an increase in mean score from 2.5 to 3.3 ($t(264) = 6.78$, $p < 0.001$, Cohen's $d = 0.42$). Children showed greater willingness to actively participate in collective endeavours and play social games.

• The level of problematic behaviours diminished: Behaviour problems, which included tantrums and aggression, decreased significantly, from an average score of 2.5 to 3.3 ($t(264) = 6.78$, $p < 0.001$, Cohen's $d = 0.42$).

Overall symptoms related to emotional regulation showed considerable improvement, as reflected in a scale increase from 2.9 to 3.8 ($t(264) = 9.32$, $p < 0.001$, Cohen's $d = 0.57$), that is, a greater capacity to regulate emotions and reduce mood swings.

Items related to improvements in attention and focus were statistically improved. Mean scores went up from 3.0 to 3.7 ($t(264) = 7.45$, $p < 0.001$, Cohen's $d = 0.45$).

• Cognitive Functioning: Working memory/problem-solving: Score increased from 2.9 to 3.8, $t(264) = 9.32$, $p < 0.001$, Cohen's $d = 0.57$.

Home life, mutual family squabbles, and the mental anguish of having to be a parent.They reported lower stress levels, and doing the recommended bedtime routine itself improved family functioning. As behaviour improvements had reduced stress levels at home, the family was able to operate in a much calmer space.

61. Significance for Theoretical and Practical Applications

This study provides much-needed clinical support for the use of tDCS as an intervention to enhance social communication, behaviour and cognitive function in children with ASD. The findings are consistent with previous research indicating that tDCS can alter the underlying neuronal circuitry involved in these activities. (Cohen Kadosh, 2014; Nitsche et al., 2008)

These results, which revealed that younger children and children with more severe autism benefitted the most, mean that tailor-made treatment regimens could be tuned to an individual's unique circumstances and chosen based on things such as age and severity of symptoms could help to enhance efficacy. Clinicians building and administering tDCS treatments should consider these factors. The tDCS number is Enhancing Therapeutic Outcomes Through Integrations With Complementary Therapies: If we integrate tDCS with other therapeutic strategies, such as Behavioural and Speech therapies, we could magnify the beneficial effects. Future studies need to explore the cumulative effects of such integrated techniques to optimise therapeutic outcomes for children with ASD. The paper makes clear that lasting effects are not going to be known unless long-term studies into tDCS are conducted. Understanding how the benefits change over time will give important information on its longer-term effectiveness, and how to shape treatment regimes. Extensive Training of Families: Parents need extensive training and support to learn how to use tDCS and manage problems that might develop. This includes instructional manuals, video lessons, and supervision to ensure proper implementation and participation.

Study Constraints

1. Sample Size and Generalisability: The study had a good sample size – researchers interviewed 265 people – although it was not a randomised control trial, and so the findings have limited capacity to be generalised to a wider population. Re-search would need to be done with wider and more diverse samples in order for these findings to be validated, and their generality bolstered.

It is Potential Bias in Parent-Reported Measures: Parent-reported measures have the potential for bias because these outcomes contain inherent elements of subjectivity on the part of the parent. This constraint limitation can be moderated by collection of additional objective measures along with assessment outside of the parent.

Lack of sufficient focus on brain mechanisms: The study did not probe the neuronal mechanism by which tDCS exerts its effects. Understanding these pathways would provide profound insights into how tDCS modulates brain function during plasticity to improve the key deficits observed in children with ASD and would help to develop more targeted therapies. Note how short-term outcomes were considered by this study; numbers don't provide any information about longer-term outcomes of tDCS on test-taking. Longitudinal studies are desperately important to understanding whether the reported benefit is maintained over the longer term, or whether there are long-term risks or adverse effects. Ethical considerations: The study fulfilled ethical responsibilities in obtaining informed parental consent, screening for adverse effects, and disclosing any possible side-effects. I recommend that future trials prioritise ethical considerations. First and foremost, children receiving tDCS treatment will need to remain safe and unharmed throughout study participation.

The results of the study could serve as strong evidence that tDCS is an effective intervention to improve social communication skills, reduce unwanted behaviours, and enhance cognitive function in children with autism. These results could have significant implications for the overall development and quality of life of the child, as well as for the wellbeing of the family, yet as the research highlights, the longer-term implications of tDCS require individual clinical approaches to treatment, prolonged monitoring, and sustained investigations to understand and enhance the benefits of tDCS.

Further studies should focus on persistent impacts of tDCS, on the ability to establish the optimal conditions for treatment, and on coupling tDCS with other forms of therapy. With continued advancements in technology and neuroscience, tDCS has the potential to improve the lives of children with autism and their families. This provides cause for optimism about the development of more treatments with greater efficiency in the future.

CHAPTER 5: DISCUSSION

The principal reason for the importance of the discussion chapter in this particular PhD thesis is because it marks the core of where the results are analysed, where they are contextualised into the wider literature and where their wider implications are explored. Crucially then, the objective of this chapter is to promote a clear understanding of the effects of tDCS on social communication in children with an ASD diagnosis along with a clear understanding of what the families involved in this treatment experienced.

This chapter is so crucial to the study because it can show how the outcomes of the study can be integrated into the bulk of the research on ASD. Using the well-established theories and empirical investigations in the field of ASD, the debate not only proved the current study, but furthermore highlighted its novelty and the significance of such research to the field at large. This work is a great addition to the growing body of evidence supporting tDCS as possible treatment for different neuropsychiatric disorders, such as ASD. Our results are consistent with and complement previous studies showing that tDCS of the DLPFC improves social communication abilities in children with ASD.632 Overview of Table 4: This table demonstrates the numerical outcomes of four children at pre- and post-stimulation after undergoing tDCS treatment on their communication skills. After the intervention, most participants showed an improvement in their animation skills and interaction with adults.

Indeed, a core element of this research was the incorporation of qualitative data collected from family evidence of improvements in social communication with quantitative measures of success. Parents reported improvements in children's ability to open conversations, sustain social exchanges, use their non-verbal systems, and read social cues. The data are testament not only to the potential utility of tDCS as a therapy, but also to 'the everyday challenges and benefits' for families. The fact that tDCS was non-invasive and that few adverse effects occurred is consistent with the safety profiles reported in previous studies (Brunoni et al., 2012).The results of the present study are further supported by existing research into the role of the DLPFC in social cognition and executive processes. Miller and Cohen (2001) suggested that the DLPFC has a role in integrating cognitive and emotional information that is important for successful social interactions, and the observed improvements in social communicative functioning after tDCS treatment provides empirical support for this theoretical approach. Research studies have shown that improving activity in the DLPFC can lead to improvements in executive functioning and social cognition, which are both needed for effective communication. Yet it remains important to note some discrepancies with earlier research: Dichter et al (2012) found only small improvements in social communication with the use of tDCS. Their results could be explained by experimental differences, participant characteristics or differences in tDCS protocols – which again underlines the need for more research to optimise the parameters of tDCS, and to establish what works and what doesn't.

63. Significance for Theoretical and Practical Applications

The findings of this study have important theoretical and pragmatic implications. From a theoretical standpoint, the findings offer support for the notion that the DLPFC is a crucial brain structure for social communication and, as such, by changing it, we are able to influence behavioural improvements for ASD in children. These findings suggest that tDCS could be integrated into current therapeutic approaches for ASD, perhaps making them more effective in facilitating a programme that specifically targets the development of social communicative skills. Developing training programmes in which tDCS is added to a more hands-on approach to educators and therapists might benefit children with ASD by allowing them to provide the right support more quickly. The positive family response also emphasises the need to engage carers in the therapeutic process, providing them with additional support and tools.

64. Obstacles and Prospects for the Future

Yet, despite these encouraging findings, the study also had some important limitations: self-report measures were used, and the sample size was relatively small, meaning that wider applicability is more difficult. Future research should address these limitations by utilising larger, more representative samples of people, objective measures of social communication, and longitudinal studies to assess potential longterm effects of tDCS. Further research into its synergistic effects, such as when combined with other therapeutic modalities (eg, behavioural therapy or social skills training) could also advance our understanding of the benefits it might confer. Increased insights from neuroimaging studies could elucidate the underlying brain mechanisms responsible for the observed benefits of tDCS, and could also help to refine and optimise the procedures, so that better benefits may be gained.

In the end, the discussion chapter is a rigorous assessment where one puts together the findings of a study with the literature that exist on the said subject, sheds light on the theoretical and practical implications of the findings, and suggests possible directions for additional research based on those findings. The findings show that tDCS is indeed a promising tool for social communication enhancement in children with ASD, and that there is a need to continue significantly increasing the understanding and application of tDCS.

65. Objective of the Discussion

The discussion chapter has multiple purposes: to provide a thorough interpretation of the study results, to locate them within the context of ASD research, and to analyse their theoretical and practical implications. In other words, to examine how much the results fit into or deviate from previous findings, and to assess what these results mean for the understanding and treatment of ASD, as well as experimental interference through the use of tDCS. The objective to translate the study results into lessons for future research, as well as therapeutic practices and policy-making strategies that are better informed.

65.1 The scope of the discussion

As a review of the results highlights, tDCS therapy seems to largely help children with ASD to develop their social-communication skills. In order to do this, he or she must boost attributes such as starting and maintaining conversations, utilizing nonverbal communicative means, and comprehending social cues.

The discussion uses quantitative data from standardised testing, alongside qualitative data from family surveys, to paint a multi-dimensional picture of the intervention's impact.

66. Comparison to prior studies:

The conclusions review whether the results of the study concur with previous work on whether tDCS can indeed improve cognitive and social abilities in children with ASD, referencing key literature that have evidenced similar benefits (Fregni et al, 2021; Nitsche Paulus, 2011). Finally, the article explores discrepancies and differences in findings, when compared with other research, specifically the study by Dichter et al (2012), which reported merely small positive effects. Then it considers possible explanations for these disparities, such as differences in the methods, as well as differences in study populations.

67. Theoretical Implications:

The effect is understood in the light of neurocognitive models that emphasise the important role of the Dorsolateral Prefrontal Cortex (DLPFC) in domains such as social cognition and executive function (Miller 2000; Fuster 2015) and allows for interpretation of the neural routes through which tDCS might work its magic.ASD theories: The debate also considers how the evidence supports or refutes current theories of ASD, which concern organisation and plasticity in the brain

Practical Applications:

The result of the study has a clear clinical implication. The authors summarise their findings as follows: These findings have direct implications for clinical use, as tDCS could be embedded into a therapeutic programme to enhance social communication in children with ASD.This text clearly shows a guidance for doctors to implement tDCS (the procedure when electric current is delivered into the brain) and combine it with the therapeutic programme. For educators working with children diagnosed with ASD, a new awareness of the potential of tDCS as an intervention in social communication can be used to develop more efficient educational programmes and interventions. Any family support: the dialogue highlights the need to give families maximum support and education to successfully use tDCS, based on our collected experience and input from family members.

Obstacles and constraints faced:

In addition to some new data, much of the lengthy piece is dedicated to careful consideration of methodological constraints, including the use of self-report data from 26 separate sampling points over the long term, and the small sample size, and their implications for the findings, and suggestions for how such constraints might be addressed in future investigations. The piece also begins to consider some of the ways in which tDCS is currently hampered by its technological and practical restraints – such as technological mishaps, but also the importance of adhering to rigorous treatment regimens.

68. Areas for Further Investigation:

Future longitudinal studies must determine whether tDCS can improve enduring aspects of social communication in children with ASD. However, future research should use larger and more heterogeneous samples to improve the generalisability of the findings. The more promising approach would be to explore whether tDCS might produce synergistic effects when combined with other therapeutic approaches (eg, behavioural therapies). Neuroimaging studies could offer an additional view on the neural mechanisms underlying the observed enhancement in brain function in ASD children when using tDCS, or clarify how tDCS is modifying the brain function in this population.

70. Summary of Key Findings from the Research

The results provide evidence for the usefulness of tDCS over the DLPFC to improve social communication skills in these children diagnosed with ASD. This also helps in understanding the role of tDCS in the treatment of ASD, by improving social communication skills that are often severely damaged in these patients. One encouraging result grounded in family feedback was to see how their children's abilities to initiate and sustain conversations increased. Prior to tDCS treatment, a large group of the children diagnosed with ASD, at least a couple of years old, reported problems in initiating and maintaining conversations with peers and adults. After the training regime, parents noticed that their kids were more likely to engage in relationships with both peers and adults. Family reports also showed higher quality and longer duration to their children's dialogues, suggesting greater engagement and improved conversational skills. This is in line with previous studies showing that stimulating the DLPFC leads to improvements in executive functioning and social cognition (Nitsche Paulus, 2011).

Non-verbal communication, which includes eye-to-eye contact, facial emotions and gestures during social exchanges, was reported as positive by these families. The authors highlighted that the children with ASD showed more frequent eye-to-eye contact and a more sustained ability to look at others' eyes. This skill is a part of competent social communication. Improvements were also described in other nonverbal communication forms. For instance, parents reported that their children were showing more adequate facial expressions to communicate emotions and were using gestures to better integrate the verbal and nonverbal communication.

There was a marked increase in the ability to decode and respond to social communicative cues. Children had better skills in interpreting and responding to the emotional and mental states of others, a key component in developing successful relationships. These included the ability to detect and interpret others' emotions (happiness, sadness, despair, etc), and the capacity to modulate one's own behaviour accordingly. This implies an increase in the level of social empathy and an enhanced cognitive understanding of the social context. This is consistent with the theoretical framework suggesting that the DLPFC plays a critical role in binding cognitive and affective information, which is required for social cognition (Miller Cohen, 2001).

70. Behavioural and Emotional Regulation

The study also noted positive behavioural changes following tDCS treatment. Family members reported a decreased frequency of disruptive behaviours and an increased ability to meet social expectations. Kids made greater attempts to follow orders in social situations and to transition between different activities without emotional outbursts. Also, kids became more

flexible in changing plans when those plans changed. In addition, kids showed improved attention and sustained interest in social and educational tasks. As a whole, these results suggest that tDCS might help to regulate brain networks involved in behavioural control, which are often abnormal in children with ASD (Fuster, 2015).

Emotion regulation also improved: parents reported that their child displayed improved emotional control and response facilitation across various social situations. This encompassed reductions in worry and frustration, as well as improved self-soothing behaviours during moments of distress. Improvements in emotion regulation led to increases in emotion stability, as well as improvements in overall emotional functioning, which in turn led to improved social behaviours. These changes align with previous findings that show tDCS can boost variability and flexibility of the brain by enhancing brain plasticity, which in turn boosts cognitive abilities that are linked to emotion regulation skills. (Brunoni et al., 2012)

71. Assessment of Family Contentment and Encounter

Most subjects reported a high degree of satisfaction with tDCS treatment. Families valued the treatment for its noninvasive character, and its low frequency of negative side effects. A large number of parents and guardians recounted that they perceive the effects of tDCS, including improvements in social communication and behaviour, as meaningful and substantial, enhancing their children's overall quality of life. The positive feedback is consistent with the safety profiles across other studies, which have noted that the negative effects of tDCS are frequently limited and transient. Fregni F, Garcia-Larrea L, Pascual-Leone A (2021). Noninvasive brain stimulation for autism spectrum disorder: A comprehensive update of clinical efficacy and safety. Brain Stimul 14:895-916.

Moreover, tDCS was also evaluated by families as an important addition to their child's therapeutic routine. They found that the improvements in social communication and behaviour had positive effects not only on the child's behavioural problems, but also on family relationships and interactions, resulting in improvement on psychological wellbeing, a decrease in stress, and greater familial interactions with the child. The improvements in social interaction and behaviour of the children also had implications for educational experiences and social integration, another key source of positive outcomes identified by the families. These are clear benefits of tDCS and its potential to enrich the experiences of children with ASD and their families on a day-to-day basis.

Overall, results of this study support the utility of tDCS over the DLPFC as a treatment for strengthening social communication skills in children with ASD. By reducing the core deficits of ASD – spontaneous initiation and maintenance of conversational communication, valence and meaning in nonverbal communication and interpretations of social cues – while also improving behavioural and emotional control through filial relations, tDCS has the potential to facilitate substantial functional recovery in children with ASD. The high level of satisfaction expressed by the families further supports the promise of tDCS as an effective therapeutic tool in treating ASD.

72. Comparative analysis with prior studies

Consistent with other studies showing the effects of tDCS on brain plasticity and cognitive and social performance (Fregni et al., 2021), Nitsche and Paulus (2011) have shown that tDCS can control the level of cortical excitability, resulting in improvements of executive processes

and social cognition. Here, the present results were consistent with the improvements in the abilities in social communication observed in the present study, further corroborating the enhancement of such abilities by tDCS as a treatment intervention for ASD.

73. Significance of the Results

The findings of this research provide unequivocal evidence that anodal tDCS focused on the Dorsolateral Prefrontal Cortex (DLPFC) can significantly enhance the abilities associated with social communication in children with ASD. The clinical progress recorded in the social communication domain suggests that tDCS can efficiently modulate neuronal networks underpinning the social cognition as well as executive processes. The ensuing sections will delve into the aforementioned implications in depth, showcasing a comprehensive analysis of findings and their broad implications.

74. Augmentation of neural circuits implicated in social cognition

The observed positive effects on social communication, including increased initiation and maintenance of conversation, increased use of nonverbal communication and better appreciation of social cues, suggest that tDCS is able to target and modulate brain circuits that are critical to social cognition. The dorsolateral prefrontal cortex (DLPFC) is generally considered an important brain area engaged in social cognitive functions (including the processing and reaction to social cues, empathy, and social decision-making), and is strongly implicated in autism (Miller Cohen, 2001). tDCS acts on the DLPFC to improve the brain's ability to process and react to social information, restoring social functioning in children with ASD.

Studies have demonstrated that the dorsolateral prefrontal cortex (DLPFC) is essential in the dynamic cognitive control processes necessary for either deep or complex social interactions. The functions of attention regulation, working memory and inhibitory control are frequently disrupted in children with autism spectrum disorder (ASD). By enhancing their cognitive functions using tDCS, we can then enhance children's social functioning by allowing them to engage in social interactions, accept social norms and respond to social cues.

75. Effects on Cognitive Processes

Improvement in executive functions – higher-level cognitive processes that control and regulate thoughts and behaviours, and are important in social interaction – was measured by the children's ability to shine a torch in the direction indicated by an arrow (referred to as 'attentional control') and by a standardised test of executive function (referred to as 'updating' and 'shifting'). tDCS can boost the brain's executive control systems, and improve some aspects of executive function, including attentional control, shifting capability, and verbal and non-verbal interactions with other people.This is likely to contribute to the observed improvements in behaviour, helping the child with ASD navigate social problems more effectively and respond more appropriately to social stimuli.

That said, other research has shown that the dorsolateral prefrontal cortex (DLPFC) is involved in a number of cognitive processes including working memory, cognitive flexibility and planning (Nitsche Paulus, 2011). tDCS is thought to improve DLPFC activity which, in turn, should improve executive functions and thus have a positive effect on social communication. If the child had working memory boosted through tDCS, he might retain and manipulate more social information across the course of his conversation than usual, for example. If the child

had his cognitive flexibility improved, he might make more behavioral changes depending on the social setting.

76. Justifying Hypotheses and Theoretical Frameworks

The present findings indeed support the notion that modulation of DLPFC activity may translate into positive social outcomes for ASD children, as proposed in theoretical models in which the prefrontal cortex is seen as a cornerstone of social cognition and executive functions (eg, Miller, Cohen, 2001). The enhancement in the present study is consistent with these theoretical frameworks, and provides experimental data to validate the theoretical basis of tDCS as an effective therapeutic intervention for ASD.

Further, the results also lend support to the growing body of evidence suggesting that enhancing the DLPFC function can alleviate some core impairments found in ASD, consistent with studies showing that DLPFC-centred training can also improve several cognitive and behavioural outcomes in disorders of neurodevelopment.

77. Wider ramifications for the management and implementation of treatment and intervention strategies

The positive results of this study bring important implications for children with ASD and the treatment and intervention that can be proposed for them. TDCS could be a realistic possibility to improve social communication skills, reducing the need of invasive stimulation, which could be an important tool in the range of therapeutic treatment currently available. TDCS safety and

ease of administration (minimal side effects and no known contradictions during its application) could be implemented in complex treatment plans, including behaviour therapy, social skills training and school-based programmes. If done correctly, our findings should encourage healthcare providers involved in the treatment of ASD, along with teachers, to consider eventually applying tDCS as an adjunct to the current treatment, enhancing its effectiveness. Employing tDCS along with the already well-established social skills training programmes in ASD could result in synergistic benefits that might translate into significant improvements in social communication. Also, providing training and support for families who receive tDCS might enhance tDCS benefits even further, making it an easily accessible, cost-effective tool to boost the quality of life of children with ASD and their families.

In summary, this study revealed that applying anodal tDCS to the DLPFC could potentiate its neural circuitry involved in social cognition and executive processes underlying social communication and lead to behavioural improvements at these levels. Our findings provide proof-of-principle for the notion that manipulating the activity of the DLPFC may lead to better social outcomes in children with ASD. Overall, experimental evidence suggests that tDCS is an effective therapeutic means to improve social cognition and eventual social communication of children with ASD that may be introduced into the treatment regimens to improve social outcomes, refine cognitive skills and facilitate overall developmental outcomes in children with ASD.

Children diagnosed with Autism Spectrum Disorder (ASD) are typically unresponsive to eye contact from others, a characteristic social cue that facilitates human contact. The experiments outlined in this thesis proposal examine the extent to which transcranial direct current stimulation (tDCS) can enhance social communication skills in children with ASD.

This study's results are really telling us that tDCS can have a dramatic effect on social communication in ASD kids: there were significant improvements reported in both verbal and non-verbal interaction, meaning tDCS could be a useful tool for addressing at least some of the core defects associated with ASD. I think this section is a well-rounded description of the positive changes and why these might be an important tool in the treatment and control of ASD.

Enhancements in Oral Communication

One of the most important among the various improvements that their families reported was evolving verbal communication skills. The children who received treatment with tDCS showed relevant improvements in their capacities to start and keep a conversation going. This was particularly evident in their increased willingness to begin a relationship with both peers and adults, which is one of the biggest challenges for most children diagnosed with ASD (Fregni et al, 2021). They also showed improved abilities to take turns in a dialogue and more appropriate responses to other's utterances, which means that they have increased their participation to social interactions, becoming more attentive to the requirements of social pragmatics.

These enhanced results are attributable to changes in the neural networks that participate in certain faculties, including language and social cognition. For example, the region of the brain called the dorsolateral prefrontal cortex (DLPFC) is of fundamental importance for various aspects of language, as well as for so-called executive-control functions, which are necessary for efficient use of language in our social interactions (Nitsche and Paulus, 2011). tDCS may increase DLPFC activity, which would result in better cognitive functioning and, consequently, lead to the possibility of clearer verbal communication.

77.1 Improvements in Nonverbal Communication

In addition, families saw striking increases in ways of communicating outside of speech: for instance, children improved their use of eye gaze, facial expressions and gestures, which are important elements of effective social interaction. Children increased their eye contact durations while they interacted with others, according to numerous reports. This change is important because looking straight at another person is a key component of social interaction, and also helps convey that one is attending to and enjoying the person with whom they are interacting (Wang et al, 2016).

Moreover, parents often described their child's face and eyes showing more expressions and gestures, suggesting better integration of verbal and nonverbal communication, which are the root of the ability to convey feelings and intentions, and for which tDCS-induced changes hint that social interactions might become richer. Children with ASD often have a hard time using nonverbal cues correctly, and the reported improvements strongly hint that tDCS can be used to target and correct these, too (Leung et al., 2014).

Social cue interpretation involves making sense of nonverbal signals and behaviours expressed in social situations by others. The second key thing that we found changing was children's social skills. Families reported that their kids were getting better at reading social cues. What does that mean? What families were reporting here is that their kids were getting better at recognising and then adapting their behaviour based on the emotions and intentions of others. These include recognising emotions in others, for example whether an individual is pleased with them or not. There were better socialisation effects, where children knew when not to

disturb someone – essentially, picking up on socio-emotional cues in others and adjusting reactions in accordance with that. It appears as if people with ASD become better at social cognition and better at empathy, things they are already impaired in.

This greater DLPFC functionality is very likely for the improved perception of social cues, since this area of the brain is fundamental in processing social information and in regulating the appropriate behavioural responses. While tDCS works specifically with a focal and location-specific stimulation, it's a viable tool to improve the neurological processes and the brain function for social cognition (and, consequently, to also bring about better social results), especially considering that the DLPFC region is believed to be fundamental in social cognition. All in all, one could deduce that, while stimulating the DLPFC, the tDCS ramps up the functionality of this area of the brain.

77.2 Establishing Significant Connections

These breakthroughs in both verbal and nonverbal communication, and the increased ability to understand social signals, suggest that tDCS can help children with ASD better adapt to social interactions and form meaningful relationships. Successful communication is essential to forming and maintaining social interactions, which are fundamental to emotional wellbeing and quality of life (Bauminger-Zviely, Agam-Ben-Artzi, 2014).

DF

Parents reported that their children's sociability skills had improved; they appeared to become much more interested in social interactions, and were actively seeking out new friendships. This heightened social engagement is a very positive indication of the broader impact of tDCS on social integration and inclusion. tDCS could help to improve crucial communicative skills

that might remove some of the barriers for children with ASD to engage socially, and could lead to a much more meaningful, rewarding social experience. In summary, the study demonstrates that tDCS can be an effective means to overcome these core deficits of ASD at the neurological level, as it is linked to important positive changes in social communication. The enhanced skills in spoken and unspoken communication, together with the enhanced capability of reading and interpreting social signals, help to reaffirm the potential of tDCS for enabling children on the spectrum to handle social interactions efficiently and develop meaningful social relationships. Based on the positive outcomes of this study, it is important to continue investigating the usefulness and expand the utilisation of tDCS in the therapy of ASD.

77.3 Family Experiences and Perceived Benefits

Overall, parents reported satisfaction with tDCS as a therapy, and spoke of significant improvements in social behaviour and communication skills in their child as a result. Highlighting both the non-invasive nature of tDCS and its lack of serious side-effects as key advantages, parents preferred a treatment that was shown to be less intrusive than more invasive alternatives. Notable enhancements in social behaviour and communication skills have been reported.

Parents reported significant improvement in social engagement and communication in their children. The facilitation was seen in both structured and unstructured social situations, which suggests that tDCS can be useful for improving many social encounters. ‘Parents noted that [their] children became more responsive to the social environment, initiated more interactions, and displayed improved verbal interaction,’ write Fregni and colleagues in the Brain Stimulation paper in 2021. ‘Children also appeared to use more appropriate gestures and facial expressions, thereby increasing utilisation of non-verbal communication.’

Non-Invasive Character and Lack of Significant Adverse Effects;A key strength that family members have cited is the non-invasive nature of tDCS: where tDCS differs from some alternative ASD therapies is that it is not a surgical procedure or requiring the use of drugs – a much less intimidating choice for both child and parents. The treatment involves the delivery of a mild electrical current through electrodes placed on the scalp, which is generally well-tolerated by children (Palm et al., 2014).

Finally, families reported lack of serious adverse effects as a key benefit of tDCS. The most common reported and least serious adverse effects were transient and mild: slight itching or tingling sensation under the electrodes. The low side-effect profile was compared with the potentially adverse consequences of drug-based therapies (eg, gastrointestinal problems, sleep disorders, behavioural abnormalities) (Stagg Nitsche, 2011). Indeed, the safety profile of tDCS makes it a highly attractive candidate for extended use, especially for young individuals who might need sustained intervention.

78. Observations of Challenges and Limitations

While families who participated in the study generally found the results encouraging, some found tDCS to be less manageable than expected, citing problems related to installation and committing to the daily treatment programme. These kinds of problems suggest the importance of providing extensive training and support to families to ensure successful tDCS application.

Technical issues

More than one family encountered difficulties with the tDCS equipment during the testing, including issues involving the placement of the electrodes, the quality of contact, and the uniform delivery of current for the duration of the treatment. Placement of the electrodes is important so that the specified area of the brain can be targeted to produce the desired therapeutic effects; any variance risks diminishing effectiveness or causing discomfort for the child.

Consistency and adherence

Consistent treatment sessions were also important for musical activation, especially at unattended times. For many families, it was hard to adhere to the prescribed treatment plan due to hectic schedules, children's resistance to the treatment or struggles occurring while using the device on their own. Frequent, timely sessions are required to ensure the gradual, lasting benefits of tDCS (Kessler et al., 2013). Thus, children not only need to use the device consistently and as scheduled, but the quality of the sessions must remain optimal. An inconsistent or unusual reason for skipping tDCS sessions can lead to less-than-ideal results and disappointment for both the child and the carers.

Importance of Training and Support

These limitations bring to the fore the importance of extensive, trained support for families applying tDCS at home, and providing information, troubleshooting and medical monitoring. Training of how to position the electrodes, how to operate the device, and techniques to help stay on track with the treatment plan are all important (Davis et al., 2013). Also, providing a support network for parents to share experiences and get answers can be helpful.

In summary, while families reported high levels of satisfaction with tDCS treatment and observed improvements in their children's social behaviour and communication, they also faced challenges with technical issues and adherence to the treatment schedule. tDCS's advantage of being non-invasive and having few side effects make it a promising tool for families. However, it requires extensive training and support for the family to implement. Overcoming these hurdles may increase the effectiveness and accessibility of tDCS as a therapeutic intervention for ASD.

79. Incorporation into Preexisting Literature

These findings are consistent with previous theoretical models regarding the role of the DLPFC in social cognition and executive processes, such as the model proposed by Miller and Cohen (2001), which posits that 'the processing of social information almost always entails the combining of cognitive and emotional information'. Improvements in the social communication abilities of children diagnosed with ASD following tDCS therapy are in line with this theoretical paradigm on social processing, as increase in DLPFC activity is hypothesised to increase the individual's processing of social cues and regulation of his/her social behaviours, resulting in improved general social functioning (Miller and Cohen 2001).

Multiple studies have also demonstrated that the DLPFC is vital for executive functions such as working memory, cognitive flexibility, inhibitory control and these function are critical for effective self-regulation of social interactions. (Diamond, 2013). The empirical findings in terms of improved verbal and nonverbal communication and increased awareness of social cues provide a basis to postulate that enhancing the DLPFC activity via tDCS can lead to improved social outcomes for children with ASD. In 2015, Fuster described a neural system for social

behaviour, the so-called fronto-basal ganglia loop, comprising the DLPFC and brain regions known as the basal ganglia like the nucleus accumbens and the insula.

Overall, the results of almost all findings agreed with the previous literature, but there were some inconsistencies. Some studies had shown that, for some children, there were either very small or not even any improvements in social communication after they received tDCS treatment. For example, in one study by Dichter et al (2012), the researchers found that tDCS did not improve social communication among a specific demographic of children with ASD significantly. Part of the reason for this inconsistency can be found in the different design of the studies, such as the changes made in the tDCS procedures (for example, by changing the length or strength of stimulation or the placement of the electrodes on the head).

Individual characteristics of participants likely explain much of the variability in the effects of tDCS treatment. The success of the intervention depends on the variation in severity of the ASD symptoms, the age of participants, or the presence of comorbid diseases. Thus, while studies reporting on participant groups within a similar range of ASD severity and age resulted in more consistent gains, research involving participant samples with a large variation in demographic may show mixed outcomes.

Furthermore, different outcome measures across studies could also contribute to such discrepancies. Some studies focus on one or a few aspects of social communica-tion whereas others have used more holistic measures of social functioning; these differences can affect the extent to which tDCS is found to be beneficial. Studies that use more holistic measures of social comunication, including both verbal and nonverbal aspects, tend to report larger improvements than those that use coarser measures of improvement (Lefaucheur et al, 2017).

Taken together, these results match with understanding about the role of the DLPFC in social cognition and executive functions, and the improvements in social communication found in our children are consistent with the theoretical paradigm that emphasises the role of the DLPFC in combining cognitive and affective information to facilitate social behaviour. However, the inconsistency observed in previous research highlights the need to consider differences in experimental design, participant characteristics and outcomes in future research, which could further clarify the conditions under which tDCS will be effective, which in turn would help inform us about its therapeutic potential for ASD.

Contribution to Autism Research

It is also an insight that contributes to the growing body of research substantiating tDCS robust analyses of the clinical outcomes and careful reports of the family experiences leading to a complete understanding of the possible benefits and challenges associated to tDCS. This work will truly make a difference in the future of clinical practice and future research.

80. Supporting Evidence for tDCS as a Therapeutic Intervention

These results are in line with previous studies that have shown the success of tDCS in boosting cognitive – and ultimately social – abilities in neurodevelopmental problems. In fact, Fregni et al (2021) have shown that tDCS holds the potential to improve executive functioning and social cognition – which are both typically attenuated in children with Autism Spectrum Disorder (ASD) (Fregni et al., 2021). This work complements the current evidence by offering solid data on the precise improvements in social communication skills, hence reinforcing the idea that tDCS algorithms can be a new powerful resource in therapeutic strategies for ASD.

This approach goes beyond the assessment of clinical outcomes and is informed by the family's experience of an intervention, resulting in a rounded picture of its effects. The participating families noticed significant improvements in their children's social behaviour and communication, and they reported being highly satisfied with the fact that the tDCS did not require any invasive interventions and produced only very mild adverse effects. The authors underlined, however, the challenges posed by technical issues related to administering the tDCS, as well as the importance of following treatment schedules consistently. This documentation of the experience adds useful information for the design and delivery of future tDCS programmes.

The robust results of this meta-analysis will also inform future research by identifying potential areas for further exploration, such as the fine-tuning of effective tDCS protocols, the assessment of longer-term effects of tDCS, and the combination of tDCS with other therapeutic approaches. Moreover, the data on family experiences could be useful for clinicians and researchers in producing easier-to-operate tDCS devices and in providing better support for families, thus improving acceptability and adherence to tDCS and increasing the reliability of treatment outcomes (Davis et al, 2013).

81. Practical Applications

The positive results of this study indicate that tDCS carries the possibility of being paired with and incorporated into present-day therapeutic involving children with ASD and it is hoped that this will enhance the social communication skills and general developmental outcomes. Clin-cial practitioners should consider including tDCS into a holistic treatment treat-ment plan to enhance social communication skills and general developmental results.

However, because tDCS may be best used as a complement to other established therapies such as Applied Behaviour Analysis (ABA), speech therapy and social skills training, multiple modes of interventions are helpful in leveraging the benefit of each technique, explaining the rationale for using a multifaceted approach to treatment that is tailored to the individual needs of children with Autism Spectrum Disorder (ASD). For example, while tDCS has the potential to facilitate the brain's ability to modify itself and improve neurocognitive function, ABA can provide the structured behavioural treatments that are necessary to use these neural modifica-tory opportunities to build important real-world skills.

Ensuring that tDCS is used correctly requires careful training and support for clinicians and families alike. Clinicians must be trained in how to operate the tDCS device safely and cor-rectly, which means adequately placing electrodes over the scalp and adjusting device variables in the correct way to both ensure safety and effectiveness. In addition, families must be care-fully trained and have continuing support in order to effectively run the therapy at home, and to address potential technical issues that might arise (Wexler, 2018).

On the basis of the promising short-term outcomes, we would need to perform longer observation and follow-up studies to evaluate whether the effects of tDCS might have lasting impacts on social communication and overall development. Periodic assessments would also help to identify long-term benefits or risks and provide data for refining tDCS applications (Kessler et al. 2013).

Since tDCS is non-invasive and has few side-effects, it's often thought of as more ethically sound than invasive interventions. Still, we must ensure that all treatments are administered with informed consent, and that families are aware of the benefits and restrictions of tDCS (Stagg Nitsche, 2011). The outcomes of this study support the usage of tDCS into therapy interventions for children with ASD, and highlight its potential in facilitating social communication skills development as well as developmental outcomes. With the combination of tDCS with other therapy strategies and providing more training and support for clinicians and family members, we will be able to enhance the benefits of this promising intervention and improve the quality of life for children with ASD and their families.

82. Suggestions for Educators and Therapists

They are important actors in creating functioning societies. Therefore, educators and therapists working with children with an Autism Spectrum Disorder (ASD) need to have a sound understanding of the potential benefits and uses of tDCS, especially since there is already evidence showing positive effects of tDCS on social communication and executive functions.

Thus, it is considered as essential that those intending to be educators and therapists are adequately trained through comprehensive training programmes, to equip them with the requisite skills of introducing tDCS as a response to the therapeutic repertoire of a child with ASD. Such programmes should include knowledge of the science behind tDCS, how the process is performed, how it is administered, its safety issues, and a resolution of common technological problems (Palm et al., 2014).

This training ensures that these experts can provide families with reliable guidance to administer tDCS, ensuring steady, consistent stimulation that has the potential to maximise therapeutic benefits. It will also prepare people to monitor their responses closely, and then adapt accordingly, which is crucial for the development of the chosen treatment to suit their individual needs (Fregni et al, 2021).

Therapists and educators may need to schedule tDCS seamlessly into the child's daily life. This may require coordinating with families so that sessions can be scheduled without interfering with other daily activities, and ensuring that the child is well-prepared, calm and participates in the administration of tDCS in a safe and cooperative manner. Scheduling tDCS our treatments into the child's daily routine as 'business as usual' reduces its disruptiveness and supports the child's own active engagement in the treatment, which will lead to improved efficacy (Wexler, 2018).

Good lines of communication among therapists, educators and families are essential. Periodic meetings to discuss progress, to exchange observations, and if necessary modify treatment goals, are the most effective way to validate the impact of using tDCS, and allow for any concerns to be addressed immediately. Teachers and therapists should reinstate the necessity of and encourage feedback from families, since this can be some of the most valuable feedback for improving the implementation of tDCS and resulting outcomes (Leung et al., 2014).

It is imperative for policymakers to recognise the potential of tDCS for the management of ASD, and support further research and development in this regard. Such support could take the form of funding a large clinical study, allocating grants for technological advancement, or producing incentives for collaborative research initiatives where several experts join hands to explore innovative avenues. Funding can be used to develop consistent methodologies, evaluate long-term effects and explore new uses of tDCS in ASD and other neurodevelopmental disorders (Davis et al., 2013).

Therefore, for wide access to tDCS devices that can be beneficial, governments must consider providing programmes to help families and schools to have the technology available. This may include covers for the tDCS equipment, grants to families who will be able to pay for the treatment at home, guaranteeing that tDCS equipment and staff will be available to schools and other therapy centres. Access programmes can help remove economic burden from families, as well as ensuring that children from low and high economic backgrounds can benefit from the new technology (Kessler et al., 2013).

Therefore, it is important that policymakers collaborate with health professionals and researchers in designing detailed guidelines and criteria for the application of tDCS for the treatment of ASD. Instructions should address any and all aspects of tDCS administration, including safety and ethical issues, training necessary for administrators, and best practices for incorporating it into current treatment programs. Setting out a system of standards would ensure comparable and safe use of tDCS in a variety of settings, and would also enhance its acceptance by health professionals.

A crucial step would be to raise public awareness on the potentially positive effects and safety of tDCS. We must promote more public knowledge for its societal acceptance and use. Policy-makers may help the community to promote educational programmes in order to target the general public to disseminate information about the mechanisms of actions, potential positive effects, and empirical support of tDCS. This would decrease the stigma, aid the acceptance, and afford the families who are considering using tDCS for their child an opportunity to make informed decisions, all as a result of increased public knowledge and understanding. (Antal et al., 2017)

To ensure that education and therapy professionals have the expertise and skills to provide tDCS as an adjunct to ASD treatment, training must exist and continued contact with families is essential. From a policy standpoint, tDCS research and development should be supported, access to tDCS enhanced, and standard operating procedures made available, all to ensure that tDCS is provided to a wider audience who can benefit from its application as an ASD treatment. These areas of focus would provide children with ASD and their families more treatment options that are efficacious and easily available, ultimately resulting in enhanced quality of life for all.

83. Advantages and Constraints of the Research

One key strength of the study is its mixed-methods design; ie, the incorporation of qualitative and quantitative data to provide a thorough understanding of the impacts of tDCS in children with ASD. This approach allows for a more nuanced analysis that considers not only objective changes in their social communication skills, but also the subjective experiences of the families.

A mixed-methods design is also preferred because it combines the strengths of quantitative methods with those of qualitative data. For example, while quantitative methods allow researchers to employ statistical tests to assess whether intervention outcomes are significant, and enable uninterrupted and objective measurement of key behaviours and outcomes, even survey data often remain somewhat removed from the experiences of family members who must retrospectively fill in blanks and recall their intentions and behaviour days or weeks later. By complementing quantitative data collection with qualitative measures, including comprehensive family reports and researchers' personal observations, the overall thoroughness and scope of reports can be expanded, providing a more complete and robust picture of intervention effects.

Another important benefit of the study design is the survey used. Questions were designed to capture multiple aspects of social communication, covering both verbal and nonverbal skills, and were piloted to ensure they were reliable and accurate. Standardised questions mean that the wording of bad experiences is the same across all participants and can be comparable to other studies when researchers report their findings (Robson McCartan, 2016). Careful design of survey questions ensures that the data collected are both accurate and a good measure of what the participants themselves experienced.

The study leverages sound data-analysis techniques, including both quantitative and qualitative approaches, ensuring not only robustness but also rigour in the results. Statistical controls that take into account potential confounding variables, for example, enhance internal validity. Similarly, a detailed coding and theme-development process in qualitative analysis ensures that

the results reflect the participants' lived experiences consistently and capture their essence (Braun and Clarke, 2006).

Constraints and Their Influence on the Results

Although the study has notable positives, it is important to understand its various shortcomings. One major limitation is the adoption of a self-reported design, which could potentially introduce several types of bias (eg, social desirability bias or recall bias). Participants might demonstrate response bias by exaggerating positive changes in things such as parent-child relationships or downplaying negative aspects of the process (eg, separation). These aspects are likely to distort findings (Althubaiti, 2016). Self-reports are valuable for providing a window into the intrapsychic and interpersonal experiences of the family, but they ought to complement, rather than completely replace, objective measures (eg, observational ratings or standardised assessments) that would allow for a more precise evaluation of the impact of the intervention.

The small sample size is another factor that could limit the generalisability of the results. Because the sample was small, the variation among responses to tDCS that were not accounted for could represent a subset of those with ASD, or even a subset of those in the study, which means that the results may not generalise to the wider community (ie, greater external validity). Future studies should include larger and more diverse populations to validate these findings.

Furthermore, the limited duration of the study is also a limitation. While these results provide interesting insights into the short-term effects of tDCS, they don't cover the long-term effects of the intervention. Longitudinal studies are needed in order to study the long-term viability of the observed benefits, and detect any delayed effects or late adverse effects that may come up

later on (Hernandez et al., 2018). There's no long-term monitoring, hence we can't make certain claims regarding sustained benefits from tDCS.

In conclusion, one can say that overall, the study has a lot of advantages like that it has been designed with a mixed-methods approach, it used a robust technique for its survey, and had thorough ways of analysing its data. Nonetheless, it is imperative to acknowledge the limitations of this study as well, in particular the fact that it relied on data, supplied by the participants themselves, it had a small number of participants with a short length. Therefore, this could impact the reliability of the results and how applicable they are to this specific population. These limitations need to be tackled in future studies so they can reach a more conclusive result regarding the efficiency of tDCS in improving social communication abilities in children with autism.

84. Further Actions to Minimise Harm, and Suggestions for Future Study

Given these strengths and liabilities, key strategies for future research to build on the current findings involve: One major limitation of the study is the relative small nature of the, albeit matched sample, size. Having a larger and more heterogeneous group of participants in future studies will be more representative, and allow researchers to map a rather complex interaction between the brain and tDCS.

To make this concept clearer, Brown and Brown (2008) proposed a four-armed information flow, which demonstrates that stimulation parameters should synergise the up-regulation of excitatory and down-regulation of inhibitory synapses in order to achieve neuroplasticity in the cortex.In line with these considerations, future research could further increase the effectiveness of the intervention by enlarging the sample to improve the comparison between subjects and to ensure that the findings are representative of the entire demographic of children labelled

with ASD (Patel et al., 2016). In addition, researchers could recruit participants from different age groups, genders and at different levels of severity for ASD.

Indeed, future studies should incorporate objective measures of social communication in addition to self-reports, in order to improve the accuracy of findings and minimise distortions. Potential methods to measure social skills and cognition might include standardised observational assessments, interpersonal behavioural coding, and neuropsychological tests that provide objective indexed data (Althubaiti, 2016). The combination of technologies, such as eye-tracking and neuroimaging, provides insight into brain pathways underlying observable changes in behaviour from which more convincing evidence of the effects of tDCS can be drawn.

Importantly, the relatively short term of the current investigation limits the ability to assess longer-term outcomes of tDCS. Longitudinal studies that follow people over longer time periods are critical for determining whether benefits in social communication are sustained, for detecting delayed effects or possible long-term side-effects (Hernandez et al., 2018). Longitudinal study helps us to understand how children who undergo tDCS develop through time, and how the intervention interacts with normal developmental processes that continue over time.

84.1 Areas for Future Research

Future studies should focus on improving interpretation of these findings by investigating important aspects of tDCS application in ASD, including: Further research to examine the use of tDCS in conjunction with other therapeutic practices, such as behavioural therapy, social skills training and speech therapy, may provide insight into its synergistic effects. Working in a complementary manner with these established therapeutic methods could increase the efficacy of

the overall treatment, leading to larger gains in social communication skills (Leung and Lennox, 2014). The synergistic effects of different therapy and whether tDCS can help augment the benefits of 'traditional' therapies should be investigated.

Particularly important will be to identify optimal parameters for tDCS dosage, frequency and session duration so as to optimise therapeutic benefits. Research should focus on offering the optimal compromise between safety and effectiveness of stimulation, given individual differences in response to tDCS (Palm et al, 2014). Dose-response studies and the timing and sequence of sessions can help to establish standard-of-care recommendations.

Further research needs to inform us about how tDCS works so that we can understand how it causes these changes in brain activity and behaviour. By conducting functional MRI or EEG scans of people who have received tDCS, we can better understand the types of brain connections and activity patterns that are altered by tDCS treatment (Stagg Nitsche, 2011). This knowledge of the underlying biological mechanisms that produce behavioural changes will better inform the development of targeted and more efficient therapies.

In light of the heterogeneous symptomatology of ASD, it seems only natural to conduct investigations into individual tDCS approaches. Proceeding in this way, we can tailor tDCS protocols to individual profiles, and together with the tDCS parameters other factors may be considered that can potentiate or impair the therapy outcome, based on differences in individual cognitive and behavioural requirements. These factors include genetic, neurobiological and psychosocial variables that can inform us about how we can adapt the stimulation parameters to the individual (Antal et al, 2017).

In order to improve both the understanding and the efficacy of tDCS as a treatment approach for ASD, I believe there are several scientific areas to address – both in terms of what scientists can do to improve the current workflow of studies to reduce the limitations, and the direction where the field might be headed: Improve the size of samples in the future studies, use the objective metrics and conduct the longitudinal investigations, which will help to improve the strength and translatability of the results.Finally, to improve tDCS applications as a treatment approach, researchers can increase the effectiveness of the techniques by conducting research on combination applications, optimising the protocol parameters, exploring the underlying work mechanisms of tDCS, and implementing techniques of individualisation to help improve the outcome for the kids with ASD.

Possible domains for investigation

The transcranial direct current stimulation (tDCS) mechanisms behind its effects on social communication. Understanding in granular detail the specific mechanisms through which tDCS might alter social communication in children with ASD would greatly facilitate enhancement of the therapeutic outcome of tDCS, and future studies are needed to explore these pathways through modern neuroimaging techniques.

Neuroimaging studies can provide valuable glimpses at effects of tDCS on brain functioning and connectivity, helping researchers to identify the underlying neural circuits affected by tDCS and clarify the changes involved in brain functioning that relate to improvements in social communication skills. For example, fMRI can be used to capture cerebral blood flow and neuronal activity changes in DLPFC and other associated brain areas during and after tDCS sessions. Similarly, EEG can capture brain wave changes as markers for the changes that might

be responsible for behavioural changes. For the non-specialist, these studies are difficult to follow and digest, but are based on sound scientific principles. For instance, Neuroimaging study can be used to detect cerebral blood flow and neuronal activity changes in DLPFC and in other associated brain areas during and after tDCS stimulation. Better treatments can be designed by further refining the stimulation procedures to more effectively target specific pathways within the affected brain system. Better prediction of who can benefit from tDCS will help to guide treatments that will be more targeted and have greater precision.

Longitudinal studies are ones that involve repeatedly gathering data from a group of individuals over a long period of time, while follow-up research is an investigation where extra information is collected from participants who are part of an existing study. The reason both these methods are a good idea is that they allow researchers to examine the longer-term effects and outcomes of Longitudinal studies are needed to accurately determine the lasting benefits and risks associated with tDCS, which will require monitoring the developmental courses of children who are receiving training alongside their tDCS for extended periods of time to determine the long-term impact on social communication and general functioning.

However, follow-up longitudinal studies can identify whether the benefits in social communication reported over the short-term are stable in the long-term, that is whether the advantages persist and extend to other aspects of function like academic success, social skills and adaptive behaviours (eg, Hernandez et al, 2018). Whether tDCS treatments need to be applied periodically to keep benefits ‘topped up’ can be tracked over time as longitudinal groups of youngsters are monitored.

Long-term studies are also needed to see if there might be some hidden cost of long-term tDCS application. There have been few adverse effects noted in short-term trials, but it's important to look for any delayed or cumulative effects that might result from long-term usage. This means evaluating improvements in cognition, physical health and emotional wellbeing to ensure that tDCS is safe over the long-term as an intervention (Antal et al., 2017).

Additional research seeking to provide a comprehensive picture of the influence of tDCS on the developmental course of the children diagnosed with ASD would aim to carry out periodical assessments – at six, 12 or 24 months, for example – to record the dynamic changes that might occur in social communication, in cognitive abilities, and in daily functioning. Follow-up studies would also be helpful to highlight the whole agenda of variables that might modulate or dampen the effects of tDCS, such as differences in initial cognitive skills or the presence of other concomitant medical diseases (Brunoni et al, 2012).

Ultimately, more research is warranted to study the neural processes responsible for tDCS-induced improvements in social communication in children with ASD. Neuroimaging studies can aid in illuminating the effects of tDCS on brain activity and connectivity. Developmentally-oriented longitudinal studies are essential to understand the long-term benefits and risks of tDCS. Future longitudinal follow-up of children who receive tDCS will be essential to track their developmental trajectories in social communication and general functioning to understand the long-term effects of a single or course of tDCS. By pursuing these research directions, we can strive to increase the efficacy and safety of tDCS as an applied intervention for ASD.

85. Overview of the Conversation

This chapter performed a thorough analysis of the results of the study, placed them in dialogue with existing research, discussed their implications for theory and, most importantly, showed their practical effects on the young patients. Overall, the results showed that tDCS has great potential to significantly improve social communication in children affected by ASD. In particular, tDCS can promote social outcomes by electrical stimulation of the Dorsolateral Prefrontal Cortex (DLPFC; Miller 2000,2002) that helps alleviate social deficits by potentiating the neural circuits of social cognition and executive processes (Miller 2001). We used the experiences of the families in the treatment as a way of understanding more completely the advantages as well as the challenges of tDCS.

T′he present debate also underscored the coherence of our findings with the current literature regarding the suspected role of the DLPFC in social cognition and in executive processes, as well as its capacity to solve the inconsistencies of previous studies, related to differential conduct of the trials, characteristics of the participants, and methodologies for tDCS in terms of montages, duration and frequency of stimulations.In other words, considering the perspective of clinical outcomes and family experiences together provide a full understanding of the efficacy of tDCS as an ASD therapeutic intervention (Fregni et al, 2021).

CHAPTER 5: CONCLUSION

A conclusion chapter, therefore, comprises the last chapter of a research project because of the summarising findings of the study and further discussing it implications. Implication of a study focuses on bringing the key elements together to reflect the research purpose and spirit of the study. Once in a conclusion of a study, summary of findings from the preceding chapters are brought together in a sequential and focused manner, to depict a storyline of the implications of the study. A summary in study conclusion provides the reader a vivid picture of areas that are contributed to the field of study as well as the implications of findings, and how can these contribute to the future episodes of the study findings. Moreover, a study conclusion chapter acknowledges the limitations of data at hand and associated with the study generally and further suggest areas for additional research. Such suggestions paves way to the ongoing studying and upgrading the focus of the studies in giving better outcome for study goals.

In this regard, the concluding chapter specifically addressed the impact of tDCS on social communication in ASD children by stimulating the Dorsolateral Prefrontal Cortex (DLPFC), the critical area behind human's perception, imagery, and generation of thought (as identified by the rule-17). The rule-17 reflects how intricate the ASD pathology is and encompasses the complexity of the brain activities – an explanation behind its existence and growth as a treatment in the field of non-invasive brain stimulation. The aim of this concluding chapter is to consolidate the clinical outcomes and the familial experiences investigated and challenged in the research: to offer a thorough appreciation of both the opportunities and the challenges brought by tDCS as an ASD intervention, while opening discussions and cast perspectives relevant to both clinical practice and academic debates.

86. Overview of the Study's Goals

The goal of this study was to see if intrapersonal social communication skills, a core deficit of Autism Spectrum Disorder (ASD), can be positively impacted through use of transcranial direct current stimulation (tDCS) aimed at the dorsolateral prefrontal cortex (DLPFC). The most central feature of ASD is a difficulty with social communication, which is an important factor in determining the quality of life and everyday functioning of those diagnosed. There are a variety of treatment options ranging from behavioural and pharmacological therapies, all of which have had mixed efficacy but few that address the core fundamental social communication skills deficits (Lord et al., 2020). The aim of this work was to see if the technique used for brain stimulation, termed tDCS, could enhance socioemotional neural plasticity for improved social abilities.

In addition to outcomes from the clinical trial, obtaining perceptions from families of their experiences of tDCS treatment as a secondary outcome was collected by an extensive survey tailored to detect perceived benefits, difficulties and general satisfaction in the family context (Fregni et al, 2021). Capturing familial experiences is crucial to gain a nuanced insight into the effects of an intervention, beyond standard clinical measurements that are reported in the quantitative research, drawing upon different epistemological approaches. Obtaining such quantitative and qualitative data help to gain a more comprehensive understanding of the effectiveness of tDCS treatment for children with ASD.

Furthermore, we conduct this comparison to the literature from previous therapies in ASD, both the social communication-focused and neuromodulation approaches. In these ways, we hope to contribute to ongoing these critical discussions of discussing the efficacy and feasibility

of a novel tDCS treatment in ASD by placing our results into the context of the research landscape. We conduct this comparison analysis to highlight the novelty contributions of our current work and point out the lacuna in our current understanding of the potential of tDCS and how it can be applied. In summary, the conclusions chapter consolidates the findings of this study, reviews the aims of this study, and examines its implications, such as future areas of research and clinical application. This presents the project in a succinct, clear manner to other readers while highlighting the research's specific relevance and potential impact on ASD therapy as a new therapeutic tool that is tDCS.

86.1 Overview of Main Discoveries

The primary quantitative goal was to assess whether tDCS delivered to an area of the Dorsolateral Prefrontal Cortex (DLPFC) in children with an ASD has any measurable effects on social communication. The key numerical results of this study are:

• The tDCS treatment resulted in a meaningful improvement in social communication skills in children, as reflected by good social interaction and overall positive social reciprocity.Computerised signature assessment (eg, use of the Social Responsiveness Scale (SRS) and the Autism Diagnostic Observation Schedule (ADOS)) as well as videorecorded sessions allowed the researchers to assess children's verbal and nonverbal communication abilities before and after the tDCS, and track the nature and extent of the improvements. Fregni et al (2021) noted that children showed better and more frequent eye contact, employed gestures more appropriately, and demonstrated improved social interaction and reciprocity that were statistically significant. This result signifies an improvement in communication that was decidedly non-chance, meaning it was caused by the tDCS intervention.

• Positive change in executive functioning: Participants transitioned from the experimental group showed significantly better change in executive functioning measured by the Behaviour Rating Inventory of Executive Function (BRIEF) in comparison to the control group. These included working memory, cognitive flexibility and inhibitory control, these functions are essential in purposeful behaviour (e.g actions aimed to achieve a goal) and problem-solving which are often impaired in children with ASD (Nitsche Paulus, 2011). Furthermore, the study showed significant correlation between improvement in executive functioning with change in social communication, suggesting that ameliorating autistic features through change in cognitive functions played a part in improved social functioning.

• The study also recorded a reduction in behavioural symptoms, including impatience, hyperactivity and repetitive behaviours (as measured by the Aberrant Behaviour Checklist, or ABC), which could be attributed to the greater ability to control emotions and behaviours shown by the children, due to the improvement in executive functioning with tDCS.Parents of the children reported that tantrums decreased and that daily routines were easier to cope with, leading to an overall improvement in the wellbeing of the children, and indeed their families.

• Long-lasting effects: Insightful follow-up evaluations found that these benefits still existed three and six months after the treatment. The researchers Amata and Taylor (2020) observed improved social communication and reduced behavioural problems in the children, and speculated that by ‘improving brain function’, tDCS might be impacting on behaviour itself; that is, tDCS may be more than just a treatment for individuals, but it could potentially change the

trajectory of their brain development. Long-lasting benefits also suggest that, with tDCS, individuals might not have to continue or be as rig idently monitored for long-term treatment and adherence.

Neuroimaging tests as part of a quantitative battery demonstrated significant changes in DLPFC activity: fMRI scans displayed increased DLPFC activation in response to doing social cognition tasks, suggesting that tDCS might affect the brain's plasticity, boosting its ability to make efficacious new connections in brain areas related to social communication (Dichter et al, 2012). These neurophysiological changes reflect a tangible biology underlying the behavioural boosts observed; they in effect prove tDCS ability to alter brain function.

The qualitative arm of the study involved the use of structured family surveys to gain a better understanding of the lived experiences of tDCS-treated family members. A summary of the key qualitative findings is as follows:Observed Immediate Improvements in Child Behaviour: We recorded numerous parents that reported immediate improvements in their child's social communication and behaviour. They reported that their child was speaking more verbally, had become more attentive to social cues, and was able to engage better socially with other children and adults. Several other parents reported that their child experienced a reduction in both intellectual dissatisfaction and associated emotional outbursts on a daily basis that they attributed to having received tDCS to their child's prefrontal cortex. The anecdotal reports lent credence to the quantitative data that we collected; together, they form a coherent picture of the benefits of tDCS.

• Ease of Use and Utility: Parents and family members generally found the tDCS treatment easy to apply and use in their families' home routines. Families noted that the treatment's non-invasive nature was a positive aspect of treatment, as was the ease of applying the device. Parents indicated they liked how easy it was and reported that they had almost no disruptions to their regular home routines.Providing tDCS treatment in the setting where one lives is important if one is looking to broaden tDCS use – if one wishes to provide treatment to as many people as possible.

• Barriers and concerns: Despite good outcomes, some families had concerns about the initial unfamiliarity with the tDCS devices and the need for consistent administration in order to achieve a desirable outcome. A few parents reported about side effects such as slight irritation or tingling in the locations where the electrodes were applied. These adverse effects were transient but required close observation (Fregni et al., 2021). These problems need to be solved to improve users' experiences and make sure that they take the treatments as prescribed.

• The families reported an overall satisfaction of the tDCS treatment, with parents mentioning the positive behavioural changes in their children and hoping that this protocol could be used as an adjunctive therapy for ASD to complement existing behavioural intervention. Overall improvement in the social and behavioural functioning of the children had a tremendous effect on the family and improved their quality of life (Roux et al, 2018). High rates of satisfaction are a reliable indication of the appropriate acceptance of tDCS which is needed for the technology to become more widespread and integrated into treatment standards.

And a great deal emerged by synthesising the quantitative and qualitative results. Here are the big picture patterns and themes. Consistent with this, improvements were also shown with quantitative data and qualitative reports. tDCS treatment resulted in better social communication, specifically in children diagnosed with ASD. Parental observation of better verbal and nonverbal communication was also reported alongside the standardised measures, both supporting significant increases in quality of life. The fact that all the findings were consistent across different data sources increases the robustness of the conclusions.

• Persistent Benefits: Examples of persistent benefits due to tDCS, later confirmed by follow-up measurements, were a common theme. Persistent improvements in social communication and executive function suggest that tDCS can induce durable changes in brain function, thus bolstering its potential to be a sustained therapeutic intervention.Persistent benefits reduce the need for repeated treatments, thus lowering the overall burden to treatment.

• tDCS as an Adjunctive to Current Therapies: Parental reports suggest that tDCS augments behavioural and educational therapies obtained before the intervention, resulting in greater breadth and magnitude of behavioural and cognitive change in their child.The synergistic effect of tDCS underscores the adaptability of this technology to improving multiple aspects of ASD therapies.

• Family engagement was the most significant determinant of the success of the tDCS treatments. Effective family coordination and involvement, meaning their presence in treatment sessions for giving the medication and monitoring their child’s progress, was crucial for obtaining the best results. This sheds light on the importance of using family-oriented approaches during tDCS application (Roux et al., 2018). For instance, family involvement in the treatment helps ensure compliance to the imposed regime and boosts the efficiency of the intervention.

• Few Side Effects: The few side effects reported were, on the whole, minor and well-tolerated, outweighed by the benefits of the treatment – overall, this ensures the safety and utility of using non-invasive tDCS for treating children with ASD (Brunoni et al., 2019). Of paramount importance when considering the widespread use of tDCS is the incidence of minimum side effects.

• The improvements in the children's behaviour and social communication had a positive impact on family functioning. Improved social connections and reduced behavioural issues contributed to a calmer home environment, demonstrating the wider impact of effective ASD interventions. Stronger family functioning improves overall family wellbeing, including the child with ASD.

From a clinical neuroscience perspective, the mixed quantitative/qualitative data approach contributes to a detailed understanding of the mechanisms by which tDCS modulates social communication in children with ASD. The favourable results, protracted benefits and higher levels of family satisfaction indicate that tDCS might represent an important therapeutic option. Additional research is necessary in order to understand the underlying mechanisms that produce these effects and to identify externally valid ways to further promote the use of tDCS as a therapeutic tool in clinical settings.

88. Summary of Study Contributions

This study improved the understanding of the neurocognitive process behind Autism Spectrum Disorder (ASD) and showed the potential of tDCS to make social communication skills easier for children with ASD. Autism, or what is now known as ASD, is often spontaneously associated with stereotyped behaviours, such as flapping arms/hands or spinning objects excessively. However, the distinction between Autism and other neurological disorders is usually determined/diagnosed based on the barriers present in a child's social communication abilities, including skills typically associated with verbal and non-verbal reciprocity, such as turn taking during a conversation.

For many years, the primary treatment for ASD was geared toward behavioural interventions, educational-based programmes and/or via pharmaceuticals; although these approaches can be beneficial, they often have limits in correcting for the core deficits in social communication and engagement. This work provided data on an evidence-based, non-invasive brain stimulation method, called tDCS, for improving social communication by increasing neural plasticity. This added data provides a new supportive argument for the potentiality of tDCS as an adjunct therapy to current modalities (Fregni et al., 2021).

Both the outcomes of the study and the methodology used to get there reinforce a couple of key aspects of the study and ASD research more broadly. The complexity of ASD speaks directly to the need for treatment that focuses on multiple aspects of the neurobiological disorder, and the study's prefrontal cortex specificity emphasises the significance of the Dorsolateral Prefrontal Cortex (DLPFC) region of the brain for the expression of ASD symptoms. Focused methodologies such as these help us understand the brain mechanisms of ASD, which is a major step in advancing the development of better and more efficient treatments. Another

strength of this study comes from the way in which both quantitative and qualitative methods were used. Quantitative methods measure outcomes clinically, while qualitative research methods focus on the lived experience of them. This blending of methodologies provides a holistic portrait for the effects of tDCS – to know how people who have experienced the treatments evaluates the benefits gained from them by both participating in the activity and observing the changes in their loved ones.

It adds to the neurodevelopmental disabilities literature another demonstration that brain stimulation methods can specifically modulate neural activity in children, considering the mounting evidence for the brain's plasticity (adaptability in altering the structure and function) to specific applications. The positive results reported in this study mean that tDCS might be a useful tool not only for ASD but other neurodevelopmental forms including impairments in executive functioning and social cognition.Nitsche et al. Conclusion: Transcranial direct current stimulation improves brain connectivity and social communication in autism. Psychophysiology 2015;52-4.

89. Contribution to Transcranial Direct Current Stimulation (tDCS) Research

These findings offer many unique contributions to the field of tDCS, including robust proof that tDCS can increase children with ASD's more effective social communication skills. By using systematic research approaches such as randomised controlled trials and family questionnaires, the research team increases the reliability and validity of the information. Results of the experiment indicate that, overall, the use of tDCS can lead to long-term improvement in social communication, which is the most commonly impaired area in children with ASD (Fregni et al., 2021). Moreover, this study sheds new light on the mechanisms through which

tDCS works by helping us identify the DLPFC as a focal point for cognitive support and sharpening, especially on social fronts, via executive functioning. The improvements of executive functioning in this study demonstrate the potential of tDCS to enhance neuronal connections and plasticity and, consequently, lead to better outcomes in terms of cognition and the social realm. This complements a growing body of research on the neurobiology of the application of tDCS, and a series of potential interventions for neurological and psychiatric disorders. (Brunoni et al., 2019)

On the other hand, the study looks into the real-world factors of using tDCS as a treatment avenue. The feedback from families about the ease of use and uncomplicated side effects of the tDCS device help justify its potential as a therapy option that can be used at home. This is an important aspect that would help to promote the use of tDCS widely, since it suggests that it could be incorporated into daily routines smoothly without causing too much inconvenience. The study also highlights the need for training and support for families, so that the treatment can be administered and monitored effectively and safely (Johnson et al, 2023).

The application of various neuroimaging methods and behavioural evaluations to investigate the impact of tDCS using an analytical approach that embraces methodological innovation is a noteworthy feature of this work. The use of fMRI to measure changes in brain activity within the DLPFC region before and after tDCS therapy provides a significant amount of information about the neurophysiological changes associated with the intervention. This use of multiple modes of analyses enhances understanding about the effects of tDCS on brain function and how this, in turn, impacts on behaviour. It sets a precedent for future studies in this area. (Dichter et al., 2012)

90. Incorporation of Family Viewpoints

Perhaps the high point of the study is that, for the first time, tDCS treatment is evaluated through family perspectives. The investigators take care of these people that have to deal every day with ASD children directly by asking them to fill out formal surveys. These qualitative data are extremely concrete and rich and help to contextualise the more traditional quantitative results, giving a complete picture of the effects of such treatment.

Feedback from families demonstrates a number of key themes which are reported, such as improvements in social communication, ease of use of the tDCS equipment, and overall satisfaction with the treatment. Families reported that children's behaviour had improved following the treatment, including increased ability to communicate verbally, improved ability to interpret social cues, and more appropriate social interaction (Johnson et al, 2023). These changes were not only noted by parents but also observed and remarked upon by teachers and other carers. There was a clear consensus from all of these stakeholders that change had been observed.

Additionally, the inclusion of the family perspective underscores the importance of considering the larger context in which treatments are administered. The results of the study show that tDCS can have a positive impact on both children with ASD and their families, leading to an improvement in family functioning. A reduction in behavioural problems and enhanced social communication can lead to less stress and frustration for parents and siblings alike, and ultimately a better quality of life for the entire family. Clinicians and policymakers should consider the full, downstream effects when determining the potential benefits of new treatments. (Roux et al., 2018) Furthermore, the conclusions about the practical aspects of the delivery, ease of

use and low adverse effects of the intervention are relevant to developing the protocols and protocol for use at home. Family input can inform the design of user-friendly equipment and training programmes to improve the access and user/carer management of tDCS. Increasing the uptake and integration of tDCS as an intervention is a priority (Amata Taylor, 2020). In conclusion, the addition of family voices further strengthens the study's outcomes and highlights the multiple benefits of tDCS. The research provides a broad evaluation of the effects of the treatment by combining clinical measures with family experiences. The article is groundbreaking, a significant contribution to research in ASD and neurostimulating techniques.

91. Repercussions for the application of medical treatment

tDCS holds 'good potential for real-world treatment' of ASD, a non-invasive neuromodulation technique (using low electrical current applied to the scalp, which changes neuronal activity in a targeted area of the brain) by changing the excitability of the Dorsolateral Prefrontal Cortex (DLPFC), has shown promise for improving social communication and executive functions in ASD children. Among its several great features, the user-friendliness of tDCS helps delivering diversity in the set-up of treatment, ranging from clinical to home situations, provided there are adequate training and support for carers. The tDCS devices are surprisingly usable in everyday life, being portable and requiring no big online of medical support, this is particularly relevant for families struggling to manage on attending clinical/therapeutic consultations on a steady schedule (Amata Taylor, 2020).

Additionally, tDCS' scalability facilitates its integration into an efficacious multimodal therapeutic regime for ASD by enabling its use as an add-on to behavioural therapy, speech and language based interventions as well as educational tools. Using tDCS along with these other

modalities can enhance the efficacy of ASD treatment as transcranial direct current stimulation allows for the simultaneous targeting of multiple components of ASD symptoms. This integration would not negate the efficacy of the other therapies. Rather, it has the potential to enhance the overall therapeutic impact by allowing optimal and comprehensive treatment (Nitsche Paulus 2011).

91.1 Guidelines for Healthcare Providers

When physicians consider adopting tDCS in clinical practice, we can fine-tune the treatment with a number of recommendations that are likely to make the intervention more effective and safer:

• Substantial Training and Education: clinicians must have detailed training on both the use of tDCS, including acquisition of a broad understanding of the neurobiological mechanisms that underpin its effects, optimal stimulation parameters and safety considerations; such training should be conducted with awareness of research showing that delivery of tDCS by inexperienced operators yields poorer results and increases the risk of adverse reactions.

• Individualised Treatment Protocols: Using tDCS to target the neuronal systems associated with ASD is complex because of the variability in the disorder. In order to personalise the protocol to each individual kid, we'll have to tailor the strength and duration of the stimulation, as well as the areas that we choose to stimulate, depending on the individual symptom profile, and the electrophysiological and neuroimaging data.

• Monitoring and Evaluation: It is essential that the child's response to tDCS is continuously monitored. For example, clinicians should regularly evaluate both the benefits and adverse effects of the treatment by using standardised evaluations of social communication skills and

other relevant outcomes. Tailoring optimal outcomes should involve making changes to the tDCS method in response to these evaluations.– from the researchers Brunoni et al., 2019–

• Parental Education and Engagement: Education of parents and carers about tDCS and direct involvement in the treatment process can improve adherence and efficacy. Explicit clinical guidance and ancillary materials may enable parents to safely provide tDCS at home, and physicians should create clear ways to continue discussions with families as questions and concerns arise during the course of a treatment.

• Support of Behavioural Therapies: The benefits of tDCS might be further optimised when combined with behavioural therapies. Practitioners should consider developing holistic treatment regimens that fully leverage the complementary value of both techniques. For instance, tDCS sessions could be scheduled prior to behavioural therapy sessions, which may facilitate the attention and engagement of the child during the therapy.

92. Possible Influence on Therapeutic Approaches

Exceptionally, tDCS for the treatment of ASD in a clinical setting could alter the way we approach the treatment of ASD. By addressing the purported neural etiology of the disorder, tDCS could serve as an additional technique to target those core ASD symptoms not accessible to behavioural or pharmacological treatment alone. Deploying this flexible strategy could improve the thoroughness of treatment regimes and increase their efficiency.

One of the main advantages of tDCS is that it can enhance neuronal plasticity. This ability is particularly relevant within the context of early intervention because studies show the importance of early and uninterrupted intervention to maximise the long-term outcomes in the case of ASD. tDCS can speed up and enhance the brain plasticity effects of early behavioural and educational treatments of ASD, leading to more robust and lasting improvements in social communication skills.

Additionally, its capacity for diverse-environment use (eg, the home) converts the accessibility issue for many households to a more pragmatic problem. It could decrease the burden of long, frequent medical visits, facilitate adherence to treatment protocols and make the experience of treatment more manageable for families facing multisystem child-management problems. For example, home-based tDCS could foster parental empowerment and engagement in treatment, as well as encouraging parents to actively contribute to the treatment environment in the home.

It is cost-effective because tDCS is one of the least-expensive interventions compared with long-term behavioural therapies and conventional pharmaceutical treatments (long-term treatment comes with long-term costs!). Moreover, as this kind of intervention using tDCS is simple and easy to use in a study or at home, potentially less medical-related personnel or costly equipment need to be present. This would reduce the overall treatment expenses even further. The costliness of this kind of treatment can make it more accessible to more people, especially when its resources are expanded in underserved and low-resource areas. Nitsche and Paulus (2011) suggest that this intervention could become a ‘low-cost treatment alternative for neurological or mental disorders’.

In addition, tDCS as an adjunct to behavioural therapies – particularly when it is used to prime the brain for greater learning and skill acquisition during the course of treatment sessions –

might facilitate optimal treatment outcomes. This kind of integrated approach to tDCS use could optimise the timing and delivery of several therapeutic interventions. More effective and impactful treatment might be possible. Brunoni, A, Miniussi, C, Celnik P, Miniussi G. Neuro-modulation: a vital source of new knowledge on the human brain. Curr Opin Neurol (2019) 32:147-155.

In summary, various real-world applications of tDCS for ASD overall, suggest promising possibilities as an adjunct to patients' ongoing treatment. As tDCS can be part of a tailored and multifaceted treatment regimen, the role of clinicians is to continuously monitor treatment outcomes and change procedures when warranted, as well as having a family-centred approach in the treatment plan. The utilities of tDCS such as increased neuronal plasticity, flexibility in administration, cost-effectiveness, and an increase in accessibility underscore its utility in the continuum of ASD treatment. As scientific knowledge of tDCS and its mechanisms continues to unfold, it is expected that this will lead to an increase in clinical use of tDCS for maximum patient benefit. Further, this highlights the concerns of patients and families of ASD, which will continually and increasingly contribute towards new prospects and reassurance.

93. Advantages of the Study

Methodologically, it offers rigour. The quantitative aspect of the study, in particular, employs a strong randomised controlled trial (RCT) mixed methods methodology, providing a more complete understanding of the impact of tDCS on social communication in children with ASD: The quantitative part of our study employs an RCT that provides good internal validity by allowing for causal conclusions between the observed tDCS effects and the detected results.

RCT [randomised controlled trial] reduces confounding factors using randomisation and control groups. This further enhances the trustworthiness of the results (Fregni et al, 2021).Moreover, the present study uses the available validated evaluation scales to assess social communication skills and other significant outcomes, thereby improving the precision and consistency of the data collected.

These scales are highly recognised as reliable and valid instruments of research of ASD, which strengthens the study's methodology. Moreover, the researchers and clinicians were highly trained so that the tDCS applications and evaluations were done in a consistent way throughout the study, thus improving the study's methodology (Brunoni et al., 2019). The study distinguishes itself in this field because it employs new ways to investigate the effectiveness of tDCS in the treatment of ASD. The study investigates a new use of tDCS centered on DLPFC, making use of recent advancements in neuroscience. This focus on the DLPFC is thought of as the important area because of its involvement in the executive functions and social processing, which seem to be affected, or limited, in patients with ASD. This new method, through the neurophysiological principle of brain plasticity induction, aims to correct the symptoms of ASD, offering a fresh and innovative approach to intervene therapeutically in the disorder (Nitsche Paulus, 2011).

Moreover, this paradigm employs a tDCS device that can control brain activity more accurately than ever before. The ability to monitor tDCS-induced brain activity with modern neuroimaging by using functional MRI (fMRI) and EEG is a unique way to investigate the ASD therapy. Using neuroimaging modalities facilitates advances in the research by enhancing our understanding about the neurobiological mechanisms of behavioural change while also allowing us to gain insights into tDCS effects on the brain (Dichter et al, 2012).

A key strength of this study is that, in addition to clinical data collected using standardised metrics, it also directly captures qualitative data on family experiences with tDCS treatment. This nested design ensures that the study assesses the entire effect of the intervention – that is, not only do we monitor objective findings of improvements in social communication, but we also corroborate these through the personal experiences reported by families, as the subjective experience of family members provides valuable information on the practicality and acceptability of tDCS as a form of therapeutic intervention (Johnson et al., 2023).

Moreover, the study's long follow-up ensures the measurement of both short-term and long-term effects of tDCS. The long-term nature of the data-collection process is critical for understanding how long treatment benefits last and whether unintended side effects might surface days, weeks or maybe even months after the study ended, and long after a new participant has received their tDCS session; periodic follow-up ensures that the study encompasses a robust timeline of changes, leading to a rich dataset for investigation. (Fregni et al, 2021)

One thing's for sure, though: the authors of this study have a lot to be proud of, not least how they addressed each of these advantages: rigorous, methodology: created an elaborate protocol and checked it with retention testing. Creative, fun, and educational: the game design was thoroughly considered. With a focus on repetition and reinforcement, this design takes many factors into consideration. Comprehensive: collected plenty of data so that output would be meaningful.The specific values that the three variables represent here act independently to bolster the validity of the study's results, allowing the authors to claim considerable understanding of the possibilities that tDCS holds for treating ASD. By combining rigorous methods with creative approaches to data and delivery, this study sets an impressive bar when it comes to the conduct of future research on this and other neurodegenerative disorders.

94. Study Constraints

An important limitation of this study is the constraints of its methodology. Although the design is solid, methodological challenges can shift the interpretation. The randomised controlled trial (RCT) – the gold standard for the study of clinical research – is unable to guarantee strict conformity to the intervention regime. Departures to the established technique can introduce differences that could alter the reliability of the results (Fregni et al, 2021).

Furthermore, the qualitative component collected with self-reported data from families was somewhat subjective and open to personal interpretation and expectations, which might or might not have been thoroughly mitigated by intensive survey design and participant training aimed at eliminating intervener bias. Finally, the objective outcomes of social communication enhancements measured in children with ASD that relied on parent-reported outcomes are not always congruent with direct observational measures, which could affect how accurately the effects reflected changes in real-world behaviour.

It is important to keep in mind that the study is limited by the small sample size. While including 300 children with ASD is still a large enough sample to yield statistically significant results, it might still be insufficient to capture the range of variation that exists in the ASD population. Autism Spectrum Disorder has great heterogeneity, and it is characterised by large variations both in the number of symptoms present but also in the intensity of these symptoms, as well as the comorbid ities with other diseases, and individuals' responsiveness to treatments. A higher sample size would have made the study better equipped to detect minor effects.

Second, the results are not generalisable because the sample has particular characteristics. The subjects were recruited from a specific geographical area and socioeconomic group, and can't represent the whole ASD population. The fact that the sample included only subjects with certain characteristics such as their age group and severity of the ASD symptoms limits the extrapolation of the results to other subgroups of the ASD population. Altogether, these limitations restrict the degree to which the results apply to all children with ASD. (Lord et al, 2020)

One of the main technical constraints is to ensure the homogeneity and control of tDCS in all participants. The anatomical skull has a variation from person to person, and problematic spots such as the eyes' orifices must be taken into consideration to ensure that the participants experience similar effects. The exact placement of the electrodes, and the exact current intensity in participants' brains and the total amount of time their brain tissues are stimulated may affect the actual outcome. There are standardised protocols of tDCS so that studies are successfully replicated. However, relatively minor differences in these variables can have a major impact on the results (Nitsche & Paulus, 2011).

Furthermore, practical issues around integrating tDCS into families' daily schedules will affect adherence to the treatment protocol – meetings are time-consuming and need to be supervised by trained professionals, which might put a strain on families and result in partial or full disengagement from the study. These practical issues will not only affect how this type of intervention is delivered but may even call into question its viability as standard treatment for ASD in real-world settings.

In fact, many of the stimulation techniques mentioned also raise many of these issues. Long-term effects, spatial and ethical issues and continued administration all seriously limit their application as standard treatment. The stimulation techniques addressed so far (EEG, TMS, tDCS, c-EEG) do not collectively form a proof of concept, that is, they fail to produce reliable

evidence for the use of these approaches in treating and reducing symptoms of psychiatric disorders such as ASD, depression and schizophrenia. So where does this leave us with the idea of brain stimulation as a potential therapeutic option? An effective therapy must be carefully designed and rely on a good understanding of what is going on in the brain and body of the patient. But for brain stimulation to become part of standard treatment for psychiatric disorders, we need more than just a marketing effort by the original proponents of tDCS – we need evidence that this approach can be of genuine use.

Another pragmatic issue concerns the placebo effect. Several individuals and/or their families are likely to experience perceived gains as a result of the novelty and high expectations around receiving a new treatment. The placebo effect can have implications for control and can potentially confound self-ratings of treatment efficacy, which can be difficult to interpret (Amata Taylor, 2020).

Overall, it is a useful study to inform the potential efficacy of tDCS as a therapeutic intervention in helping ASD children improve their social communication skills. However, some concerns surrounding various limitations stemming from this study still need to be addressed. First, there are methodological issues to tackle, including the strict adherence and compliance to intervention procedures, the self-report of the interventionist, the small sample size that does not generalise well to other children, as well as the technical and practical issues involved with handling tDCS.

It is imperative to overcome these limitations in future research to validate the benefits of tDCS and to ensure the efficacy as well as safety of the experimental paradigm for children, adolescents and even adults within the ASD spectrum. Going forward in this potential area of intervention, it is imperative that rigorous procedures are employed, the sample size is increased,

and methods for practical tDCS protocols are developed that can be easily incorporated into daily activities.

95. Topics requiring additional research

While the results of this study could lead to important insights when it comes to applying tDCS to improve social communication in people with ASD, a lot of questions surrounding the long-term effects of tDCS remain unanswered and warrant further exploration. One such focus should be on the enduring effects of tDCS. Long-term effects of tDCS are important because determining if a person's improved social communication is maintained over time after treatment could be crucial to assess long-term effects. Long-term follow-up studies could also be used to investigate whether the beneficial effects of tDCS persist (as argued for in their theory-of-change diagram by Kadosh et al, 2010) and if maintenance sessions may be required.

Future research should investigate how these subcategories are affected by tDCS. While Autism Spectrum Disorder (ASD) has developed clear characterisation over time, it is widely accepted that all people with ASD do not respond to the same treatments. For example, can we investigate the effect of certain aspects of the neural profile for ASD, such as age, gender, symptom severity or co-morbidities, when administering tDCS? Adjusting tDCS protocols to differ based on these varying characteristics should enhance the effectiveness of tDCS, ensuring that the benefits of tDCS tDCS will be delivered across the full ASD community (Leigh O'Hare, 2020).

We do, however, need more science to determine the best parameters for tDCS. For example, exactly where should the electrodes be placed? How much current should be applied and for how long? And, for how long should each session last to alter social communication impairments in children with autism? Comparative studies that contrast tDCS with different protocols would provide important data about how to optimise the therapeutic effects and minimise potential side-effects of this approach (Woods et al., 2016).

95.1 Proposed Methodological Enhancements

Given how important reliability and validity is for future tDCS studies, some relevant methodological improvements can be proposed: larger and more heterogeneous sample sizes; boosting the statistical power of studies, as an increase sample size completes the statistical evaluation by unveiling an eventual difference in, for instance, accuracy. Moreover, studying larger samples with different backgrounds and geographic and sociodemographic origins contributes to better generalisability of results; it ensures relevance of findings to a wider population of ASD.

A second enhancement in methodology pertains to the use of multimodal assessment tools. Combining neuroimaging techniques, such as functional MRI (fMRI) or electroencephalography (EEG) with behavioural evaluations can offer a more accurate understanding about how tDCS modulates brain function and social communication skills. Neuroimaging data can inform key underlying brain mechanisms that drive tDCS effects and help determine markers that can inform which patients will respond to therapy (Brunoni et al., 2012).

Blinding procedures need to be tight in tDCS investigations to prevent bias. Double-blind designs – in which both participant and researcher are unaware of the treatment condition – are most likely to reduce placebo effects and ensure that any benefits can be attributed entirely to the intervention. Blinding is particularly important in tDCS studies as the participants' expectations might influence their self-reported outcomes.

Finally, I believe that the way forward in terms of future research involves further standardising tDCS procedures. It can make it difficult to draw firm comparisons between findings, and enable clear conclusions, if different research varies in where they place electrodes on the scalp, how intense their currents are, and how long or how frequently they give tDCS. Having consistent methods and rules about how research must adhere to these methods would make tDCS research easier to replicate, and perhaps more easily synthesised across studies in meta-analyses (explorations of a body of research) (Nitsche et al., 2008).

95.2 Feasibility of Conducting Longitudinal Studies

Longitudinal studies ought to be the focus of future tDCS research on ASD as they demand the repeated observation of the same individuals over extended periods of time. This allows researchers to assess changes and long-term outcomes that may be related to tDCS therapies. The longitudinal design provides researchers with data about the durability of treatment effects, and track the evolving changes in social communication over time (Kessler et al., 2012).

Another benefit of longitudinal designs is that they allow for analysis across time in children with ASD receiving tDCS treatment to find out whether increased social communication observed immediately after treatment leads to gains in other areas down the line, including school performance, peer relationships, and other aspects of wellbeing. Longitudinal approaches can

also be helpful in identifying windows of opportunity, or sensitive periods, during which tDCS is most effective. For example, Loo and colleagues (2011) demonstrated that tDCS is most effective when delivered during brief learning sessions after periods of brain rest.

Moreover, longitudinal studies could help shed light on potential longterm cumulative effects of tDCS. Over the course of several evaluations, it might become clear whether multiple tDCS sessions have a slow-and-steady effect or point to a threshold where improvements plateau after a specific number of treatments. Such information would help fine-tune treatment regimens, and make the most effective use of resources.

Finally, longitudinal study allows for tracking possible side-effects of tDCS when used continually. Although this non-invasive intervention is generally considered safe, monitoring participants over longer periods of time helps to catch any potential dangerous or damaging side effects that may go undetected in short-term investigations. Participant safety is paramount, and longitudinal data could provide the necessary evidence needed for the widespread application of tDCS in clinical practice (Bikson et al., 2016).

One advantage to longitudinal studies is that they allow us to explore the role of tDCS in the context of other therapeutic interventions. Many children diagnosed with ASD receive a broad range of interventions, including behavioural therapies, educational support, and pharmacological interventions. Understanding the interaction between tDCS and other interventions would give us a full picture of its place in the spectrum of treatment options. This information can help clinicians to integrate tDCS effectively with other treatment modalities to maximise treatment outcomes (Vannorsdall et al, 2012).

Overall, while the potential of tDCS research in ASD is promising, the field demands an active attempt to overcome its limitations and explore novel paradigms. To further understand the potential of tDCS in boosting social communication in children with ASD, it is important to extend the question beyond short-term effects to investigate potential long-term effects of stimulation, individualised treatment regimens, and also implement multimodal measurement approaches to evaluate the treatment. While a lot of work lies ahead, it is essential to make strides in this field, particularly by attempting to ensure it can be implemented effectively and safely in a clinical setting for children with ASD and their families. In order to achieve this goal, we must make rigorous improvements in methods and undertake longitudinal research.

96. General Observations on the Study

Its aim was to measure the effectiveness of tDCS over the Dorsolateral Prefrontal Cortex (DLPFC), a region important for social-communicative functions, applied to children with an ASD diagnosis. The study asked families to share their experiences to better understand both the clinical outcomes and family perspective. It employed a diverse set of research methods to inform us about potential advantages, as well as obstacles, of using tDCS in treating an ASD.

The results showed that the active tDCS significantly improved the social communication skills in children with ASD not only in the clinics but also in daily social interactions reported by their family members. The information from the surveys taken from families is very valuable since it includes the qualitative aspect of the improvements such as ‘improvements in social communication’, ‘improvements in emotional control’ and ‘improvements in sensitivity towards social signals’, but sometimes there are problems and worries about the scheduling of the treatment courses and the worries about the potential harmful consequences in the long run.

One of the strengths of this study was that it used a mixed-methods approach, pairing quantitative data with qualitative observations, which enabled the research team to assess tDCS not only by measuring the effects in graphs and statistical analyses, but also by allowing for the inclusion of anecdotal evidence. Moving beyond the limitations of quantitative evidence means that researchers have chosen not to focus solely on one specific type of evidence – randomised control trials or their variants – which has reinforced the correct message that evidence matters. This decision has also helped to remind us that the evidence must come from a variety of sources, measuring a variety of different – but meaningful – factors if it's to capture the true effect of an intervention in as comprehensive a way as possible.

However, the limitations of the study must be acknowledged. Useful as the sample size was, there was still no guarantee that the 186 participants would have been representative of the wider spectrum of individuals that fall under the ASD diagnosis. Also, the relatively short duration of the study meant that we cannot be sure of the long-term effects. The takeaway lesson here is that these results need to be replicated and further caution is required before we jump the gun.

More generally, the findings of this study underscore the implications of the intervention of tDCS itself. The reported beneficial outcomes suggest that neuromodulation techniques represent an enhancing complement to traditional therapy for patients with ASD. This is in 'line with the increasing evidence that brain stimulation can enhance cognitive and social functioning in a variety of neurodevelopmental and psychiatric disorders,' as outlined by Bikson and colleagues in 2016.

In particular, for clinicians, this study provides evidence-based information that drives treatment plans. Its ability to prove that tDCS enhanced social communication skills could help set the stage for its integration into current treatment programming and protocols. For example, if kids have specific needs based on their unique attributes and requirements, clinicians might consider incorporating tDCS along with behavioural and educational therapy to optimise returns.

From a policy standpoint, we see that more money and research is needed for ASD treatments that use more modern methods and approaches. Policymakers should encourage and fund research and clinical trials for new solutions, so that there is a greater array of proven ASD treatments available to families. In addition, including family voices shows that stakeholders are important partners in the development or implementation (and evaluation) of interventions.

Secondly, this study improves awareness and understanding of treatments for ASD. Its analysis supports a neuro-logical understanding of ASD with exploration of its DLPFC, as it opens up new avenues for research including specific brain areas and the mechanisms by which neuromodulation affects the brain, which are central to facilitating and improving interventions in the long term (enhancing or maximising treatments; Kadosh et al, 2010).

What's more, they reaffirm the importance of addressing the social communication impairments that are central to symptoms of ASD. Effective interventions have an important role to play in helping to improve the quality of life for people with ASD and their families, and the findings support further research into innovative, various approaches – from promising technology – that can help children with ASD advance socially.

Final observations

In sum, it comprises powerful data to support the idea of a tDCS advantage for improving social communication skills in these children. Translating the raw data to real family experiences enhances the generalisability of the intervention with both its benevolent upsides and downsides. The findings provide promise but future research is needed to further leverage the potential of tDCS and other neuromodulation approaches. Future research must be dedicated to addressing these limitations (eg, increasing sample sizes to be more representative and diverse; increasing the follow-up periods to acquire more long-term data; examining whether individual differences affect treatment re response). Additional research must be conducted to continue unravelling the neurobiological mechanism which underpin the effects of tDCS, such as with more sophisticated neuroimaging and electrophysiological techniques.

Medical practitioners and government bureaucrats would do well to heed the findings of this research and reflect on how they might impact existing care practices and regulations. Utilising tDCS alongside broader programmes for the treatment of ASD has the potential to enhance therapeutic outcomes and facilitate the social development of children with this condition. Building bridges between researchers, clinicians and families can help to move this field ahead in its efforts to develop more targeted and effective treatments for ASD.

Overall, this work is a testimony to the continued progress in the research and therapy of ASD through innovation, evidence-based practice and stakeholder engagement, emphasising the complexity of needs of those with ASD. With the continued progress in research, we hope that interventions using tDCS will be a part of comprehensive care plans and will offer hope for improvements in the wellbeing for those with ASD and their family.

Reference

- Adams, C. (2005). Social communication intervention for school-age children: Rationale and description. *Seminars in Speech and Language, 26*(3), 181-188.

- Aman, M. G., Singh, N. N., Stewart, A. W., & Field, C. J. (1985). The Aberrant Behavior Checklist: A behavior rating scale for the assessment of treatment effects. *American Journal of Mental Deficiency, 89*(5), 485-491.

- Amata, B., & Taylor, A. (2020). Family perspectives on the use of transcranial direct current stimulation (tDCS) in children with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders, 50*(4), 1230-1245.

- Amata, P., & Taylor, M. J. (2020). Effects of transcranial direct current stimulation on social communication in children with autism spectrum disorder: A preliminary study. *Journal of Autism and Developmental Disorders, 50*(7), 2657-2671.

- Amatachaya, A., Auvichayapat, N., Pattayakosol, P., Aree-Uea, B., Wantha, O., & Lee-lacharas, S. (2015). Transcranial direct current stimulation to enhance cognitive function in children with autism: A randomized trial. *European Journal of Paediatric Neurology, 19*(3), 223-229.

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.

- Ashburner, J., Ziviani, J., & Rodger, S. (2010). Surviving in the mainstream: Capacity of children with autism spectrum disorders to perform academically and regulate their emotions and behavior at school. *Research in Autism Spectrum Disorders, 4*(1), 18-27.

- Baio, J., Wiggins, L., Christensen, D. L., Maenner, M. J., Daniels, J., Warren, Z., ... & Dowling, N. F. (2018). Prevalence of autism spectrum disorder among children aged 8 years—Autism and Developmental Disabilities Monitoring Network, 11 sites, United States, 2014. *MMWR Surveillance Summaries, 67*(6), 1.

- Baranek, G. T., David, F. J., Poe, M. D., Stone, W. L., & Watson, L. R. (2006). Sensory experiences questionnaire: Discriminating sensory features in young children with autism, developmental delays, and typical development. *Journal of Child Psychology and Psychiatry, 47*(6), 591-601.

- Beauchamp, T. L., & Childress, J. F. (2001). *Principles of biomedical ethics*. Oxford University Press.

- Bellini, S., Peters, J. K., Benner, L., & Hopf, A. (2007). A meta-analysis of school-based social skills interventions for children with autism spectrum disorders. *Remedial and Special Education, 28*(3), 153-162.

- Ben-Sasson, A., Hen, L., Fluss, R., Cermak, S. A., Engel-Yeger, B., & Gal, E. (2009). A meta-analysis of sensory modulation symptoms in individuals with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 39*(1), 1-11.

- Bikson, M., Brunoni, A. R., Charvet, L. E., Clark, V. P., Cohen, L. G., Deng, Z. D., ... & Woods, A. J. (2016). Rigor and reproducibility in research with transcranial electrical stimulation: An NIMH-sponsored workshop. *Brain Stimulation, 9*(3), 545-550.

- Bodfish, J. W., Symons, F. J., Parker, D. E., & Lewis, M. H. (1999). Varieties of repetitive behavior in autism: Comparisons to mental retardation. *Journal of Autism and Developmental Disorders, 29*(5), 365-373.

- Boggio, P. S., Ferrucci, R., Rigonatti, S. P., Covre, P., Nitsche, M., Pascual-Leone, A., & Fregni, F. (2006). Effects of transcranial direct current stimulation on working memory in patients with Parkinson's disease. *Journal of Neurology Neurosurgery & Psychiatry, 77*(1), 43-50.

- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology, 3*(2), 77-101. https://doi.org/10.1191/1478088706qp063oa

- Brunoni, A. R., Nitsche, M. A., Bolognini, N., Bikson, M., Wagner, T., & Merabet, L. (2012). Clinical research with transcranial direct current stimulation (tDCS): Challenges and future directions. *Brain Stimulation, 5*(3), 175-195.

- Bryman, A. (2016). *Social research methods*. Oxford University Press.

- Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences, 4*(6), 215-222.

- Capps, L., Sigman, M., & Yirmiya, N. (1998). Self-competence and emotional understanding in high-functioning children with autism. *Development and Psychopathology, 10*(2), 193-210.

- Carter, A. S., Messinger, D. S., Stone, W. L., Celimli, S., Nahmias, A. S., & Yoder, P. (2011). A randomized controlled trial of Hanen's ‘More Than Words’ in toddlers with early autism symptoms. *Journal of Child Psychology and Psychiatry, 52*(7), 741-752.

- Carter, E. W., & Hughes, C. (2005). Increasing social interaction among adolescents with intellectual disabilities and their general education peers: Effective interventions. *Research and Practice for Persons with Severe Disabilities, 30*(4), 179-193.

- Clark, V. P., Coffman, B. A., Trumbo, M. C., & Gasparovic, C. (2011). Transcranial direct current stimulation (tDCS) produces localized and specific alterations in neurochemistry: A 1H magnetic resonance spectroscopy study. *Neuroscience Letters, 500*(1), 67-71.

- Coben, R., & Myers, T. E. (2010). The relative efficacy of connectivity guided and symptom-based EEG biofeedback for autistic disorders. *Applied Psychophysiology and Biofeedback, 35*(1), 13-23.

- Cohen Kadosh, R. (2014). Using transcranial direct-current stimulation (tDCS) to enhance cognitive function in the typically developing brain. *Current Directions in Psychological Science, 23*(2), 101-107.

- Cohen, J. D., Perlstein, W. M., Braver, T. S., Nystrom, L. E., Noll, D. C., Jonides, J., & Smith, E. E. (1997). Temporal dynamics of brain activation during a working memory task. *Nature, 386*(6625), 604-608. https://doi.org/10.1038/386604a0

- Constantino, J. N., & Gruber, C. P. (2005). *Social Responsiveness Scale (SRS)*. Western Psychological Services.

- Cooper, J. O., Heron, T. E., & Heward, W. L. (2007). *Applied behavior analysis* (2nd ed.). Pearson.

- Courchesne, E., Campbell, K., & Solso, S. (2011). Brain growth across the life span in autism: Age-specific changes in anatomical pathology. *Brain Research, 1380*, 138-145.

- Coury, D. L., Anagnostou, E., Manning-Courtney, P., Reynolds, A., Cole, L., McCoy, R., & Whitaker, A. (2012). Use of psychotropic medication in children and adolescents with autism spectrum disorders. *Pediatrics, 130*(Supplement 2), S69-S76.

- Creswell, J. W., & Plano Clark, V. L. (2011). *Designing and conducting mixed methods research*. Sage Publications.

- Creswell, J. W., & Plano Clark, V. L. (2017). *Designing and conducting mixed methods research*. Sage Publications.

- Decety, J., & Jackson, P. L. (2004). The functional architecture of human empathy. *Behavioral and Cognitive Neuroscience Reviews, 3*(2), 71-100. https://doi.org/10.1177/1534582304267187

- Diehl, J. J., Schmitt, L. M., Villano, M., & Crowell, C. R. (2012). The clinical use of robots for individuals with autism spectrum disorders: A critical review. *Research in Autism Spectrum Disorders, 6*(1), 249-262.

- Dichter, G. S., Felder, J. N., Green, S. R., Rittenberg, A. M., Sasson, N. J., & Bodfish, J. W. (2012). Reward circuitry function in autism spectrum disorders. *Social Cognitive and Affective Neuroscience, 7*(2), 160-172.

- Duda, M., Ma, R., Haber, N., & Wall, D. P. (2014). Use of machine learning for behavioral distinction of autism and ADHD. *Translational Psychiatry, 4*(12), e424.

- Elsner, B., Kugler, J., Pohl, M., & Mehrholz, J. (2017). Transcranial direct current stimulation (tDCS) for improving aphasia in patients after stroke. *Cochrane Database of Systematic Reviews*.

- Estes, A., Munson, J., Dawson, G., Koehler, E., Zhou, X.-H., & Abbott, R. (2009). Parenting stress and psychological functioning among mothers of preschool children with autism and developmental delay. *Autism, 13*(4), 375-387.

- Eapen, V. (2016). Neurodevelopmental basis of social impairment and autism spectrum disorders. *Pediatric Clinics, 63*(5), 851-859.

- Feinberg, E., Silverstein, M., Donahue, S., & Bliss, R. (2002). The impact of parent expectations on treatment acceptability for children with ADHD. *Journal of Attention Disorders, 5*(2), 119-128.

- Fischbach, G. D., & Lord, C. (2010). The Simons Simplex Collection: A resource for identification of autism genetic risk factors. *Neuron, 68*(2), 192-195.

- Flick, U. (2018). *An introduction to qualitative research*. Sage Publications.

- Fox, M. D., Halko, M. A., Eldaief, M. C., & Pascual-Leone, A. (2014). Measuring and manipulating brain connectivity with resting state functional connectivity magnetic resonance imaging (fcMRI) and transcranial magnetic stimulation (TMS). *NeuroImage, 62*(4), 2232-2243.

- Fregni, F., & Pascual-Leone, A. (2007). Technology insight: noninvasive brain stimulation in neurology--perspectives on the therapeutic potential of rTMS and tDCS. *Nature Clinical Practice Neurology, 3*(7), 383-393.

- Fregni, F., Pascual-Leone, A., Freedman, S. D., Garrett, C., & Mozaffarian, D. (2021). Transcranial direct current stimulation in autism spectrum disorder: Challenges and future directions. *Neuroscience & Biobehavioral Reviews, 125*, 158-170.

- Friend, M., & Bursuck, W. D. (2018). *Including students with special needs: A practical guide for classroom teachers*. Pearson.

- Frith, C. D., & Frith, U. (2007). Social cognition in humans. *Current Biology, 17*(16), R724-R732.

- Fuster, J. M. (2015). *The prefrontal cortex* (5th ed.). Academic Press.

- Gargiulo, R. M., & Metcalf, D. (2013). *Teaching in today's inclusive classrooms: A universal design for learning approach*. Cengage Learning.

- Geretsegger, M., Elefant, C., Mossler, K. A., & Gold, C. (2014). Music therapy for people with autism spectrum disorder. *Cochrane Database of Systematic Reviews*.

- Geschwind, D. H., & State, M. W. (2015). Gene hunting in autism spectrum disorder: On the path to precision medicine. *The Lancet Neurology, 14*(11), 1109-1120.

- Gillis, J. M., & Butler, R. C. (2007). Social skills interventions for preschoolers with autism spectrum disorder: A description of single-subject design studies. *Journal of Early and Intensive Behavior Intervention, 4*(3), 532.

- Gray, C. (2010). *The new social story book*. Future Horizons.

- Greene, J. C. (2007). *Mixed methods in social inquiry*. John Wiley & Sons.

- Haber, S. N. (2016). Corticostriatal circuitry. *Dialogues in Clinical Neuroscience, 18*(1), 7-21. https://doi.org/10.31887/DCNS.2016.18.1/shaber

- Hansen, S. N., Schendel, D. E., & Parner, E. T. (2015). Explaining the increase in the prevalence of autism spectrum disorders: The proportion attributable to changes in reporting practices. *JAMA Pediatrics, 169*(1), 56-62.

- Harris, J. C., Mahone, E. M., & Singer, H. S. (2020). *Neurodevelopmental neuropsychiatry: A comprehensive clinical handbook*. Oxford University Press.

- Hastings, R. P., & Johnson, E. (2002). Stress in UK families conducting home-based behavioral intervention for their young child with autism. *Journal of Autism and Developmental Disorders, 31*(3), 327-336.

- Hodgdon, L. Q. (1995). *Visual strategies for improving communication: Practical supports for school and home*. QuirkRoberts Publishing.

- Hollander, E., Phillips, A. T., & Yeh, C. C. (2005). Targeted treatments for symptom domains in child and adolescent autism. *The Lancet, 362*(9385), 732-734.

- Hollander, E., Phillips, A. T., & Yeh, C. C. (2012). Targeted treatments for symptom domains in child and adolescent autism. *The Lancet, 379*(9831), 1447-1457.

- Howlin, P., Goode, S., Hutton, J., & Rutter, M. (2004). Adult outcome for children with autism. *Journal of Child Psychology and Psychiatry, 45*(2), 212-229.

- Howlin, P., Magiati, I., & Charman, T. (2009). Systematic review of early intensive behavioral interventions for children with autism. *American Journal on Intellectual and Developmental Disabilities, 114*(1), 23-41.

- Humphrey, N., & Symes, W. (2010). Perceptions of social support and experience of bullying among pupils with autistic spectrum disorders in mainstream secondary schools. *European Journal of Special Needs Education, 25*(1), 77-91.

- Individuals with Disabilities Education Act, 20 U.S.C. § 1400 (2004).

- Järbrink, K., Fombonne, E., & Knapp, M. (2003). Measuring the parental service and cost impacts of children with autism spectrum disorder: A pilot study. *Journal of Autism and Developmental Disorders, 33*(4), 395-402.

- Johnson, C. P., & Myers, S. M. (2007). Identification and evaluation of children with autism spectrum disorders. *Pediatrics, 120*(5), 1183-1215.

- Johnson, K., Westermann, G., Sirota, M., & Baker, D. A. (2023). Structured survey on family experiences with tDCS treatment. *Journal of Autism and Developmental Disorders*.

- Just, M. A., Cherkassky, V. L., Keller, T. A., Kana, R. K., & Minshew, N. J. (2007). Functional and anatomical cortical underconnectivity in autism: Evidence from an fMRI study of an executive function task and corpus callosum morphometry. *Cerebral Cortex, 17*(4), 951-961.

- Kaminsky, L., & Dewey, D. (2001). Siblings of children with autism: Social support and family quality of life. *Autism, 5*(2), 193-204.

- Kasari, C., Freeman, S. F., & Paparella, T. (2011). Joint attention and symbolic play in young children with autism: A randomized controlled intervention study. *Journal of Child Psychology and Psychiatry, 47*(6), 611-620.

- Kasari, C., Rotheram-Fuller, E., Locke, J., & Gulsrud, A. (2012). Making the connection: Randomized controlled trial of social skills at school for children with autism spectrum disorders. *Journal of Child Psychology and Psychiatry, 53*(4), 431-439.

- Keshavan, M. S., Vinogradov, S., Rumsey, J., Sherrill, J., & Wagner, A. (2013). Cognitive training in mental disorders: Update and future directions. *American Journal of Psychiatry, 171*(5), 510-522.

- Kim, Y. S., Leventhal, B. L., Koh, Y. J., Fombonne, E., Laska, E., Lim, E. C., ... & Grinker, R. R. (2011). Prevalence of autism spectrum disorders in a total population sample. *American Journal of Psychiatry, 168*(9), 904-912.

- Klin, A., McPartland, J., & Volkmar, F. R. (2007). Asperger syndrome. In F. R. Volkmar, R. Paul, A. Klin, & D. Cohen (Eds.), *Handbook of Autism and Pervasive Developmental Disorders* (3rd ed., pp. 88-125). Wiley.

- Koegel, L. K., Bimbela, A., & Schreibman, L. (2014). Collateral effects of parent training on family interactions. *Journal of Autism and Developmental Disorders, 24*(6), 347-359.

- Koegel, R. L., & Koegel, L. K. (2006). *Pivotal response treatments for autism: Communication, social, & academic development*. Paul H Brookes Publishing.

- Kurth, J. A., & Mastergeorge, A. M. (2010). Academic and cognitive profiles of students with autism: The impact of placement in general education versus special education classrooms. *Journal of Autism and Developmental Disorders, 40*(5), 617-625.

- Kurth, J. A., & Mastergeorge, A. M. (2012). Academic and cognitive profiles of students with autism: Implications for classroom practice and placement. *International Journal of Special Education, 27*(2), 8-14.

- Lai, M. C., Lombardo, M. V., & Baron-Cohen, S. (2014). Autism. *The Lancet, 383*(9920), 896-910.

- Leitner, Y. (2014). The co-occurrence of autism and attention deficit hyperactivity disorder in children–what do we know? *Frontiers in Human Neuroscience, 8*, 268.

- Leaf, J. B., Oppenheim-Leaf, M. L., Leaf, R., McEachin, J., Taubman, M., & Parker, T. (2016). Concerns about ABA-based intervention: An evaluation and recommendations. *Focus on Autism and Other Developmental Disabilities, 31*(1), 70-78.

- Loomes, R., Hull, L., & Mandy, W. P. (2017). What is the male-to-female ratio in autism spectrum disorder? A systematic review and meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry, 56*(6), 466-474.

- Lord, C., Elsabbagh, M., Baird, G., & Veenstra-Vanderweele, J. (2018). Autism spectrum disorder. *The Lancet, 392*(10146), 508-520.

- Lord, C., Rutter, M., DiLavore, P. C., & Risi, S. (2000). *Autism Diagnostic Observation Schedule (ADOS)*. Western Psychological Services.

- Lovaas, O. I. (1987). Behavioral treatment and normal educational and intellectual functioning in young autistic children. *Journal of Consulting and Clinical Psychology, 55*(1), 3-9.

- Maenner, M. J., Shaw, K. A., Baio, J., Washington, A., Patrick, M., DiRienzo, M., ... & Dietz, P. M. (2020). Prevalence of autism spectrum disorder among children aged 8 years—Autism and Developmental Disabilities Monitoring Network, 11 sites, United States, 2016. *MMWR Surveillance Summaries, 69*(4), 1-12.

- Mandell, D. S., Morales, K. H., Xie, M., Lawer, L. J., Stahmer, A. C., & Marcus, S. C. (2010). Age of diagnosis among medicaid-enrolled children with autism, 2001–2004. *Psychiatric Services, 61*(8), 822-829.

- Marcus, R. N., Owen, R., Kamen, L., Manos, G., McQuade, R. D., Carson, W. H., & Findling, R. L. (2009). A placebo-controlled, fixed-dose study of aripiprazole in children and adolescents with irritability associated with autistic disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 48*(11), 1110-1119.

- Matson, J. L., & Smith, K. R. (2012). Current status of intensive behavioral interventions for young children with autism and PDD-NOS. *Research in Autism Spectrum Disorders, 2*(1), 60-74.

- McConachie, H., & Diggle, T. (2007). Parent implemented early intervention for young children with autism spectrum disorder: A systematic review. *Journal of Evaluation in Clinical Practice, 13*(1), 120-129.

- McCracken, J. T., McGough, J., Shah, B., Cronin, P., Hong, D., Aman, M. G., ... & Research Units on Pediatric Psychopharmacology Autism Network. (2002). Risperidone in children with autism and serious behavioral problems. *New England Journal of Medicine, 347*(5), 314-321.

- McElhanon, B. O., McCracken, C., Karpen, S., & Sharp, W. G. (2014). Gastrointestinal symptoms in autism spectrum disorder: A meta-analysis. *Pediatrics, 133*(5), 872-883.

- McLeskey, J., Waldron, N. L., & Redd, L. (2012). *Inclusion programs for students with autism spectrum disorders: Research-based approaches*. Guilford Press.

- McPheeters, M. L., Warren, Z., Sathe, N., Bruzek, J. L., Krishnaswami, S., Jerome, R. N., & Veenstra-VanderWeele, J. (2011). A systematic review of medical treatments for children with autism spectrum disorders. *Pediatrics, 127*(5), e1312-e1321.

- Miniussi, C., Ruzzoli, M., & Walsh, V. (2013). The mechanism of transcranial direct current stimulation: From cortical plasticity to network interaction. *Neuropsychologia, 51*(8), 1260-1266.

- Mitchell, P., Parsons, S., & Leonard, A. (2007). Using virtual environments for teaching social understanding to adolescents with autistic spectrum disorders. *Journal of Autism and Developmental Disorders, 37*(3), 589-600.

- Morningstar, M. E., Kurth, J. A., & Johnson, P. (2017). Examining the meaning of participation for students with significant disabilities in general education classrooms. *Exceptional Children, 84*(1), 45-60.

- National Autism Center. (2015). *Findings and conclusions: National standards project, phase 2*. Randolph, MA: Author.

- National Research Council. (2001). *Educating children with autism*. National Academies Press.

- Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *Journal of Physiology, 527*(3), 633-639.

- Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology, 57*(10), 1899-1901.

- Nitsche, M. A., & Paulus, W. (2008). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *Journal of Physiology, 527*(3), 633-639.

- Nitsche, M. A., & Paulus, W. (2011). Transcranial direct current stimulation—update 2011. *Restorative Neurology and Neuroscience, 29*(6), 463-492.

- Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences, 9*(5), 242-249.

- Odom, S. L., Boyd, B. A., Hall, L. J., & Hume, K. A. (2014). Comprehensive treatment models for children and youth with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 44*(1), 425-437.

- Osborne, L. A., McHugh, L., Saunders, J., & Reed, P. (2008). Parenting stress reduces the effectiveness of early teaching interventions for autistic spectrum disorders. *Journal of Autism and Developmental Disorders, 38*(6), 1092-1103.

- Oswald, D. P., & Sonenklar, N. A. (2007). Medication use among children with autism spectrum disorders. *Journal of Child and Adolescent Psychopharmacology, 17*(3), 348-355.

- Paul, R., Norbury, C. F., & Gosse, C. (2014). Language deficits in children with autism spectrum disorders. *Autism Spectrum Disorder: A Review of the Current Understanding and Treatments*.

- Paul, R., Orlovski, S. M., Marcinko, H. C., & Volkmar, F. (2014). Conversational behaviors in youth with high-functioning ASD and Asperger syndrome. *Journal of Autism and Developmental Disorders, 39*(1), 115-125.

- Pascual-Leone, A., Bartres-Faz, D., & Keenan, J. P. (2011). Transcranial magnetic stimulation: Studying the brain–behaviour relationship by induction of ‘virtual lesions’. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences, 354*(1387), 1229-1238.

- Pisula, E., & Porebowicz-Dorsmann, A. (2017). Family functioning, parenting stress, and quality of life in mothers and fathers of Polish children with high functioning autism or Asperger syndrome. *Journal of Autism and Developmental Disorders, 47*(2), 353-365.

- Polanía, R., Nitsche, M. A., & Paulus, W. (2012). Modulating functional connectivity patterns and topological functional organization of the human brain with transcranial direct current stimulation. *Human Brain Mapping, 32*(8), 1234-1249.

- Posey, D. J., Stigler, K. A., Erickson, C. A., & McDougle, C. J. (2008). Antipsychotics in the treatment of autism. *Journal of Clinical Investigation, 118*(1), 6-14.

- Reaven, J., Blakeley-Smith, A., Culhane-Shelburne, K., & Hepburn, S. (2012). Group cognitive behavior therapy for children with high-functioning autism spectrum disorders and anxiety: A randomized trial. *Journal of Child Psychology and Psychiatry, 53*(4), 410-419.

- Reichow, B. (2012). Overview of meta-analyses on early intensive behavioral intervention for young children with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 42*(4), 512-520.

- Reichow, B., Barton, E. E., Boyd, B. A., & Hume, K. (2018). Early intensive behavioral intervention (EIBI) for young children with autism spectrum disorders (ASD). *Cochrane Database of Systematic Reviews*.

- Research Units on Pediatric Psychopharmacology Autism Network. (2005). Randomized, controlled, crossover trial of methylphenidate in pervasive developmental disorders with hyperactivity. *Archives of General Psychiatry, 62*(11), 1266-1274.

- Research Units on Pediatric Psychopharmacology Autism Network. (2005). Risperidone treatment of autistic disorder: Longer-term benefits and blinded discontinuation after 6 months. *American Journal of Psychiatry, 162*(7), 1361-1369.

- Riley, K., Davis, N. J., Bentley, P., & Cohen Kadosh, R. (2018). Comparing the effects of tDCS and TMS on social communication in children with ASD. *Brain Stimulation, 11*(3), 655-661.

- Rolls, E. T. (2004). The functions of the orbitofrontal cortex. *Brain and Cognition, 55*(1), 11-29.

- Sandbank, M., Bottema-Beutel, K., Crowley, S., Cassidy, M., Dunham, K., Feldman, J. I., ... & Woynaroski, T. G. (2020). Project AIM: Autism intervention meta-analysis for studies of young children. *Psychological Bulletin, 146*(1), 1-29.

- Sansosti, F. J., & Powell-Smith, K. A. (2010). High-functioning autism/Asperger syndrome in schools: Assessment and intervention. *Psychology in the Schools, 47*(9), 891-896.

- Saxe, R., & Kanwisher, N. (2003). People thinking about thinking people: The role of the temporo-parietal junction in "theory of mind". *NeuroImage, 19*(4), 1835-1842.

- Scahill, L., Bearss, K., McCracken, C. E., et al. (2015). Parent training for young children with autism and disruptive behavior: Response to treatment and maintenance of gains. *Journal of the American Academy of Child and Adolescent Psychiatry, 54*(8), 610-619.

- Schneider, H. D., & Hopp, J. P. (2011). The use of tDCS to enhance cognitive and behavioral therapy outcomes for individuals with neurodevelopmental disorders. *Journal of Visualized Experiments, 1*(54), e3024. https://doi.org/10.3791/3024

- Schneider, H. D., & Hopp, J. P. (2019). The effects of transcranial direct current stimulation on cognitive functions in children with autism spectrum disorder: A pilot study. *Neuropsychologia, 123*, 14-23.

- Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T., & Baird, G. (2008). Psychiatric disorders in children with autism spectrum disorders: Prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child & Adolescent Psychiatry, 47*(8), 921-929.

- Smith, T. (2001). Discrete trial training in the treatment of autism. *Focus on Autism and Other Developmental Disabilities, 16*(2), 86-92.

- Smith, T. (2007). Outcome of early intervention for children with autism. *Clinical Psychology: Science and Practice, 4*(1), 33-49.

- Smith, T., Eikeseth, S., Groen, A. D., & Wynn, J. W. (2000). Randomized trial of intensive early intervention for children with pervasive developmental disorder. *American Journal on Mental Retardation, 105*(4), 269-285.

- Smith, T., Killian, J., & Evans, E. (2019). Telehealth delivery of services for families of children with ASD in rural areas. *Journal of Autism and Developmental Disorders, 49*(2), 1-12.

- Solomon, M., Van Egeren, L. A., Mahoney, G., Quon Huber, M. S., & Zimmerman, P. (2018). Empowerment and competency in parents of children with autism spectrum disorder. *Research in Autism Spectrum Disorders, 2*(2), 223-237.

- Sundberg, M. L., & Partington, J. W. (1998). *Teaching language to children with autism or other developmental disabilities*. Behavior Analysts, Inc.

- Tager-Flusberg, H. (2000). Language and understanding minds: Connections in autism. In S. Baron-Cohen, H. Tager-Flusberg, & D. J. Cohen (Eds.), *Understanding other minds: Perspectives from developmental cognitive neuroscience* (pp. 124-149). Oxford University Press.

- Tashakkori, A., & Teddlie, C. (2010). *SAGE handbook of mixed methods in social & behavioral research*. Sage.

- Taylor, J. L., & Seltzer, M. M. (2011). Changes in the autism behavioral phenotype during the transition to adulthood. *Journal of Autism and Developmental Disorders, 41*(10), 1375-1390.

- Turkeltaub, P. E., et al. (2012). Enhanced phonological processing after tDCS stimulation of the left temporoparietal cortex in dyslexia. *Brain and Language, 121*(1), 1-11.

- Uddin, L. Q., Supekar, K., & Menon, V. (2013). Reconceptualizing functional brain connectivity in autism from a developmental perspective. *Frontiers in Human Neuroscience, 7*, 458.

- Vorstman, J. A. S., Parr, J. R., Moreno-De-Luca, D., Anney, R. J. L., Nurnberger, J. I., & Hallmayer, J. F. (2017). Autism genetics: Opportunities and challenges for clinical translation. *Nature Reviews Genetics, 18*(6), 362-376.

- Vygotsky, L. S. (1978). *Mind in society: The development of higher psychological processes*. Harvard University Press.

- Wehman, P., Schall, C., McDonough, J., Molinelli, A., Riehle, J. E., Ham, W., & Thiss, W. (2014). Competitive employment for youth with autism spectrum disorders: Early results from a randomized clinical trial. *Journal of Autism and Developmental Disorders, 44*(3), 487-500.

- Whitaker, P. (2002). Supporting families of preschool children with autism: What parents want and what helps. *Autism: The International Journal of Research and Practice, 6*(4), 411-426.

- Williams, K., Brignell, A., Randall, M., Silove, N., & Hazell, P. (2010). Selective serotonin reuptake inhibitors (SSRIs) for autism spectrum disorders (ASD). *Cochrane Database of Systematic Reviews*.

- Woods, A. J., Antal, A., Bikson, M., Boggio, P. S., Brunoni, A. R., Celnik, P., ... & Nitsche, M. A. (2016). A technical guide to tDCS, and related non-invasive brain stimulation tools

. *Clinical Neurophysiology, 127*(2), 1031-1048.

- Woodgate, R. L., Ateah, C., & Secco, L. (2008). Living in a world of our own: The experience of parents who have a child with autism. *Qualitative Health Research, 18*(8), 1075-1083.

- Zhu, C., et al. (2014). tDCS over the primary motor cortex enhances motor coordination in developmental coordination disorder. *Developmental Medicine & Child Neurology, 56*(4), 328-334.

- Zwaigenbaum, L., Bauman, M. L., Choueiri, R., Kasari, C., Carter, A., Granpeesheh, D., ... & Wagner, S. (2015). Early intervention for children with autism spectrum disorder under 3 years of age: Recommendations for practice and research. *Pediatrics, 136*(Supplement 1), S60-S81.

Appendix

Appendix A

Ethics form

AIU
Atlantic International University

June 17, 2024
Honolulu, Hawaii

Re: Ethics Approval and Request Permission to Conduct Research and Gather Data

To Whom it may concern,

Alptekin Aydin is a student of the Doctorate Program in *Clinical Psychology* at our Atlantic International University.
The thesis proposal of Alptekin Aydin has been approved. His research methodology has been approved by the Ethics Review Board at AIU. He is in the process of gathering data for his thesis.
Completing an approved thesis is a requirement to graduate with a Doctoral degree at AIU.
Upon completion of the written thesis, there will also be an oral thesis defense.
The title of his thesis will be "*Assessing the Impact of tDCS Treatment Over the Dorsolateral Prefrontal Cortex (DLPFC) on Social Communication in Children with ASD: A Survey of Family Experiences*".

We humbly request that your organization support his academic and research efforts.
We ask respectfully that you inform us of any problems or concerns connected with the thesis research of Alptekin Aydin. Please contact us with any concern that you

Sincerely,

Edward J Lambert EdL

AIU

Dr. Edward Lambert
Academic Director
Atlantic International University
Honolulu, Hawaii, USA
edward@aiu.edu
studentservices@aiu.edu

Atlantic International University, Pioneer Plaza 900 Fort Street Mall -905, Honolulu, Hawaii 96813
Phone: 808-521-1868 Fax: 808-421-1678

Appendix B

Research Protocol

Assessing the Impact of tDCS Treatment Over the Dorsolateral Prefrontal Cortex (DLPFC) on Social Communication in Children with ASD: A Survey of Family Experiences

1. Introduction

The research aims to evaluate the impact of transcranial Direct Current Stimulation (tDCS) on social communication in children with Autism Spectrum Disorder (ASD). Additionally, it will survey family experiences with this treatment.

2. Background and Rationale

Autism Spectrum Disorder (ASD): ASD is a neurodevelopmental disorder with challenges in social interaction, communication, and repetitive behaviors. Traditional treatments involve behavioral interventions and educational support, but recent advances suggest neurostimulation techniques like tDCS may offer additional benefits.

Transcranial Direct Current Stimulation (tDCS): tDCS is a non-invasive brain stimulation technique that uses low electrical currents to modulate neuronal activity. The DLPFC is targeted due to its role in social cognition and executive functions.

Previous Research: The PlatoWork tDCS headset has shown potential in cognitive enhancement and therapeutic applications, indicating possible improvements in cognitive and social outcomes for children with ASD.

3. Research Questions and Objectives

Research Questions:

1. Does tDCS targeting the DLPFC improve social communication in children with ASD?

2. What are the family experiences and perceptions of the tDCS treatment?

Objectives:

• Primary Objective: Assess the impact of tDCS on social communication in children with ASD.

• Secondary Objective: Explore family experiences and satisfaction with the tDCStreatment.

4. Methodology

Participants: Families of the 300 children from the previous QEEG-guided tDCS study.

Survey Design: A structured survey will gather qualitative data on family experiences, perceived benefits, challenges, and overall satisfaction with the tDCS treatment.

Data Collection:

• Survey Administration: Distributed via online platforms and in-person sessions at rehabilitation centers.

• Survey Content: Includes questions on treatment effectiveness, observed improvements in social communication, ease of use, and overall satisfaction.

5. Data Collection and Analysis

Quantitative Analysis:

• Descriptive Statistics: Summarize survey responses.

• Software Tools: SPSS and R for statistical analysis.

Qualitative Analysis:

• Thematic Analysis: Identify common themes and insights from survey responses.

• Software Tools: NVivo for organizing and analyzing qualitative data.

6. Expected Outcomes

The study aims to provide a comprehensive understanding of the impact of tDCS on social communication in children with ASD and gather insights into family experiences. Expected

outcomes include improved social communication in children and detailed insights into family perceptions and satisfaction with the tDCS treatment.

7. Ethical Considerations

Ethical approval will be obtained from the Ethics Committee. Informed consent will be secured from all participants' parents or guardians. Confidentiality and anonymity of survey responses will be maintained.

8. Timeline

Timeline:

- Month 1-2: Ethical approval and participant recruitment.
- Month 3-4: Survey design and pilot testing.
- Month 5-6: Data collection.
- Month 7-8: Data analysis.
- Month 9: Reporting and dissemination of findings.

Appendix C

Information Sheet

Title of the Study: Assessing the Impact of tDCS Treatment Over the Dorsolateral Prefrontal Cortex (DLPFC) on Social Communication in Children with ASD: A Survey of Family Experiences

Researcher: Alptekin Aydin

Institution: Atlantic International University, Honolulu, Hawaii

Contact Information:

Introduction

You are being invited to participate in a research study conducted by Alptekin AYDIN from Atlantic International University. This study aims to evaluate the impact of transcranial Direct Current Stimulation (tDCS) targeting the Dorsolateral Prefrontal Cortex (DLPFC) on social communication in children with Autism Spectrum Disorder (ASD), and to explore the experiences of their families.

Purpose of the Study

The purpose of this study is twofold:

1. To assess the effectiveness of tDCS in improving social communication in children with ASD.
2. To understand family experiences and satisfaction with the tDCS treatment.

Your participation is crucial as it will contribute to a better understanding of how tDCS can be used to support children with ASD. The data collected will help improve future treatments and provide valuable insights into family experiences, which can inform healthcare professionals and policymakers to better support ASD families.

What is tDCS?

Transcranial Direct Current Stimulation (tDCS) is a non-invasive brain stimulation technique that applies a low electrical current to the scalp to modulate neuronal activity. The DLPFC is targeted in this study due to its role in social cognition and executive functions.

Why Have I Been Invited?

You have been invited to participate because you are a parent or guardian of a child who has undergone tDCS treatment. Your insights and experiences are valuable in understanding the broader impact of this treatment.

What Will Happen During the Study?

If you decide to participate, you will be asked to complete a detailed survey that includes questions about:

- Demographic information
- Details of the tDCS treatment your child received
- Observed effects on your child's social communication, behaviour, cognitive abilities, and emotional regulation
- The impact of the treatment on family dynamics, parental stress, and well-being
- Overall satisfaction with the treatment and any feedback or suggestions

The survey will take approximately 30-45 minutes to complete and can be done online or in person at your convenience.

Are There Any Risks?

There are no known risks associated with participating in this study. Some questions may touch on sensitive topics, but you are free to skip any questions that make you uncomfortable.

Are There Any Benefits?

While there are no direct benefits to you, your participation will contribute to a better understanding of tDCS treatment's effectiveness and the experiences of families. This information could help improve future treatments and support for children with ASD.

Will My Information Be Kept Confidential?

Yes, all responses will be kept confidential. Data will be anonymized and stored securely. Only the research team will have access to the data. No identifying information will be used in any publications or presentations.

Do I Have to Take Part?

No, your participation is entirely voluntary. You can withdraw from the study at any time without any consequences or loss of benefits.

What If I Have Questions?

If you have any questions or concerns about the study, please contact Alptekin AYDIN at [Researcher’s Email] or [Researcher’s Phone Number]. For questions about your rights as a participant, you may contact the Ethics Committee at Atlantic International University.

Consent

Your participation is voluntary, and you will be asked to sign a consent form before taking part in the study. This form will confirm that you understand the study's purpose, procedures, risks, benefits, and your rights as a participant.

Thank you for considering participation in this study. Your insights and experiences are invaluable to us.

Appendix D

Informed Consent Form

Title of the Study: Assessing the Impact of tDCS Treatment Over the Dorsolateral Prefrontal Cortex (DLPFC) on Social Communication in Children with ASD: A Survey of Family Experiences

Researcher: Alptekin AYDIN

Institution: Atlantic International University, Honolulu, Hawaii

Contact Information:

Introduction

You are invited to participate in a research study conducted by Alptekin AYDIN at Atlantic International University. The purpose of this study is to evaluate the impact of transcranial Direct Current Stimulation (tDCS) targeting the Dorsolateral Prefrontal Cortex (DLPFC) on social communication in children with Autism Spectrum Disorder (ASD), and to explore family experiences with this treatment.

Participation

Your participation in this study is entirely voluntary. You may decline to participate or withdraw from the study at any time without penalty. If you choose not to participate or withdraw, you will not be penalised or lose ay advantages to which you are otherwise entitled.

Procedures

If you agree to participate in this study, you will be asked to complete a survey that covers various aspects of your child's treatment and your family's experiences with tDCS. The survey will include questions about:

- Demographic information
- Treatment details
- Observed effects on your child
- Impact on family dynamics
- Satisfaction with the treatment and feedback
- Open-ended questions about your personal experiences

The survey will take approximately 30-45 minutes to complete.

Risks and Discomforts

There are no known risks associated with participating in this study. However, some questions may be of a sensitive nature and could potentially cause discomfort. You are free to skip any questions that you do not wish to answer.

Benefits

While there are no direct benefits to you for participating in this study, your responses will contribute to a better understanding of the effects of tDCS treatment on social communication in children with ASD and the experiences of their families. This information may help improve future treatments and support for children with ASD and their families.

Confidentiality

Your responses will be kept confidential. All data collected will be anonymized and stored securely. Only the research team will have access to the data. Your identity will not be revealed in any publications or presentations resulting from this study. Direct quotes from the survey may be used in publications, but these will be anonymized to ensure confidentiality.

Contact Information

If you have any questions or concerns about this study, please contact Alptekin AYDIN at If you have questions about your rights as a research participant, you may contact the Ethics Committee at Atlantic International University.

Consent

Please read the statements below and indicate your agreement by checking the appropriate boxes:

1. I have read and understood the information provided about this survey.
    - ☐ Yes
    - ☐ No
2. I understand that participation is voluntary and that I can withdraw from the survey at any time without any consequences.
    - ☐ Yes
    - ☐ No
3. I understand that my responses will be kept confidential and that no identifying information will be shared.
    - ☐ Yes
    - ☐ No
4. I understand that the data collected will be anonymized before analysis and publication.
    - ☐ Yes
    - ☐ No
5. I consent to participate in this survey and understand that my responses will be used for research purposes only.
    - ☐ Yes

- ☐ No

6. I agree to the use of my anonymized data in future related research studies.
   - ☐ Yes
   - ☐ No
7. I understand that my participation or non-participation will not affect any services or support I receive.
   - ☐ Yes
   - ☐ No
8. I understand that the results of this study may be published, but my identity will remain confidential.
   - ☐ Yes
   - ☐ No
9. I know whom to contact if I have any questions about the survey or my rights as a participant.
   - ☐ Yes
   - ☐ No

Participant's Signature: _____

Date: _____

Researcher's Signature: _____

Date: _____

Appendix E

Survey

Section A: Demographics

1. Child Information:

   Age: ___

   Gender: ___

   Age at ASD diagnosis: ___

   Current severity level of ASD:

   - Mild
   - Moderate
   - Severe

   Co-existing conditions (if any): ___

   Primary language spoken at home: ___

   Educational setting:

   - Mainstream
   - Special education
   - Home-schooled
   - Other (please specify): ___

2. Parent/Guardian Information:

   Relationship to child:

   - Mother
   - Father

- Guardian
- Other (please specify): ___

Age: ___

Highest level of education:

- Less than high school
- High school diploma or equivalent
- Some college
- Associate degree
- Bachelor’s degree
- Master’s degree
- Doctorate

Occupation: ___

Marital status:

- Single
- Married
- Divorced
- Widowed
- Other (please specify): ___

Number of children in the household: ___

Primary caregiver:

- Yes

- No

Involvement in ASD support groups or communities:

- Yes
- No

Section B: Treatment Details

1. tDCS Treatment Information:

Duration of treatment (weeks/months): ___

Frequency of sessions per week: ___

Duration of each session (minutes): ___

Total number of sessions completed: ___

Any previous treatments tried (Yes/No): ___

- If yes, please specify: ___

Who administers the treatment:

- Parent
- Therapist
- Other (please specify): ___

Were there any variations in the prescribed treatment schedule (Yes/No): ___

- If yes, please explain the reason for the variation and its impact: ___

How did you track the treatment sessions (e.g., journal, app, calendar): ___

Any side effects observed during the treatment (Yes/No): ___

- If yes, please describe: ___

2. PlatoWork Headset Usage:

Ease of use:

- Very easy
- Easy
- Neutral
- Difficult
- Very difficult

Any technical issues experienced (Yes/No): ___

- If yes, please describe: ___

Comfort level for the child during sessions:

- Very comfortable
- Comfortable
- Neutral
- Uncomfortable
- Very uncomfortable

Compliance with treatment schedule:

- Always

- Most of the time
- Sometimes
- Rarely
- Never

Child's willingness to participate in the treatment:

- Very willing
- Willing
- Neutral
- Unwilling
- Very unwilling

Did the child express any discomfort or pain during the sessions (Yes/No): ___

- If yes, please describe: ___

Did you use any additional tools or techniques to help your child remain comfortable during the sessions (e.g., distraction methods, comfort items): ___

How did you ensure the proper placement and use of the headset during each session: ___

Were there any interruptions or difficulties in completing the treatment sessions as scheduled (Yes/No): ___

- If yes, please describe: ___

Section C: Observed Effects

1. Social Communication Improvements:

   Noticed improvements in verbal communication:

   - Strongly agree
   - Agree
   - Neutral
   - Disagree
   - Strongly disagree

   Noticed improvements in non-verbal communication:

   - Strongly agree
   - Agree
   - Neutral
   - Disagree
   - Strongly disagree

   Specific examples of observed improvements in verbal communication: ___

   Specific examples of observed improvements in non-verbal communication: ___

   Impact on child's ability to initiate conversations:

   - Significant improvement
   - Some improvement
   - No change
   - Worsened

Impact on child's ability to maintain conversations:

- Significant improvement
- Some improvement
- No change
- Worsened

Changes in child's use of gestures and facial expressions:

- Significant improvement
- Some improvement
- No change
- Worsened

2. Behavioural Changes:

Reduction in problematic behaviours:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Improvements in social interactions with peers and family members:

- Strongly agree
- Agree
- Neutral

- Disagree
- Strongly disagree

Any new challenging behaviours observed:

- Yes
- No
- If yes, please describe: ___

Specific examples of improved social interactions with peers: ___

Specific examples of improved social interactions with family members: ___

Changes in the child's ability to share and take turns:

- Significant improvement
- Some improvement
- No change
- Worsened

3. Cognitive and Emotional Changes:

Improvements in attention span and focus:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Changes in emotional regulation:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Specific examples of cognitive changes observed: ___

Specific examples of emotional changes observed: ___

Changes in academic performance (if applicable):

- Significant improvement
- Some improvement
- No change
- Worsened

Impact on child's ability to complete tasks and follow instructions:

- Significant improvement
- Some improvement
- No change
- Worsened

Changes in child's mood and overall happiness:

- Significant improvement
- Some improvement

- No change
- Worsened

Any observed side effects or negative outcomes from the treatment:

- Yes
- No
- If yes, please describe: ___

Section D: Family Impact

1. Family Dynamics:

Changes in family routines and dynamics due to the treatment:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Impact on sibling relationships:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Specific examples of changes in family dynamics: ___

Support received from extended family or community:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Impact on family activities and outings:

- Significant improvement
- Some improvement
- No change
- Worsened

Changes in family communication patterns:

- Significant improvement
- Some improvement
- No change
- Worsened

Support received from healthcare professionals:

- Strongly agree
- Agree
- Neutral
- Disagree

- Strongly disagree

Level of involvement of other family members in the treatment process:

- Very high
- High
- Moderate
- Low
- Very low

Impact on overall family harmony and cohesiveness:

- Significant improvement
- Some improvement
- No change
- Worsened

2. Parental Stress and Well-being:

Level of stress related to managing the treatment:

- Very high
- High
- Moderate
- Low
- Very low

Overall well-being of the parent/guardian:

- Improved
- No change
- Deteriorated

Impact on work-life balance:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Specific examples of any changes in parental well-being: ___

Access to respite care or other support services:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Changes in social support network (friends, support groups, etc.):

- Significant improvement
- Some improvement
- No change
- Worsened

Availability of mental health resources for parents:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

Impact on financial stability due to treatment costs:

- Significant impact
- Some impact
- No impact

Parent/guardian's satisfaction with the information and resources provided about the treatment:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied
- Very dissatisfied

Strategies used by parents to manage stress related to the treatment process: __

Section E: Satisfaction and Feedback

1. Overall Satisfaction:

Satisfaction with the overall treatment experience:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied
- Very dissatisfied

Satisfaction with the results of the tDCS treatment:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied
- Very dissatisfied

Would you recommend this treatment to other families:

- Yes
- No

Any additional comments or feedback: ___

Satisfaction with the support received from healthcare professionals:

- Very satisfied
- Satisfied

- Neutral
- Dissatisfied
- Very dissatisfied

Satisfaction with the information provided about the treatment:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied
- Very dissatisfied

Satisfaction with the ease of integrating the treatment into daily routines:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied
- Very dissatisfied

Overall effectiveness of the tDCS treatment on your child's quality of life:

- Very effective
- Effective
- Neutral
- Ineffective

- Very ineffective

Comparison of tDCS treatment to other treatments tried (if applicable):

- Much better
- Better
- About the same
- Worse
- Much worse

2. Suggestions for Improvement:

Suggestions for improving the PlatoWork headset or treatment process: ___

Any additional support or resources that would be helpful: ___

Ideas for improving the communication and support from healthcare professionals: ___

Recommendations for future research or areas of interest: ___

Additional tools or technologies that could assist in the treatment process: ___

Any logistical challenges faced and how they could be mitigated: ___

Recommendations for other families starting this treatment: ___

How satisfied are you with the customization and personalization options of the treatment protocol:

- Very satisfied
- Satisfied
- Neutral
- Dissatisfied

- Very dissatisfied

Section F: Open-Ended Questions

1. Personal Experiences:

   Describe any notable changes you have observed in your child since starting the tDCS treatment:

   - Improvements in specific skills (e.g., communication, social interaction, academic performance):
   - Changes in behaviour (e.g., reduced meltdowns, increased compliance):
   - Emotional changes (e.g., mood, frustration tolerance):
   - Any regressions or negative changes observed:

   Share any challenges you faced while administering the treatment and how you overcame them:

   - Initial setup and learning curve:
   - Child's resistance or discomfort during sessions:
   - Managing the treatment schedule alongside daily routines:
   - Technical issues with the PlatoWork headset:
   - Any support or resources that were particularly helpful:

   Provide any additional comments on how the treatment has impacted your family's daily life:

   - Changes in family routines and dynamics:
   - Impact on sibling relationships:
   - Effects on parental stress and well-being:

- Any changes in social activities or participation in community events:
- Financial implications of the treatment:

Describe your child's attitude towards the tDCS treatment:

- Initial reactions:
- Any changes in attitude over time:
- Specific aspects of the treatment your child liked or disliked:

How do you feel about the long-term prospects of using tDCS treatment for your child?

- Expectations for future progress:
- Concerns about long-term use:
- Hopes for continued improvements:

In what ways has the tDCS treatment influenced your approach to managing your child's ASD?

- Changes in parenting strategies:
- Adoption of new therapies or interventions:
- Integration with other treatments or educational plans:

Share any memorable moments or breakthroughs during the tDCS treatment period:

- Positive interactions or milestones:
- Moments of significant progress:
- Any notable feedback from teachers or therapists:

Any suggestions for other parents considering tDCS treatment for their child:

- Tips for getting started:

- Managing expectations:
- Building a support network:

How do you envision the role of tDCS treatment in your child's future care plan?

- Plans for continued use:
- Integration with other therapies:
- Desired outcomes:

Additional comments or reflections:

- Anything else you would like to share about your experience with the tDCS treatment:
- Thoughts on the importance of research in this area:

Appendix F

Turkish version filled out Survey

DLPFC Üzerindeki tDCS Tedavisinin Özel Çocuklarda Sosyal İletişim Üzerindeki Etkisinin Değerlendirilmesi: Aile deneyimleri anketi — SurveyMonkey

#276

COMPLETE

Collector: Web Link 1 (Web Link)
Started: Thursday, June 27, 2024 8:37:27 PM
Last Modified: Thursday, June 27, 2024 8:45:56 PM
Time Spent: 00:08:29
IP Address: 92.72.38.180

Page 1: Bölüm A: Demografik Bilgiler

Q1

Çocuğun yaşı

6

Q2 — **Erkek**

Cinsiyet:

Q3

Otizm teşhis yaşı

4

Q4 — **Ağır**

Mevcut Otizm şiddet seviyesi

Q5 — **Respondent skipped this question**

Eşlik eden durumlar (varsa):

Q6

Evde konuşulan ana dil:

Türkçe

Q7 — **Özel eğitim**

Eğitim durumu:

Q8 — **Anne**

Ebeveyn/Vasi Bilgileri: Çocuk ile ilişkisi

1 / 9

Q9

Veli Yaşı:

35

Q10 **Lisans (4 yıllık)**

En yüksek eğitim düzeyi:

Q11

Meslek:

Sosyoloji

Q12 **Evli**

Medeni durum:

Q13 **3**

Evdeki çocuk sayısı:

Q14 **Aile**

Çocuğun bakımında birinci derece sorumlu

Q15 **Evet**

Otizm destek gruplarına veya topluluklarına katılım:

Page 2: Bölüm B: Tedavi Detayları

Q16 **3 ay**

Tedavi süresi

Q17

Haftalık seans sayısı

5

Q18

Her bir seans süresi (dakika):

30

Q19 Respondent skipped this question

Tamamlanan toplam seans sayısı:

Q20

Daha önce denenmiş tedaviler:

Özel eğitiö

Q21 **Terapist**

Tedaviyi uygulayan kişi:

Q22 **Hayır**

Öngörülen tedavi programında değişiklik oldu mu?

Q23

Tedavi seanslarını nasıl takip ettiniz?

Plato progamı ile

Q24

Tedavi sırasında gözlenen yan etkiler oldu mu?

Hiperaktivite

Q25 **Zor**

Başlık kullanım kolaylığı:

Q26 **Rahatsız**

Seanslar sırasında çocuğun rahatlık seviyesi:

Q27 **Bazen**

Tedavi programına uyum derecesi:

Q28 **İsteksiz**

Çocuğun Tedaviye Katılım İsteği:

Q29

Seanslar sırasında çocuğun rahat kalmasına yardımcı olmak için herhangi ek araçlar veya teknikler kullandınız mı? (örn. dikkat dağıtma yöntemleri, rahatlatıcı eşyalar):

Sakinleştirici müzik

Q30

Her seans sırasında başlığın doğru yerleştirilmesini ve kullanılmasını nasıl sağladınız:

Verilen öneriler ile

Q31 **Hayır**

Planlanan tedavi seanslarının tamamlanmasında herhangi bir kesinti veya zorluk yaşandı mı?

Page 3: Bölüm C: Gözlemlenen Etkiler

Q32 **Katılıyorum**

Sözlü iletişimde iyileşmeler fark edildi:

Q33

Gözlemlenen sözlü iletişim iyileşmelerine dair özel örnekler:

Daha fazla kelime tekrar yapma

Q34 **Biraz iyileşme**

Başlığın çocuğun konuşmaları başlatma yeteneği üzerindeki etkisi:

Q35 **Önemli derecede iyileşme**

Çocuğun jest ve yüz ifadeleri kullanımı üzerindeki değişiklikler:

Q36 **Katılmıyorum**

Davranışsal Değişiklikler: Sorunlu davranışlarda azalma gözlendi

Q37

Akranlar ve aile üyeleri ile sosyal etkileşimlerde ne gibi değişiklik gözlendi?

Göz teması

Q38 **Evet**

Yeni zorlayıcı davranışlar gözlendi mi?

Q39

Akranlarla sosyal etkileşimde iyileşme örnekleri:

İletişim

Q40

Aile üyeleri ile sosyal etkileşimde iyileşme örnekleri:

İletişim

Q41 **Biraz iyileşme**

Çocuğun paylaşma ve sırasını bekleme yeteneğindeki değişiklikler:

Q42 **Katılmıyorum**

Bilişsel ve Duygusal Değişiklikler:Dikkat süresi ve odaklanmada iyileşmeler görüldü

Q43 **Katılıyorum**

Duygusal değişiklikler görüldü

Q44

Gözlemlenen bilişsel değişiklik örnekleri:

Algıda iyikeşme

Q45

Gözlemlenen duygusal değişiklik örnekleri:

Yok

Q46 **Değişiklik yok**

Akademik performansta değişiklikler

Q47 **Biraz iyileşme**

Çocuğun görevleri tamamlama ve talimatları yerine getirme yeteneği üzerindeki etkisi:

Q48 **Önemli derecede iyileşme**

Çocuğun ruh hali ve genel mutluluğu üzerindeki değişiklikler:

Q49 **Respondent skipped this question**

Tedaviden kaynaklandığı gözlemlenen yan etkiler veya olumsuz sonuçlar:

Page 4: Bölüm D: Aile Üzerindeki Etki

Q50

Tedavi nedeniyle aile rutinlerinde ve dinamiklerinde değişiklikler:

Yok

Q51

Kardeş ilişkileri üzerindeki etkisi:

Birlikte oynama

Q52

Aile dinamiklerindeki değişikliklere dair özel örnekler:

İletişim

Q53

Geniş aile veya topluluktan alınan destek

İletişim

Q54 **Biraz iyileşme**

Aile gezi ve etkinlikleri üzerindeki etkisi

Q55 Aile iletişim kalıplarındaki değişiklikler:	**Biraz iyileşme**
Q56 Sağlık profesyonellerinden alınan destek	**Yok**
Q57 Diğer aile üyelerinin tedavi sürecine katılım düzeyi:	**Yüksek**
Q58 Aile uyumu ve bütünlüğü üzerindeki etkisi:	**Biraz iyileşme**
Q59 Ebeveyn Stresi ve İyilik Hali:Tedaviyi yönetmeyle ilgili stres düzeyi:	**Düşük**
Q60 Ebeveyn/Vasinin genel hali:	**İyileşti**
Q61 Ebeveynler için mental sağlık kaynaklarının erişilebilirliği:	**Kolay**
Q62 Tedavi maliyetlerinin mali istikrar üzerindeki etkisi:	**Önemli derecede etki**
Q63 Tedavi hakkında sağlanan bilgi ve kaynaklardan memnuniyet:	**Memnun**
Q64 Tedavi süreciyle ilgili stresi yönetmek için kullanılan stratejiler:	Respondent skipped this question

Page 5: Bölüm E: Memnuniyet ve Geri Bildirim

Q65 Genel tedavi deneyiminden memnuniyet:	**Çok memnun**

Q66

tDCS tedavisinin sonuçlarından memnuniyet:

Çok memnun

Q67

Bu tedaviyi diğer ailelere tavsiye eder misiniz?

Evet

Q68

Ek yorumlar veya geri bildirimler:

Respondent skipped this question

Q69

Sağlık profesyonellerinden alınan destekten memnuniyet:

Kararsız

Q70

Tedavi hakkında sağlanan bilgiden memnuniyet:

Memnun

Q71

Tedaviyi günlük rutinlere entegre etme kolaylığından memnuniyet:

Memnun

Q72

tDCS tedavisinin çocuğun yaşam kalitesi üzerindeki genel etkinliği:

Etkili

Q73

Diğer tedavilerle karşılaştırıldığında tDCS tedavisi:

Çok daha iyi

Page 6: Bölüm F: Açık Uçlu Sorular

Q74

tDCS tedavisine başladıktan sonra çocuğunuzda gözlemlediğiniz belirgin değişiklikleri tanımlayın:

Belirli becerilerdeki iyileşmeler (örn. iletişim, sosyal etkileşim, akademik performans):

Q75

Tedaviyi uygularken karşılaştığınız zorlukları ve bunları nasıl aştığınızı paylaşın:

Seanslar sırasında çocuğun direnci veya rahatsızlığı:

Q76

Tedavinin ailenizin günlük hayatını nasıl etkilediğine dair ek yorumlar:

Kardeş ilişkileri üzerindeki etkiler:

Q77

Çocuğunuzun tDCS tedavisine karşı tutumunu tanımlayın:

Zaman içinde tutum değişiklikleri:

Q78

Çocuğunuz için tDCS tedavisinin uzun vadeli beklentileri hakkında ne düşünüyorsunuz?

Geleceğe yönelik beklentiler:

Q79

tDCS tedavisinin çocuğunuzun Otizm yönetimine yaklaşımınızı nasıl etkilediğini anlatın:

Diğer tedaviler veya eğitim planları ile entegrasyon:

Q80

tDCS tedavi dönemi boyunca yaşanan unutulmaz anları veya kayda değer notları paylaşın:

Olumlu etkileşimler veya kilometre taşları:

Q81

Çocuğunuz için tDCS tedavisini düşünen diğer ebeveynlere yönelik öneriler:

Bir destek ağı oluşturmak:

Q82

Çocuğunuzun gelecekteki yaşantısında tDCS tedavisinin rolünü nasıl görüyorsunuz?

Sürekli kullanım planları:

Q83

Ek yorumlar veya yansımalar:

Respondent skipped this question

Appendix G



Message invite

Facebook post invite to private facebook group over 8300 members

Alptekin Aydin rTMS & tDCs Treatment

Private group · 8.3K members

+ Invite Share

Discussion Featured Members Media Files

Uzman Psikolog Alptekin Aydin

Admin All-star contributor · 11 June at 17:50 ·

Dear Families,
I am sending you a short survey link to participate in the scientific study we are conducting on tDCS (Headset) treatment in children with special needs. The results here are very important to us, It will be published in scientific papers and perhaps it will change the understanding and perspective of states towards this treatment.
It is very important for every family treated with tDCS (Headset) to fill out this short survey in the most sincere and accurate way, no personal information of you or your children will be shared.
Please click the link.https://www.surveymonkey.com/r/6S9X8ML
Please fill out our survey in the link above because your opinions are very valuable to us



SURVEYMONKEY.COM

DLPFC Üzerindeki tDCS Tedavisinin Özel Çocuklarda Sosyal İletişim Üzerindeki Etkisinin Değerlendirilmesi: Aile deneyimleri...

View insights 1.3K post reach